

THE UNIVERSITY OF TORONTO
UNDERGRADUATE MATHEMATICS COMPETITION

Sunday, March 18, 2001

Time: 3 hours

1. Let $a, b, c > 0$, $a < bc$ and $1 + a^3 = b^3 + c^3$. Prove that $1 + a < b + c$.
2. Let $O = (0, 0)$ and $Q = (1, 0)$. Find the point P on the line with equation $y = x + 1$ for which the angle OPQ is a maximum.
3. (a) Consider the infinite integer lattice in the plane (*i.e.*, the set of points with integer coordinates) as a graph, with the edges being the lines of unit length connecting nearby points. What is the minimum number of colours that can be used to colour all the vertices and edges of this graph, so that
 - (i) each pair of adjacent vertices gets two distinct colours;
 - (ii) each pair of edges that meet at a vertex gets two distinct colours; and
 - (iii) an edge is coloured differently than either of the two vertices at the ends?(b) Extend this result to lattices in real n -dimensional space.

4. Let \mathbf{V} be the vector space of all continuous real-valued functions defined on the open interval $(-\pi/2, \pi/2)$, with the sum of two functions and the product of a function and a real scalar defined in the usual way.

(a) Prove that the set $\{\sin x, \cos x, \tan x, \sec x\}$ is linearly independent.

(b) Let \mathbf{W} be the linear space generated by the four trigonometric functions given in (a), and let T be the linear transformation determined on \mathbf{W} into \mathbf{V} by $T(\sin x) = \sin^2 x$, $T(\cos x) = \cos^2 x$, $T(\tan x) = \tan^2 x$ and $T(\sec x) = \sec^2 x$. Determine a basis for the kernel of T .

Notes. A subset $\{v_1, v_2, \dots, v_k\}$ of a vector space is *linearly independent* iff $c_1v_1 + c_2v_2 + \dots + c_kv_k = 0$ for scalars c_i implies that $c_1 = c_2 = \dots = c_k = 0$. The *kernel* of a linear transformation is the subspace that T maps to the zero vector. A *basis* for a vector space is a linearly independent set of vectors for which every element of the space is some linear combination.

5. Let n be a positive integer and x a real number not equal to a nonnegative integer. Prove that

$$\frac{n}{x} + \frac{n(n-1)}{x(x-1)} + \frac{n(n-1)(n-2)}{x(x-1)(x-2)} + \dots + \frac{n(n-1)(n-2)\dots 1}{x(x-1)(x-2)\dots(x-n+1)} = \frac{n}{x-n+1}.$$

[This was a problem given by Samuel Beatty on a regular problem assignment to first year honours mathematics students in the 1930s.]

6. Prove that, for each positive integer n , the series

$$\sum_{k=1}^{\infty} \frac{k^n}{2^k}$$

converges to twice an odd integer not less than $(n+1)!$.

7. Suppose that $x \geq 1$ and that $x = [x] + \{x\}$, where $[x]$ is the greatest integer not exceeding x and the fractional part $\{x\}$ satisfies $0 \leq \{x\} < 1$. Define

$$f(x) = \frac{\sqrt{[x]} + \sqrt{\{x\}}}{\sqrt{x}}.$$

- (a) Determine the supremum, *i.e.*, the least upper bound, of the values of $f(x)$ for $1 \leq x$.
- (b) Let $x_0 \geq 1$ be given, and for $n \geq 1$, define $x_n = f(x_{n-1})$. Prove that $\lim_{n \rightarrow \infty} x_n$ exists.
8. A regular heptagon (polygon with seven equal sides and seven equal angles) has diagonals of two different lengths. Let a be the length of a side, b be the length of a shorter diagonal and c be the length of a longer diagonal of a regular heptagon (so that $a < b < c$). Prove ONE of the following relationships:

$$\frac{a^2}{b^2} + \frac{b^2}{c^2} + \frac{c^2}{a^2} = 6$$

or

$$\frac{b^2}{a^2} + \frac{c^2}{b^2} + \frac{a^2}{c^2} = 5 .$$

Saturday, March 9, 2002

Time: $3\frac{1}{2}$ hours

1. Let A, B, C be three pairwise orthogonal faces of a tetrahedron meeting at one of its vertices and having respective areas a, b, c . Let the face D opposite this vertex have area d . Prove that

$$d^2 = a^2 + b^2 + c^2 .$$

2. Angus likes to go to the movies. On Monday, standing in line, he noted that the fraction x of the line was in front of him, while $1/n$ of the line was behind him. On Tuesday, the same fraction x of the line was in front of him, while $1/(n+1)$ of the line was behind him. On Wednesday, the same fraction x of the line was in front of him, while $1/(n+2)$ of the line was behind him. Determine a value of n for which this is possible.
3. In how many ways can the rational $2002/2001$ be written as the product of two rationals of the form $(n+1)/n$, where n is a positive integer?
4. Consider the parabola of equation $y = x^2$. The normal is constructed at a variable point P and meets the parabola again in Q . Determine the location of P for which the arc length along the parabola between P and Q is minimized.
5. Let n be a positive integer. Suppose that f is a function defined and continuous on $[0, 1]$ that is differentiable on $(0, 1)$ and satisfies $f(0) = 0$ and $f(1) = 1$. Prove that, there exist n [distinct] numbers x_i ($1 \leq i \leq n$) in $(0, 1)$ for which

$$\sum_{i=1}^n \frac{1}{f'(x_i)} = n .$$

6. Let $x, y > 0$ be such that $x^3 + y^3 \leq x - y$. Prove that $x^2 + y^2 \leq 1$.
7. Prove that no vector space over \mathbb{R} is a finite union of proper subspaces.
8. (a) Suppose that P is an $n \times n$ nonsingular matrix and that u and v are column vectors with n components. The matrix $v^T P^{-1} u$ is 1×1 , and so can be identified with a scalar. Suppose that its value is not equal to -1 . Prove that the matrix $P + uv^T$ is nonsingular and that

$$(P + uv^T)^{-1} = P^{-1} - \frac{1}{\alpha} P^{-1} uv^T P^{-1}$$

where v^T denotes the transpose of v and $\alpha = 1 + v^T P^{-1} u$.

(b) Explain the situation when $\alpha = 0$.

9. A sequence whose entries are 0 and 1 has the property that, if each 0 is replaced by 01 and each 1 by 001, then the sequence remains unchanged. Thus, it starts out as 010010101001 \dots . What is the 2002th term of the sequence?

Sunday, March 16, 2003

Time: $3\frac{1}{2}$ hours

No aids or calculators permitted.

1. Evaluate

$$\sum_{n=1}^{\infty} \tan^{-1} \left(\frac{2}{n^2} \right).$$

[\tan^{-1} denotes the (composition) inverse function for \tan .]

2. Let a, b, c be positive real numbers for which $a + b + c = abc$. Prove that

$$\frac{1}{\sqrt{1+a^2}} + \frac{1}{\sqrt{1+b^2}} + \frac{1}{\sqrt{1+c^2}} \leq \frac{3}{2}.$$

3. Solve the differential equation

$$y'' = yy'.$$

4. Show that n divides the integer nearest to

$$\frac{(n+1)!}{e}.$$

5. For $x > 0$, $y > 0$, let $g(x, y)$ denote the minimum of the three quantities, x , $y + 1/x$ and $1/y$. Determine the maximum value of $g(x, y)$ and where this maximum is assumed.

6. A set of n lightbulbs, each with an *on-off* switch, numbered $1, 2, \dots, n$ are arranged in a line. All are initially off. Switch 1 can be operated at any time to turn its bulb on or off. Switch 2 can turn bulb 2 on or off if and only if bulb 1 is off; otherwise, it does not function. For $k \geq 3$, switch k can turn bulb k on or off if and only if bulb $k-1$ is off and bulbs $1, 2, \dots, k-2$ are all on; otherwise it does not function.

(a) Prove that there is an algorithm that will turn all of the bulbs on.

(b) If x_n is the length of the shortest algorithm that will turn on all n bulbs when they are initially off, determine the largest prime divisor of $3x_n + 1$ when n is odd.

7. Suppose that the polynomial $f(x)$ of degree $n \geq 1$ has all real roots and that $\lambda > 0$. Prove that the set $\{x \in \mathbb{R} : |f(x)| \leq \lambda |f'(x)|\}$ is a finite union of closed intervals whose total length is equal to $2n\lambda$.

8. Three matrices A , B and $A + B$ have rank 1. Prove that either all the rows of A and B are multiples of one and the same vector, or that all of the columns of A and B are multiples of one and the same vector.

9. Prove that the integral

$$\int_0^{\infty} \frac{\sin^2 x}{\pi^2 - x^2} dx$$

exists and evaluate it.

10. Let G be a finite group of order n . Show that n is odd if and only if each element of G is a square.

Sunday, March 14, 2004

Time: $3\frac{1}{2}$ hours

1. Prove that, for any complex numbers z and w ,

$$(|z| + |w|) \left| \frac{z}{|z|} + \frac{w}{|w|} \right| \leq 2|z + w| .$$

2. Prove that

$$\int_0^1 x^x dx = 1 - \frac{1}{2^2} + \frac{1}{3^3} - \frac{1}{4^4} + \frac{1}{5^5} + \cdots .$$

3. Suppose that u and v are positive integer divisors of the positive integer n and that $uv < n$. Is it necessarily so that the greatest common divisor of n/u and n/v exceeds 1?

4. Let n be a positive integer exceeding 1. How many permutations $\{a_1, a_2, \dots, a_n\}$ of $\{1, 2, \dots, n\}$ are there which maximize the value of the sum

$$|a_2 - a_1| + |a_3 - a_2| + \cdots + |a_{i+1} - a_i| + \cdots + |a_n - a_{n-1}|$$

over all permutations? What is the value of this maximum sum?

5. Let A be a $n \times n$ matrix with determinant equal to 1. Let B be the matrix obtained by adding 1 to every entry of A . Prove that the determinant of B is equal to $1 + s$, where s is the sum of the n^2 entries of A^{-1} .

6. Determine

$$\left(\int_0^1 \frac{dt}{\sqrt{1-t^4}} \right) \div \left(\int_0^1 \frac{dt}{\sqrt{1+t^4}} \right) .$$

7. Let a be a parameter. Define the sequence $\{f_n(x) : n = 0, 1, 2, \dots\}$ of polynomials by

$$f_0(x) \equiv 1$$

$$f_{n+1}(x) = xf_n(x) + f_n(ax)$$

for $n \geq 0$.

- (a) Prove that, for all n, x ,

$$f_n(x) = x^n f_n(1/x) .$$

- (b) Determine a formula for the coefficient of x^k ($0 \leq k \leq n$) in $f_n(x)$.

8. Let V be a complex n -dimensional inner product space. Prove that

$$|u|^2|v|^2 - \frac{1}{4}|u-v|^2|u+v|^2 \leq |(u, v)|^2 \leq |u|^2|v|^2 .$$

9. Let $ABCD$ be a convex quadrilateral for which all sides and diagonals have rational length and AC and BD intersect at P . Prove that AP, BP, CP, DP all have rational length.

Saturday, March 12, 2005

Time: $3\frac{1}{2}$ hours

1. Show that, if $-\pi/2 < \theta < \pi/2$, then

$$\int_0^\theta \log(1 + \tan \theta \tan x) dx = \theta \log \sec \theta .$$

2. Suppose that f is continuously differentiable on $[0, 1]$ and that $\int_0^1 f(x) dx = 0$. Prove that

$$2 \int_0^1 f(x)^2 dx \leq \int_0^1 |f'(x)| dx \cdot \int_0^1 |f(x)| dx .$$

3. How many $n \times n$ invertible matrices A are there for which all the entries of both A and A^{-1} are either 0 or 1?
4. Let a be a nonzero real and \mathbf{u} and \mathbf{v} be real 3-vectors. Solve the equation

$$2a\mathbf{x} + (\mathbf{v} \times \mathbf{x}) + \mathbf{u} = \mathbf{0}$$

for the vector \mathbf{x} .

5. Let $f(x)$ be a polynomial with real coefficients, evenly many of which are nonzero, which is *palindromic*. This means that the coefficients read the same in either direction, *i.e.* $a_k = a_{n-k}$ if $f(x) = \sum_{k=0}^n a_k x^k$, or, alternatively $f(x) = x^n f(1/x)$, where n is the degree of the polynomial. Prove that $f(x)$ has at least one root of absolute value 1.
6. Let G be a subgroup of index 2 contained in S_n , the group of all permutations of n elements. Prove that $G = A_n$, the alternating group of all even permutations.
7. Let $f(x)$ be a nonconstant polynomial that takes only integer values when x is an integer, and let P be the set of all primes that divide $f(m)$ for at least one integer m . Prove that P is an infinite set.
8. Let $AX = B$ represent a system of m linear equations in n unknowns, where $A = (a_{ij})$ is an $m \times n$ matrix, $X = (x_1, \dots, x_n)^t$ is an $n \times 1$ vector and $B = (b_1, \dots, b_m)^t$ is an $m \times 1$ vector. Suppose that there exists at least one solution for $AX = B$. Given $1 \leq j \leq n$, prove that the value of the j th component is the same for every solution X of $AX = B$ if and only if the rank of A is decreased if the j th column of A is removed.
9. Let S be the set of all real-valued functions that are defined, positive and twice continuously differentiable on a neighbourhood of 0. Suppose that a and b are real parameters with $ab \neq 0, b < 0$. Define operators from S to \mathbf{R} as follows:

$$A(f) = f(0) + af'(0) + bf''(0) ;$$

$$G(f) = \exp A(\log f) .$$

- (a) Prove that $A(f) \leq G(f)$ for $f \in S$;
- (b) Prove that $G(f + g) \leq G(f) + G(g)$ for $f, g \in S$;
- (c) Suppose that H is the set of functions in S for which $G(f) \leq f(0)$. Give examples of nonconstant functions, one in H and one not in H . Prove that, if $\lambda > 0$ and $f, g \in H$, then $\lambda f, f + g$ and fg all belong to H .

10. Let n be a positive integer exceeding 1. Prove that, if a graph with $2n + 1$ vertices has at least $3n + 1$ edges, then the graph contains a circuit (*i.e.*, a closed non-self-intersecting chain of edges whose terminal point is its initial point) with an even number of edges. Prove that this statement does not hold if the number of edges is only $3n$.

Sunday, March 12, 2006

Time: $3\frac{1}{2}$ hours

1. (a) Suppose that a 6×6 square grid of unit squares (chessboard) is tiled by 1×2 rectangles (dominoes). Prove that it can be decomposed into two rectangles, tiled by disjoint subsets of the dominoes.
 (b) Is the same thing true for an 8×8 array?
2. Let \mathbf{u} be a unit vector in \mathbb{R}^3 and define the operator P by $P(\mathbf{x}) = \mathbf{u} \times \mathbf{x}$ for $\mathbf{x} \in \mathbf{R}^3$ (where \times denotes the cross product).
 (a) Describe the operator $I + P^2$.
 (b) Describe the action of the operator $I + (\sin \theta)P + (1 - \cos \theta)P^2$.
3. Let $p(x)$ be a polynomial of positive degree n with n distinct real roots $a_1 < a_2 < \dots < a_n$. Let b be a real number for which $2b < a_1 + a_2$. Prove that

$$2^{n-1}|p(b)| \geq |p'(a_1)(b - a_1)|.$$

4. Two parabolas have parallel axes and intersect in two points. Prove that their common chord bisects the segments whose endpoints are the points of contact of their common tangent.
5. Suppose that you have a 3×3 grid of squares. A *line* is a set of three squares in the same row, the same column or the same diagonal; thus, there are eight lines.

Two players A and B play a game. They take alternate turns, A putting a 0 in any unoccupied square of the grid and B putting a 1. The first player is A , and the game cannot go on for more than nine moves. (The play is similar to noughts-and-crosses, or tictactoe.) A move is *legitimate* if it does not result in two lines of squares being filled in with different sums. The winner is the last player to make a legitimate move.

(For example, if there are three 0s down the diagonal, then B can place a 1 in any vacant square provided it completes no other line, for then the sum would differ from the diagonal sum. If there are two zeros at the top of the main diagonal and two ones at the left of the bottom line, then the lower right square cannot be filled by either player, as it would result in two lines with different sums.)

- (a) What is the maximum number of legitimate moves possible in a game?
 - (b) What is the minimum number of legitimate moves possible in a game that would not leave a legitimate move available for the next player?
 - (c) Which player has a winning strategy? Explain.
6. Suppose that k is a positive integer and that

$$f(t) = a_1 e^{\lambda_1 t} + a_2 e^{\lambda_2 t} + \dots + a_k e^{\lambda_k t}$$

where $a_1, \dots, a_k, \lambda_1, \dots, \lambda_k$ are real numbers with $\lambda_1 < \lambda_2 < \dots < \lambda_k$. Prove that $f(t) = 0$ has finitely many real solutions. What is the maximum number of solutions possible, as a function of k ?

7. Let A be a real 3×3 invertible matrix for which the sums of the rows, columns and two diagonals are all equal. Prove that the rows, columns and diagonal sums of A^{-1} are all equal.
8. Let $f(x)$ be a real function defined and twice differentiable on an open interval containing $[-1, 1]$. Suppose that $0 < \alpha \leq \gamma$ and that $|f(x)| \leq \alpha$ and $|f''(x)| \leq \gamma$ for $-1 \leq x \leq 1$. Prove that

$$|f'(x)| \leq 2\sqrt{\alpha\gamma}$$

for $-1 \leq x \leq 1$. (Part marks are possible for the weaker inequality $|f'(x)| \leq \alpha + \gamma$.)

9. A high school student asked to solve the surd equation

$$\sqrt{3x-2} - \sqrt{2x-3} = 1$$

gave the following answer: *Squaring both sides leads to*

$$3x - 2 - 2x - 3 = 1$$

so $x = 6$. The answer is, in fact, correct.

Show that there are infinitely many real quadruples (a, b, c, d) for which this method leads to a correct solution of the surd equation

$$\sqrt{ax-b} - \sqrt{cx-d} = 1.$$

10. Let P be a planar polygon that is not convex. The vertices can be classified as either *convex* or *concave* according as to whether the angle at the vertex is less than or greater than 180° respectively. There must be at least two convex vertices. Select two consecutive convex vertices (*i.e.*, two interior angles less than 180° for which all interior angles in between exceed 180°) and join them by a segment. Reflect the edges between these two convex angles in the segment to form along with the other edges of P a polygon P_1 . If P_1 is not convex, repeat the process, reflecting some of the edges of P_1 in a segment joining two consecutive convex vertices, to form a polygon P_2 . Repeat the process. Prove that, after a finite number of steps, we arrive at a polygon P_n that is convex.

Sunday, March 11, 2007

Time: $3\frac{1}{2}$ hours

1. A $m \times n$ rectangular array of distinct real numbers has the property that the numbers in each row increase from left to right. The entries in each column, individually, are rearranged so that the numbers in each column increase from top to bottom. Prove that in the final array, the numbers in each row will increase from left to right.
2. Determine distinct positive integers a, b, c, d, e such that the five numbers a, b^2, c^3, d^4, e^5 constitute an arithmetic progression. (The difference between adjacent pairs is the same.)
3. Prove that the set $\{1, 2, \dots, n\}$ can be partitioned into k subsets with the same sum if and only if k divides $\frac{1}{2}n(n+1)$ and $n \geq 2k-1$.
4. Suppose that $f(x)$ is a continuous real-valued function defined on the interval $[0, 1]$ that is twice differentiable on $(0, 1)$ and satisfies (i) $f(0) = 0$ and (ii) $f''(x) > 0$ for $0 < x < 1$.
 - (a) Prove that there exists a number a for which $0 < a < 1$ and $f'(a) < f(1)$;
 - (b) Prove that there exists a unique number b for which $a < b < 1$ and $f'(a) = f(b)/b$.

5. For $x \leq 1$ and $x \neq 0$, let

$$f(x) = \frac{-8[1 - (1 - x)^{1/2}]^3}{x^2}.$$

- (a) Prove that $\lim_{x \rightarrow 0} f(x)$ exists. Take this as the value of $f(0)$.
(b) Determine the smallest closed interval that contains the set of all values assumed by $f(x)$ on its domain.
(c) Prove that $f(f(f(x))) = f(x)$ for all $x \leq 1$.
6. Let $h(n)$ denote the number of finite sequences $\{a_1, a_2, \dots, a_k\}$ of positive integers exceeding 1 for which $k \geq 1$, $a_1 \geq a_2 \geq \dots \geq a_k$ and $n = a_1 a_2 \dots a_k$. (For example, if $n = 20$, there are four such sequences $\{20\}$, $\{10, 2\}$, $\{5, 4\}$ and $\{5, 2, 2\}$ and $h(20) = 4$.)

Prove that

$$\sum_{n=1}^{\infty} \frac{h(n)}{n^2} = 1.$$

7. Find the Jordan canonical form of the matrix $\mathbf{u}\mathbf{v}^t$ where \mathbf{u} and \mathbf{v} are column vectors in \mathbf{C}^n . (The superscript t denotes the transpose.)
8. Suppose that n points are given in the plane, not all collinear. Prove that there are at least n distinct straight lines that can be drawn through pairs of the points.
9. Which integers can be written in the form

$$\frac{(x + y + z)^2}{xyz}$$

where x, y, z are positive integers?

10. Solve the following differential equation

$$2y' = 3|y|^{1/3}$$

subject to the initial conditions

$$y(-2) = -1 \quad \text{and} \quad y(3) = 1.$$

Your solution should be everywhere differentiable.

Sunday, March 9, 2008

Time: $3\frac{1}{2}$ hours

1. Three angles of a heptagon (7-sided polygon) inscribed in a circle are equal to 120° . Prove that at least two of its sides are equal.
2. (a) Determine a real-valued function g defined on the real numbers that is decreasing and for which $g(g(x)) = 2x + 2$.
(b) Prove that there is no real-valued function f defined on the real numbers that is decreasing and for which $f(f(x)) = x + 1$.
3. Suppose that a is a real number and the sequence $\{a_n\}$ is defined recursively by $a_0 = a$ and

$$a_{n+1} = a_n(a_n - 1)$$

for $n \geq 0$. Find the values of a for which the sequence $\{a_n\}$ converges.

4. Suppose that u, v, w, z are complex numbers for which $u + v + w + z = u^2 + v^2 + w^2 + z^2 = 0$. Prove that

$$(u^4 + v^4 + w^4 + z^4)^2 = 4(u^8 + v^8 + w^8 + z^8).$$

5. Suppose that $a, b, c \in \mathbf{C}$ with $ab = 1$. Evaluate the determinant of

$$\begin{pmatrix} c & a & a^2 & \cdots & a^{n-1} \\ b & c & a & \cdots & a^{n-2} \\ b^2 & b & c & \cdots & a^{n-3} \\ \vdots & \vdots & \vdots & & \vdots \\ b^{n-1} & b^{n-2} & \cdots & & c \end{pmatrix}$$

6. 2008 circular coins, possibly of different diameters, are placed on the surface of a flat table in such a way that no coin is on top of another coin. What is the largest number of points at which two of the coins could be touching?
7. Let G be a group of finite order and identity e . Suppose that ϕ is an automorphism of G onto itself with the following properties: (1) $\phi(x) = x$ if and only if $x = e$; (2) $\phi(\phi(x)) = x$ for each element x of G . (The mapping ϕ has the property that it is one-one onto and that $\phi(xy) = \phi(x)\phi(y)$ for each pair x, y of elements of G .)
- (a) Give an example of a group and automorphism for which these conditions are satisfied.
- (b) Prove that G is commutative (*i.e.*, $xy = yx$ for each pair x, y of elements in G).
8. Let $b \geq 2$ be an integer base of numeration and let $1 \leq r \leq b - 1$. Determine the sum of all r -digit numbers of the form

$$\overline{a_{r-1}a_{r-2}\cdots a_2a_1a_0} \equiv a_{r-1}b^{r-1} + a_{r-2}b^{r-2} + \cdots + a_1b + a_0$$

whose digits increase strictly from left to right: $1 \leq a_{r-1} < a_{r-2} < \cdots < a_1 < a_0 \leq b - 1$.

9. For each positive integer n , let

$$S(n) = \sum_{k=1}^n \frac{2^k}{k^2}.$$

Prove that $S(n+1)/S(n)$ is not a rational function of n . [A *rational function* is one that can be written as a ratio of two polynomials.]

10. A point is chosen at random (with the uniform distribution) on each side of a unit square. What is the probability that the four points are the vertices of a quadrilateral with area exceeding $\frac{1}{2}$?

Sunday, March 8, 2009

Time: $3\frac{1}{2}$ hours

1. Determine the supremum and the infimum of

$$\frac{(x-1)^{x-1}x^x}{(x-(1/2))^{2x-1}}$$

for $x > 1$.

- Let n and k be integers with $n \geq 0$ and $k \geq 1$. Let $\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_n$ be $n + 1$ distinct points in \mathbf{R}^k and let y_0, y_1, \dots, y_n be $n + 1$ real numbers (not necessarily distinct). Prove that there exists a polynomial p of degree at most n in the coordinates of \mathbf{x} with respect to the standard basis for which $p(\mathbf{x}_i) = y_i$ for $0 \leq i \leq n$.
- For each positive integer n , let $p(n)$ be the product of all positive integral divisors of n . Is it possible to find two distinct positive integers m and n for which $p(m) = p(n)$?
- Let $\{a_n\}$ be a real sequence for which

$$\sum_{n=1}^{\infty} \frac{a_n}{n}$$

converges. Prove that

$$\lim_{n \rightarrow \infty} \frac{a_1 + a_2 + \dots + a_n}{n} = 0 .$$

- Find a 3×3 matrix A with elements in \mathbb{Z}_2 for which $A^7 = I$ and $A \neq I$. (Here, I is the identity matrix and \mathbb{Z}_2 is the field of two elements 0 and 1 where addition and multiplication are defined modulo 2.)
- Determine all solutions in nonnegative integers (x, y, z, w) to the equation

$$2^x 3^y - 5^z 7^w = 1 .$$

- Let $n \geq 2$. Minimize $a_1 + a_2 + \dots + a_n$ subject to the constraints $0 \leq a_1 \leq a_2 \leq \dots \leq a_n$ and $a_1 a_2 + a_2 a_3 + \dots + a_{n-1} a_n + a_n a_1 = 1$. (When $n = 2$, the latter condition is $a_1 a_2 = 1$; when $n \geq 3$, the sum on the left has exactly n terms.)
- Let a, b, c be members of a real inner-product space (V, \langle, \rangle) whose norm is given by $\|x\|^2 = \langle x, x \rangle$. (You may assume that V is \mathbf{R}^n if you wish. Prove that

$$\|a + b\| + \|b + c\| + \|c + a\| \leq \|a\| + \|b\| + \|c\| + \|a + b + c\|$$

for $a, b, c, \in V$.

- Let p be a prime congruent to 1 modulo 4. For each real number x , let $\{x\} = x - [x]$ denote the fractional part of x . Determine

$$\sum \left\{ \left\{ \frac{k^2}{p} \right\} : 1 \leq k \leq \frac{1}{2}(p-1) \right\} .$$

- Suppose that a path on a $m \times n$ grid consisting of the lattice points $\{(x, y) : 1 \leq x \leq m, 1 \leq y \leq n\}$ (x and y both integers) consisting of $mn - 1$ unit segments begins at the point $(1, 1)$, passes through each point of the grid exactly once, does not intersect itself and finishes at the point (m, n) . Show that the path partitions the rectangle bounded by the lines $x = 1, x = m, y = 1, y = n$ into two subsets of equal area, the first consisting of regions opening to the left or up, and the second consisting of regions opening to the right or down.

Sunday, March 7, 2010

Time: $3\frac{1}{2}$ hours

vskip 0.5cm

- Let F_1 and F_2 be the foci of an ellipse and P be a point in the plane of the ellipse. Suppose that G_1 and G_2 are points on the ellipse for which PG_1 and PG_2 are tangents to the ellipse. Prove that $\angle F_1 P G_1 = \angle F_2 P G_2$.

2. Let $u_0 = 1$, $u_1 = 2$ and $u_{n+1} = 2u_n + u_{n-1}$ for $n \geq 1$. Prove that, for every nonnegative integer n ,

$$u_n = \sum \left\{ \frac{(i+j+k)!}{i!j!k!} : i, j, k \geq 0, i+j+2k = n \right\}.$$

3. Let \mathbf{a} and \mathbf{b} , the latter nonzero, be vectors in \mathbb{R}^3 . Determine the value of λ for which the vector equation

$$\mathbf{a} - (\mathbf{x} \times \mathbf{b}) = \lambda \mathbf{b}$$

is solvable, and then solve it.

4. The plane is partitioned into n regions by three families of parallel lines. What is the least number of lines to ensure that $n \geq 2010$?
5. Let m be a natural number, and let c, a_1, a_2, \dots, a_m be complex numbers for which $|a_i| = 1$ for $i = 1, 2, \dots, m$. Suppose also that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^m a_i^n = c.$$

Prove that $c = m$ and that $a_i = 1$ for $i = 1, 2, \dots, m$.

6. Let $f(x)$ be a quadratic polynomial. Prove that there exist quadratic polynomials $g(x)$ and $h(x)$ for which

$$f(x)f(x+1) = g(h(x)),$$

7. Suppose that f is a continuous real-valued function defined on the closed interval $[0, 1]$ and that

$$\left(\int_0^1 xf(x)dx \right)^2 = \left(\int_0^1 f(x)dx \right) \left(\int_0^1 x^2 f(x)dx \right).$$

Prove that there is a point $c \in (0, 1)$ for which $f(c) = 0$.

8. Let A be an invertible symmetric $n \times n$ matrix with entries $\{a_{i,j}\}$ in \mathbb{Z}_2 . Prove that there is an $n \times n$ matrix with entries in \mathbb{Z}_2 such that $A = M^t M$ only if $a_{i,i} \neq 0$ for some i .

[\mathbb{Z}_2 refers to the field of integers modulo 2 with two elements 0, 1 for which $1 + 1 = 0$. M^t refers to the transpose of the matrix M .]

9. Let f be a real-valued functions defined on \mathbb{R} with a continuous third derivative, let $S_0 = \{x : f(x) = 0\}$, and, for $k = 1, 2, 3$, $S_k = \{x : f^{(k)}(x) = 0\}$, where $f^{(k)}$ denotes the k th derivative of f . Suppose also that $\mathbb{R} = S_0 \cup S_1 \cup S_2 \cup S_3$. Must f be a polynomial of degree not exceeding 2?
10. Prove that the set \mathbb{Q} of rationals can be written as the union of countably many subsets of \mathbb{Q} each of which is dense in the set \mathbb{R} of real numbers.

Sunday, March 6, 2011

Time: $3\frac{1}{2}$ hours

1. Let S be a nonvoid set of real numbers with the property that, for each real number x , there is a unique real number $f(x)$ belonging to S that is farthest from x , i.e., for each y in S distinct from $f(x)$, $|x - f(x)| > |x - y|$. Prove that S must be a singleton.
2. Let u and v be positive reals. Minimize the larger of the two values

$$2u + \frac{1}{v^2} \quad \text{and} \quad 2v + \frac{1}{u^2}.$$

3. Suppose that S is a set of n nonzero real numbers such that exactly p of them are positive and exactly q are negative. Determine all the pairs (n, p) such that exactly half of the threefold products abc of distinct elements a, b, c of S are positive.
4. Let $\{b_n : n \geq 1\}$ be a sequence of positive real numbers such that

$$3b_{n+2} \geq b_{n+1} + 2b_n$$

for every positive integer n . Prove that either the sequence converges or that it diverges to infinity.

5. Solve the system

$$x + xy + xyz = 12$$

$$y + yz + yzx = 21$$

$$z + zx + zxy = 30 .$$

6. Two competitors play badminton. They play two games, each winning one of them. They then play a third game to determine the overall winner of the match. The winner of a game of badminton is the first player to score at least 21 points with a lead of at least 2 points over the other player.

In this particular match, it is observed that the scores of each player listed in order of the games form an arithmetic progression with a nonzero common difference. What are the scores of the two players in the third game?

7. Suppose that there are 2011 students in a school and that each student has a certain number of friends among his schoolmates. It is assumed that if A is a friend of B , then B is a friend of A , and also that there may exist certain pairs that are not friends. Prove that there is a nonvoid subset S of these students for which every student in the school has an even number of friends in S .
8. The set of transpositions of the symmetric group S_5 on $\{1, 2, 3, 4, 5\}$ is

$$\{(12), (13), (14), (15), (23), (24), (25), (34), (35), (45)\}$$

where (ab) denotes the permutation that interchanges a and b and leaves every other element fixed. Determine a product of all transpositions, each occurring exactly once, that is equal to the identity permutation ϵ , which leaves every element fixed.

9. Suppose that A and B are two square matrices of the same order for which the indicated inverses exist. Prove that

$$(A + AB^{-1}A)^{-1} + (A + B)^{-1} = A^{-1} .$$

10. Suppose that p is an odd prime. Determine the number of subsets S contained in $\{1, 2, \dots, 2p - 1, 2p\}$ for which (a) S has exactly p elements, and (b) the sum of the elements of S is a multiple of p .

Saturday, March 10, 2012

Time: $3\frac{1}{2}$ hours

1. An equilateral triangle of side length 1 can be covered by five equilateral triangles of side length u . Prove that it can be covered by four equilateral triangles of side length u . (A triangle is a closed convex set that contains its three sides along with its interior.)
2. Suppose that f is a function defined on the set Z of integers that takes integer values and satisfies the condition that $f(b) - f(a)$ is a multiple of $b - a$ for every pair a, b , of integers. Suppose also that p is a polynomial with integer coefficients such that $p(n) = f(n)$ for infinitely many integers n . Prove that $p(x) = f(x)$ for every positive integer x .

3. Given the real numbers a, b, c not all zero, determine the real solutions x, y, z, u, v, w for the system of equations:

$$x^2 + v^2 + w^2 = a^2$$

$$u^2 + y^2 + w^2 = b^2$$

$$u^2 + v^2 + z^2 = c^2$$

$$u(y + z) + vw = bc$$

$$v(x + z) + wu = ca$$

$$w(x + y) + uv = ab.$$

4. (a) Let n and k be positive integers. Prove that the least common multiple of $\{n, n + 1, n + 2, \dots, n + k\}$ is equal to

$$rn \binom{n+k}{k}$$

for some positive integer r .

(b) For each positive integer k , prove that there exist infinitely many positive integers n , for which the number r defined in part (a) is equal to 1.

5. Let \mathfrak{C} be a circle and Q a point in the plane. Determine the locus of the centres of those circles that are tangent to \mathfrak{C} and whose circumference passes through Q .
6. Find all continuous real-valued functions defined on \mathbb{R} that satisfy $f(0) = 0$ and

$$f(x) - f(y) = (x - y)g(x + y)$$

for some real valued function $g(x)$.

7. Consider the following problem:

Suppose that $f(x)$ is a continuous real-valued function defined on the interval $[0, 2]$ for which

$$\int_0^2 f(x) dx = \int_0^2 (f(x))^2 dx .$$

Prove that there exists a number $c \in [0, 2]$ for which either $f(c) = 0$ or $f(c) = 1$.

(a) Criticize the following solution:

Solution. Clearly $\int_0^2 f(x) dx \geq 0$. By the extreme value theorem, there exist numbers u and v in $[0, 2]$ for which $f(u) \leq f(x) \leq f(v)$ for $0 \leq x \leq 2$. Hence

$$f(u) \int_0^2 f(x) dx \leq \int_0^2 f(x)^2 dx \leq f(v) \int_0^2 f(x) dx .$$

Since $\int_0^2 f(x)^2 dx = 1 \cdot \int_0^2 f(x) dx$, by the intermediate value theorem, there exists a number $c \in [0, 2]$ for which $f(c) = 1$. \square

(b) Show that there is a nontrivial function f that satisfies the conditions of the problem but that never assumes the value 1.

(c) Provide a complete solution of the problem.

8. Determine the area of the set of points (x, y) in the plane that satisfy the two inequalities:

$$\begin{aligned} x^2 + y^2 &\leq 2 \\ x^4 + x^3 y^3 &\leq xy + y^4. \end{aligned}$$

9. In a round-robin tournament of $n \geq 2$ teams, each pair of teams plays exactly one game that results in a win for one team and a loss for the other (there are no ties).
- (a) Prove that the teams can be labelled t_1, t_2, \dots, t_n , so that, for each i with $1 \leq i \leq n - 1$, team t_i beats t_{i+1} .
- (b) Suppose that a team t has the property that, for each other team u , one can find a chain u_1, u_2, \dots, u_m of (possibly zero) distinct teams for which t beats u_1 , u_i beats u_{i+1} for $1 \leq i \leq m - 1$ and u_m beats u . Prove that *all* of the n teams can be ordered as in (a) so that $t = t_1$ and each t_i beats t_{i+1} for $1 \leq i \leq n - 1$.
- (c) Let T denote the set of teams who can be labelled as t_1 in an ordering of teams as in (a). Prove that, in any ordering of teams as in (a), all the teams in T occur before all the teams that are not in T .
10. Let A be a square matrix whose entries are complex numbers. Prove that $A^* = A$ if and only if $AA^* = A^2$.

Notes. For any $m \times n$ matrix M with entries m_{ij} , the *hermitian transpose* M^* is the $n \times m$ matrix M^* obtained by taking the complex conjugates of entries of M and transposing; thus, the (i, j) th element of M^* is $\overline{m_{ji}}$. In particular, for the complex column vector x with i th entry x_i , x^* is a row vector whose i th entry is $\overline{x_i}$. The inner product $\langle x, y \rangle$ of two column vectors is $\sum \overline{x_i}y_i = x^*y$, and we have that $\langle x, Ay \rangle = \langle A^*x, y \rangle$. A matrix for which $A^* = A$ is said to be *hermitian*.

Saturday, March 9, 2013

Time: $3\frac{1}{2}$ hours

1. (a) Let a be an odd positive integer exceeding 3, and let n be a positive integer. Prove that

$$a^{2^n} - 1$$

has at least $n + 1$ distinct prime divisors.

- (b) When $a = 3$, determine all the positive integers n for which the assertion in (a) is false.

2. ABCD is a square; points U and V are situated on the respective sides BC and CD . Prove that the perimeter of triangle CUV is equal to twice the sidelength of the square if and only if $\angle UAV = 45^\circ$.
3. Let $f(x)$ be a convex increasing realvalued function defined on the closed interval $[0, 1]$ for which $f(0) = 0$ and $f(1) = 1$. Suppose that $0 < a < 1$ and that $b = f(a)$.

- (a) Prove that f is continuous on $(0, 1)$.

- (b) Prove that

$$0 \leq a - b \leq 2 \int_0^1 (x - f(x))dx \leq 1 - 4b(1 - a).$$

Notes. $f(x)$ is *increasing* if and only if $f(u) \leq f(v)$ whenever $u \leq v$, and is *convex* if and only if

$$f((1 - t)u + tv) \leq (1 - t)f(u) + tf(v)$$

whenever $0 < t < 1$.

4. Let S be the set of integers of the form $x^2 + xy + y^2$, where x and y are integers.
- (a) Prove that any prime p in S is either equal to 3 or is congruent to 1 modulo 6.
- (b) Prove that S includes all squares.
- (c) Prove that S is closed under multiplication.

5. A point on an ellipse is joined to the ends of its major axis. Prove that the portion of a directrix intercepted by the two joining lines subtends a right angle at the corresponding focus.

Notes. The *directrix* corresponding to a focus F of an ellipse is a line with the property that, for any point P on the ellipse, the distance from P to F divided by the distance from P to the directrix is a constant e , called the *eccentricity*, less than 1. The major axis is the chord of the ellipse that passes through the two foci.

6. Let $p(x) = x^4 + ax^3 + bx^2 + cx + d$ be a polynomial with rational coefficients. Suppose that $p(x)$ has exactly one real zero r . Prove that r is rational.
7. Let $(V, \langle \cdot, \cdot \rangle)$ be a two-dimensional inner product space over the complex field \mathbf{C} and let z_1 and z_2 be unit vectors in V . Prove that

$$\sup\{|\langle z, z_1 \rangle \langle z, z_2 \rangle| : \|z\| = 1\} \geq \frac{1}{2}$$

with equality if and only if $\langle z_1, z_2 \rangle = 0$.

Note: The inner product $\langle z, w \rangle$ is linear in the left variable and satisfies $\langle w, z \rangle = \overline{\langle z, w \rangle}$. Also, $\|z\|^2 = \langle z, z \rangle$.

8. For any real square matrix A , the adjugate matrix, $\text{adj } A$, has as its elements that cofactors of the transpose of A , so that

$$A \cdot \text{adj } A = \text{adj } A \cdot A = (\det A)I .$$

- (a) Suppose that A is an invertible square matrix. Show that

$$(\text{adj } (A^t))^{-1} = (\text{adj } (A^{-1}))^t .$$

- (b) Suppose that $\text{adj } (A^t)$ is orthogonal (*i.e.*, its inverse is its transpose). Prove that A is invertible.

- (c) Let A be an invertible $n \times n$ square matrix and let $\det (tI - A) = t^n + c_1 t^{n-1} + \cdots + c_{n-1} t + c_n$ be the characteristic polynomial of the matrix A . Determine the characteristic polynomial of $\text{adj } A$.

Note. A real square matrix M is *orthogonal* if and only if the product of M and its transpose M^t is the identity matrix.

9. Let S be a set upon whose elements there is a binary operation $(x, y) \rightarrow xy$ which is associative (*i.e.* $x(yz) = (xy)z$). Suppose that there exists an element $e \in S$ for which $e^2 = e$ and that for each $a \in S$, there is at least one element b for which $ba = e$ and at most one element c for which $ac = e$. Prove that S is a group with this binary operation.

Note. A group G is a set with an associative binary operation that contains an identity element u for which, given any element $x \in G$, $xu = ux = x$ and there exists $y \in G$ for which $yx = xy = u$.

10. (a) Let f be a real-valued function defined on the real number field \mathbb{R} for which $|f(x) - f(y)| < |x - y|$ for any pair (x, y) of distinct elements of \mathbb{R} . Let $f^{(n)}$ denote the n th composite of f defined by $f^{(1)}(x) = f(x)$ and $f^{(n+1)}(x) = f(f^{(n)}(x))$ for $n \geq 2$. Prove that exactly one of the following situations must occur:
- (i) $\lim_{n \rightarrow +\infty} f^{(n)}(x) = +\infty$ for each real x ;
 - (ii) $\lim_{n \rightarrow +\infty} f^{(n)}(x) = -\infty$ for each real x ;
 - (iii) there is a real number z such that

$$\lim_{n \rightarrow +\infty} f^{(n)}(x) = z$$

for each real x .

- (b) Give examples to show that each of the three cases in (a) can occur.

UNDERGRADUATE MATHEMATICS COMPETITION

Sunday, March 9, 2014

Time: $3\frac{1}{2}$ hours

1. The *permanent*, $\text{per } A$, of a $n \times n$ matrix $A = (a_{i,j})$, is equal to the sum of all possible products of the form $a_{1,\sigma(1)}a_{2,\sigma(2)} \cdots a_{n,\sigma(n)}$, where σ runs over all the permutations on the set $\{1, 2, \dots, n\}$. (This is similar to the definition of determinant, but there is no sign factor.) Show that, for any $n \times n$ matrix $A = (a_{i,j})$ with positive real terms,

$$\text{per } A \geq n! \left(\prod_{1 \leq i, j \leq n} a_{i,j} \right)^{\frac{1}{n}}.$$

2. For a positive integer N written in base 10 numeration, N' denotes the integer with the digits of N written in reverse order. There are pairs of integers (A, B) for which A, A', B, B' are all distinct and $A \times B = B' \times A'$. For example,

$$3516 \times 8274 = 4728 \times 6153.$$

- (a) Determine a pair (A, B) as described above for which both A and B have two digits, and all four digits involved are distinct.
- (b) Are there any pairs (A, B) as described above for which A has two and B has three digits?
3. Let n be a positive integer. A finite sequence $\{a_1, a_2, \dots, a_n\}$ of positive integers a_i is said to be *tight* if and only if $1 \leq a_1 < a_2 < \dots < a_n$, all $\binom{n}{2}$ differences $a_j - a_i$ with $i < j$ are distinct, and a_n is as small as possible.
- (a) Determine a tight sequence for $n = 5$.
- (b) Prove that there is a polynomial $p(n)$ of degree not exceeding 3 such that $a_n \leq p(n)$ for every tight sequence $\{a_i\}$ with n entries.

4. Let $f(x)$ be a continuous realvalued function on $[0, 1]$ for which

$$\int_0^1 f(x)dx = 0 \quad \text{and} \quad \int_0^1 xf(x)dx = 1.$$

- (a) Give an example of such a function.
- (b) Prove that there is a nontrivial open interval I contained in $(0, 1)$ for which $|f(x)| > 4$ for $x \in I$.
5. Let n be a positive integer. Prove that

$$\sum_{k=1}^n \frac{1}{k \binom{n}{k}} = \sum_{k=1}^n \frac{1}{k 2^{n-k}} = \frac{1}{2^{n-1}} \sum_{k=1}^n \frac{2^{k-1}}{k} = \frac{1}{2^{n-1}} \sum \left\{ \frac{\binom{n}{k}}{k} : k \text{ odd}, 1 \leq k \leq n \right\}.$$

6. Let $f(x) = x^6 - x^4 + 2x^3 - x^2 + 1$.
- (a) Prove that $f(x)$ has no positive real roots.
- (b) Determine a nonzero polynomial $g(x)$ of minimum degree for which all the coefficients of $f(x)g(x)$ are nonnegative rational numbers.
- (c) Determine a polynomial $h(x)$ of minimum degree for which all the coefficients of $f(x)h(x)$ are positive rational numbers.

7. Suppose that x_0, x_1, \dots, x_n are real numbers. For $0 \leq i \leq n$, define

$$y_i = \max(x_0, x_1, \dots, x_i).$$

Prove that

$$y_n^2 \leq 4x_n^2 - 4 \sum_{i=0}^{n-1} y_i(x_{i+1} - x_i).$$

When does equality occur?

8. The hyperbola with equation $x^2 - y^2 = 1$ has two branches, as does the hyperbola with equation $y^2 - x^2 = 1$. Choose one point from each of the four branches of the locus of $(x^2 - y^2)^2 = 1$ such that area of the quadrilateral with these four vertices is minimized.
9. Let $\{a_n\}$ and $\{b_n\}$ be positive real sequences such that

$$\lim_{n \rightarrow \infty} \frac{a_n}{n} = u > 0$$

and

$$\lim_{n \rightarrow \infty} \left(\frac{b_n}{a_n} \right)^n = v > 0.$$

Prove that

$$\lim_{n \rightarrow \infty} \left(\frac{b_n}{a_n} \right) = 1$$

and

$$\lim_{n \rightarrow \infty} (b_n - a_n) = u \log v.$$

10. Does there exist a continuous realvalued function defined on \mathbf{R} for which $f(f(x)) = -x$ for all $x \in \mathbf{R}$?

UNDERGRADUATE MATHEMATICS COMPETITION

March 8, 2015

Time: $3\frac{1}{2}$ hours

1. Suppose that u and v are two real-valued functions defined on the set of reals. Let $f(x) = u(v(x))$ and $g(x) = u(-v(x))$ for each real x . If $f(x)$ is continuous, must $g(x)$ also be continuous?
2. Given $2n$ distinct points in space, the sum S of the lengths of all the segments joining pairs of them is calculated. Then n of the points are removed along with all the segments having at least one endpoint from among them. Prove that the sum of the lengths of all the remaining segments is less than $\frac{1}{2}S$.
3. Let $f : [0, 1] \rightarrow \mathbf{R}$ be continuously differentiable. Prove that

$$\left| \frac{f(0) + f(1)}{2} - \int_0^1 f(x) dx \right| \leq \frac{1}{4} \sup\{|f'(x)| : 0 \leq x \leq 1\}.$$

4. Determine all the values of the positive integer $n \geq 2$ for which the following statement is true, and for each, indicate when equality holds.

For any nonnegative real numbers x_1, x_2, \dots, x_n ,

$$(x_1 + x_2 + \dots + x_n)^2 \geq n(x_1x_2 + x_2x_3 + \dots + x_{n-1}x_n + x_nx_1),$$

where the right side has n summands.

5. Let $f(x)$ be a real polynomial of degree 4 whose graph has two real inflection points. There are three regions bounded by the graph and the line passing through these inflection points. Prove that two of these regions have equal area and that the area of the third region is equal to the sum of the other two areas.
6. Using the digits 1, 2, 3, 4, 5, 6, 7, 8, each exactly once, create two numbers and form their product. For example, $472 \times 83156 = 39249632$. What are the smallest and the largest values such a product can have?

7. Determine

$$\int_0^2 \frac{e^x dx}{e^{1-x} + e^{x-1}}.$$

8. Let $\{a_n\}$ and $\{b_n\}$ be two *decreasing* positive real sequences for which

$$\sum_{n=1}^{\infty} a_n = \infty$$

and

$$\sum_{n=1}^{\infty} b_n = \infty.$$

Let I be a subset of the natural numbers, and define the sequence $\{c_n\}$ by

$$c_n = \begin{cases} a_n, & \text{if } n \in I \\ b_n, & \text{if } n \notin I. \end{cases}$$

Is it possible for $\sum_{n=1}^{\infty} c_n$ to converge?

9. What is the dimension of the vector subspace of \mathbf{R}^n generated by the set of vectors

$$(\sigma(1), \sigma(2), \sigma(3), \dots, \sigma(n))$$

where σ runs through all $n!$ of the permutations of the first n natural numbers.

10. (a) Let

$$g(x, y) = x^2y + xy^2 + xy + x + y + 1.$$

We form a sequence $\{x_n\}$ as follows: $x_0 = 0$. The next term x_1 is the unique root -1 of the linear equation $g(t, 0) = 0$. For each $n \geq 2$, x_n is the root other than x_{n-2} of the equation $g(t, x_{n-1}) = 0$.

Let $\{f_n\}$ be the Fibonacci sequence determined by $f_0 = 0$, $f_1 = 1$ and $f_n = f_{n-1} + f_{n-2}$ for $n \geq 2$. Prove that, for any nonnegative integer k ,

$$x_{2k} = \frac{f_k}{f_{k+1}} \quad \text{and} \quad x_{2k+1} = -\frac{f_{k+2}}{f_{k+1}}.$$

- (b) Let

$$h(x, y) = x^2y + xy^2 + \beta xy + \gamma(x + y) + \delta$$

be a polynomial with real coefficients β, γ, δ . We form a bilateral sequence $\{x_n : n \in \mathbf{Z}\}$ as follows. Let $x_0 \neq 0$ be given arbitrarily. We select x_{-1} and x_1 to be the two roots of the quadratic equation $h(t, x_0) = 0$ in either order. From here, we can define inductively the terms of the sequence for positive

and negative values of the index so that x_{n-1} and x_{n+1} are the two roots of the equation $h(t, x_n) = 0$. We suppose that at each stage, neither of these roots is zero.

Prove that the sequence $\{x_n\}$ has period 5 (i.e. $x_{n+5} = x_n$ for each index n) if and only if $\gamma^3 + \delta^2 - \beta\gamma\delta = 0$.

UNDERGRADUATE MATHEMATICS COMPETITION

Saturday, March 5, 2016

Time: $3\frac{1}{2}$ hours

1. Let a be a positive real number that is not an integer and let

$$n = \left\lfloor \frac{1}{a - [a]} \right\rfloor.$$

Prove that $\lfloor (n+1)a \rfloor - 1$ is divisible by $n+1$. [Note: $\lfloor x \rfloor$ denotes the largest integer that is not greater than x , so that $\lfloor x \rfloor \leq x < \lfloor x \rfloor + 1$.]

2. Determine all polynomial solutions $f(x)$ to the identity

$$f(x + y - xy) = f(x)f(y).$$

3. Let $n = \prod p^a$ be the prime factor decomposition of the positive integer n and define $s(n) = \sum ap$, the sum of all the primes involved in the decomposition counting repetitions. For each positive integer m exceeding 1, let $h(m)$ be the number of positive integers n for which $s(n) = m$.

(a) Prove that $\lim_{n \rightarrow \infty} s(n) = \infty$.

(b) Prove that $s(n)$ assumes every value exceeding 4 at least twice and that $\lim_{n \rightarrow \infty} h(n) = \infty$.

4. Let $p(x)$ be a monic polynomial of degree 3 with three distinct real roots. How many real roots does the polynomial $(p'(x))^2 - 2p(x)p''(x)$ have?

5. (a) Determine the largest positive integer n for which the following statement is NOT true:

There exists a finite set $\{a_1, a_2, \dots, a_k\}$ ($k \geq 1$) of positive integers for which $n < a_1 < a_2 < \dots < a_k \leq 2n$ and $n \times a_1 \times a_2 \times \dots \times a_k$ is a perfect square.

(b) Determine infinitely many integers n for which $n < a_1 < a_2 < \dots < a_k$ and $n \times a_1 \times a_2 \times \dots \times a_k$ is square implies that $a_k \geq 2n$.

(c) Let $n = m^2$. Is it possible to determine an integer m for which integers a_1, a_2, \dots, a_k can be chosen in the open interval $(m^2, (m+1)^2)$ for which the product $a_1 \times a_2 \times \dots \times a_k$ is square?

6. Suppose that f is a strictly increasing convex real-valued continuous function on $[0, 1]$ for which $f(0) = 0$ and $f(1) = 1$ and $g(x)$ is a function that satisfies $g(f(x)) = x$ for each $x \in [0, 1]$. Prove that

$$\int_0^1 f(x)g(x)dx \leq \frac{1}{3}.$$

When does equality occur?

[Note: A function is *convex* if for any $t \in [0, 1]$

$$f((1-t)x + ty) \leq (1-t)f(x) + tf(y)$$

whenever $x, y, (1-t)x + ty$ belong to the domain of f .]

7. Let m, n be integers for which $0 \leq m < n$ and let $p(x)$ be a polynomial of degree n over a field \mathbf{F} . What is the dimension over \mathbf{F} of the vector space generated by the set of functions

$$\{1, x, x^2, \dots, x^{n-m-1}, p(x), p(x+1), \dots, p(x+m)\}?$$

8. Let S be a set of the positive integers that is closed under addition (i.e., $x, y \in S \Rightarrow x+y \in S$) for which the set T of positive integers not contained in S is finite with $m \geq 1$ elements. Prove that the sum of the numbers in T is not greater than m^2 and determine all the sets S for which this sum is equal to m^2 .
9. (a) Prove that every polyhedron has at least two faces with the same number of edges.
(b) Suppose that $k \geq 3$ and that all the faces in a polyhedron have at least k edges. Prove that there are k pairs of faces with the same number of edges (the pairs need not be disjoint).
10. Let X be a subset of the group G such that

$$\bigcap \{x^{-1}X : x \in X\}$$

contains an element a of finite order other than the identity. Prove that X is the union of cosets with respect to some subgroup of G . [Note: for any set S and element g of G , $gS = \{gs : s \in S\}$.]

UNDERGRADUATE MATHEMATICS COMPETITION

March 12, 2017

Time: $3\frac{1}{2}$ hours

1. Determine the value of the infinite product

$$\prod_{n=2}^{\infty} \left(1 - \frac{2}{1+n^3}\right).$$

2. Let S be a set of n points in the plane, no two pairs the same distance apart. Each point is joined by a straight line segment to the point that is nearest to it; no other segments are drawn. Prove:
- (a) No two segments have a point in common except possibly a point in S ;
(b) No point can be joined to more than five other points;
(c) The set of segments contains no cycle. In other words, there is no set $\{A_1, A_2, \dots, A_k\}$ of points in S , with $k \geq 3$, such that A_k is joined to A_1 and A_i is joined to A_{i+1} for $1 \leq i \leq k-1$.
3. (a) Given six irrational real numbers, prove that there are always two subsets of three (not necessarily disjoint) such that the sum of any two numbers in each of the subsets is irrational.
(b) Give an example of a set of six irrational numbers for which there are exactly two subsets of three numbers with all pair sums irrational.
4. 54 and 96 are two nonsquare positive integers whose product is a square; the squares 64 and 81 lie between them. Prove or disprove: *if m and n are two distinct nonsquare positive integers such that mn is a square, then there exists a square integer between them.*
5. Let $f(x)$ be a real continuous periodic function defined on the real numbers such that, for each positive integer n ,

$$\frac{|f(1)|}{1} + \frac{|f(2)|}{2} + \dots + \frac{|f(n)|}{n} \leq 1.$$

Prove that there exists a real number r such that $f(r) = f(r + 1) = 0$.

- For $n \geq 2$, let M be a $n \times n$ matrix with n distinct eigenvalues (exactly) one of which is 0. Suppose that u is a nonzero row n -vector and v is a nonzero column n -vector for which both uM and Mv are zero N -vectors. Prove that $uv \neq 0$ (i.e., u is not orthogonal to v).
- Let $p(z)$ be a polynomial of degree n , all of whose roots have absolute value 1. Prove that $|p'(1)| \geq \frac{n}{2}|p(1)|$.
- Suppose that the real function $y = f(x)$ defined on $[0, \infty)$ satisfies the differential equation

$$y' = \frac{1}{x^2 + y^2}$$

with the initial condition $f(0) > 0$. Prove that $\lim_{x \rightarrow \infty} f(x)$ exists and that this limit exceeds 1.

- Let $f(x)$ be a real-valued continuous function defined on the closed interval $[0, 1]$ for which

$$1 = \int_0^1 f(x) dx = \int_0^1 x f(x) dx.$$

Prove that

$$\int_0^1 (f(x))^2 dx \geq 4.$$

- Prove that

$$0 < \int_{\pi/8}^{3\pi/8} \frac{\cos 2x dx}{1 + \tan x} < \frac{\pi}{8\sqrt{2}}.$$

UNDERGRADUATE MATHEMATICS COMPETITION

Sunday, March 10, 2019

Time: 3½ hours

No aids or calculators permitted.

The grading is designed to encourage only the stronger students to attempt more than five problems. Each solution is graded out of 10. If the sum of the scores for the solutions to the five best problems does not exceed 30, this sum will be the final grade. If the sum of these scores does exceed 30, then all solutions will be graded for credit.

- (a) Determine necessary and sufficient conditions on the sextuple (a, b, c, d, e, f) with $a \leq b \leq c \leq d \leq e \leq f$ in order that there exist four numbers for which a, b, c, d, e, f are the pairwise sums.
(b) Where the set of four numbers exist as in (a), determine when there is more than one possibility and show that the sum of the squares of the numbers is the same for the different possibilities.
- For $n = 1, 2, \dots$, let

$$x_n = \frac{n+1}{2^{n+1}} \sum_{k=1}^n \frac{2^k}{k}.$$

Prove that $\lim_{n \rightarrow \infty} x_n$ exists and find it.

- The positive integer n is said to be an SP (square-pair) number if the set $\{1, 2, \dots, 2n\}$ can be partitioned into n pairs such that the sum of the numbers in each pair is a perfect square.

(a) Prove that n is an SP number for which the sums of the members of the pairs are equal if and only if $n = 2m(m + 1)$ for some positive integer m .

(b) Show that 10 is not a SP number.

(c) Prove that there are infinitely many SP numbers that are not of the form $2m(m + 1)$.

4. Determine

$$\int_0^1 \left(4 \arctan x + \pi \tan \frac{\pi x}{4} \right) dx.$$

5. Let $n \geq 2$ and let x_1, x_2, \dots, x_n be positive real numbers. Determine the minimum value of

$$S = \frac{x_1}{x_2 + x_3 + \dots + x_n} + \frac{x_2}{x_1 + x_3 + \dots + x_n} + \dots + \frac{x_n}{x_1 + x_2 + \dots + x_{n-1}}$$

(n terms on the right side) and the value of (x_1, x_2, \dots, x_n) where this minimum is attained.

6. Let m, n be positive integers; let A be a $m \times n$ matrix, B be a $n \times m$ matrix, and BA an invertible $n \times n$ matrix.

(a) What are the eigenvalues of the matrix $A(BA)^{-1}B$?

(b) Give an example of this situation when $(m, n) = (3, 2)$.

Please turn over.

7. Let $p(z) = z^3 + az^2 + bz + c$ be a cubic polynomial with complex coefficients. Prove that

$$\sup\{|p(z)| : |z| \leq 1\} \geq 1.$$

8. Suppose a and b are elements of a group and r is a positive integer for which $(ab)^r = a$. Prove that $ab = ba$.

9. Suppose that $\{f_n(x) : n = 1, 2, 3, \dots\}$ is a sequence of differentiable functions defined on an open interval I for which $|f'_n(x)| \leq M$ for some constant M , all $x \in I$ and all positive integers n . Suppose further that $\lim_{n \rightarrow \infty} f_n(x) = g(x)$. Prove that $g(x)$ is continuous on I .

10. Let S be the subset of the open interval $(0, 1)$ consisting of those numbers that do not have a terminating binary (base 2) expansion (*i.e.*, are not rationals of the form $r/2^s$ for integers r and s). For $x \in (0, 1)$, let $f_n(x)$ be the number of 0's among the first n digits after the decimal point in the binary expansion of x , divided by n . Let A be the set of numbers x in S for which

$$f(x) = \lim_{n \rightarrow \infty} f_n(x)$$

exists, and let B be the set of numbers in S for which the limit does not exist.

(a) Prove that both A and B are uncountable.

(b) Does there exist a number in A for which $f(x) = x$?

END