


What are w-Trivalent Tangles? (PA := Planar Algebra)

$\{ \text{knots} \} = \text{PA} \langle \text{R123} : \text{R123} \rangle$
 $\{ \text{trivalent tangles} \} = \text{PA} \langle \text{R23, R4} \rangle$
 $\{ \text{w-tangles} \} = \text{PA} \langle \text{w-generators} \mid \text{w-relations} \mid \text{unary w-operations} \rangle$

The w-generators.

Crossing, Broken surface, 2D Symbol, Dim. reduc., Virtual crossing, Movie



Cap, Wen, w, Vertices, singular, smooth


Homomorphic expansions for a filtered algebraic structure \mathcal{K} :

$$\text{ops}^{\leftarrow} \mathcal{K} = \mathcal{K}_0 \supset \mathcal{K}_1 \supset \mathcal{K}_2 \supset \mathcal{K}_3 \supset \dots$$

$$\text{ops}^{\leftarrow} \text{gr } \mathcal{K} := \mathcal{K}_0/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \dots$$

An **expansion** is a filtration $Z : \mathcal{K} \rightarrow \text{gr } \mathcal{K}$ that "covers" the identity on $\text{gr } \mathcal{K}$. A **homomorphic expansion** is an expansion that respects all relevant "extra" operations.

Filtered algebraic structures are cheap and plenty. In any \mathcal{K} , allow formal linear combinations, let \mathcal{K}_1 be the ideal generated by differences (the "augmentation ideal"), and let $\mathcal{K}_m := \langle (\mathcal{K}_1)^m \rangle$ (using all available "products").



- Has kinds, objects, operations, and maybe constants.
- Perhaps subject to some axioms.
- We always allow formal linear combinations.


Example: Pure Braids. PB_n is generated by x_{ij} , "strand i goes around strand j once", modulo "Reidemeister moves". $A_n := \text{gr } PB_n$ is generated by $t_{ij} := x_{ij} - 1$, modulo the 4T relations $[t_{ij}, t_{ik} + t_{jk}] = 0$ (and some lesser ones too). Much happens in A_n , including the Drinfel'd theory of associators.

Our case(s).

$\mathcal{K} \xrightarrow{Z: \text{high algebra}} \mathcal{A} := \text{gr } \mathcal{K} \xrightarrow{\text{given a "Lie" algebra } \mathfrak{g}} \mathcal{U}(\mathfrak{g})$
 solving finitely many equations in finitely many unknowns low algebra: pictures represent formulas

\mathcal{K} is knot theory or topology; $\text{gr } \mathcal{K}$ is finite combinatorics: bounded-complexity diagrams modulo simple relations.

A **Ribbon 2-Knot** is a surface S embedded in \mathbb{R}^4 that bounds an immersed handlebody B , with only "ribbon singularities"; a ribbon singularity is a disk D of transverse double points, whose preimages in B are a disk D_1 in the interior of B and a disk D_2 with $D_2 \cap \partial B = \partial D_2$, modulo isotopies of S alone.



The **w-relations** include R234, VR1234, M, Overcrossings Commute (OC) but not UC, $W^2 = 1$, and funny interactions between the wen and the cap and over- and under-crossings:

OC:

yet not UC:

Challenge: Do the Reidemeister!

Reidemeister Winter

The unary w-operations

Unzip along an annulus

Unzip along a disk

Just for fun.

$\mathcal{K} = \{ \text{Reidemeister} \} = \left(\text{The set of all b/w 2D projections of reality} \right)$
 $\mathcal{K}/\mathcal{K}_1 \leftarrow \mathcal{K}/\mathcal{K}_2 \leftarrow \mathcal{K}/\mathcal{K}_3 \leftarrow \mathcal{K}/\mathcal{K}_4 \leftarrow \dots$


Crop Rotate Adjoin

An expansion Z is a choice of a "progressive scan" algorithm.


$\mathcal{K}/\mathcal{K}_1 \oplus \mathcal{K}_1/\mathcal{K}_2 \oplus \mathcal{K}_2/\mathcal{K}_3 \oplus \mathcal{K}_3/\mathcal{K}_4 \oplus \mathcal{K}_4/\mathcal{K}_5 \oplus \mathcal{K}_5/\mathcal{K}_6 \oplus \dots$
 $\parallel \quad \parallel$
 $\mathbb{R} \quad \ker(\mathcal{K}/\mathcal{K}_4 \rightarrow \mathcal{K}/\mathcal{K}_3)$

Convolutions on Lie Groups and Lie Algebras and Ribbon 2-Knots, Page 2

Knot-Theoretic statement. There exists a homomorphic expansion Z for trivalent w-tangles. In particular, Z should respect $R4$ and intertwine annulus and disk unzips:



Diagrammatic statement. Let $R = \exp \uparrow \in \mathcal{A}^w(\uparrow\uparrow)$. There exist $\omega \in \mathcal{A}^w(\uparrow)$ and $V \in \mathcal{A}^w(\uparrow\uparrow)$ so that



Algebraic statement. With $I\mathfrak{g} := \mathfrak{g}^* \rtimes \mathfrak{g}$, with $c : \hat{U}(I\mathfrak{g}) \rightarrow \hat{U}(I\mathfrak{g})/\hat{U}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$ the obvious projection, with S the antipode of $\hat{U}(I\mathfrak{g})$, with W the automorphism of $\hat{U}(I\mathfrak{g})$ induced by flipping the sign of \mathfrak{g}^* , with $r \in \mathfrak{g}^* \otimes \mathfrak{g}$ the identity element and with $R = e^r \in \hat{U}(I\mathfrak{g}) \otimes \hat{U}(\mathfrak{g})$ there exist $\omega \in \hat{S}(\mathfrak{g}^*)$ and $V \in \hat{U}(I\mathfrak{g})^{\otimes 2}$ so that

(1) $V(\Delta \otimes 1)(R) = R^{13}R^{23}V$ in $\hat{U}(I\mathfrak{g})^{\otimes 2} \otimes \hat{U}(\mathfrak{g})$
 (2) $V \cdot SWV = 1$ (3) $(c \otimes c)(V\Delta(\omega)) = \omega \otimes \omega$

Unitary statement. There exists $\omega \in \text{Fun}(\mathfrak{g})^G$ and an (infinite order) tangential differential operator V defined on $\text{Fun}(\mathfrak{g}_x \times \mathfrak{g}_y)$ so that

(1) $V\widehat{e^{x+y}} = \widehat{e^x e^y} V$ (allowing $\hat{U}(\mathfrak{g})$ -valued functions)
 (2) $VV^* = I$ (3) $V\omega_{x+y} = \omega_x \omega_y$


Group-Algebra statement. There exists $\omega^2 \in \text{Fun}(\mathfrak{g})^G$ so that for every $\phi, \psi \in \text{Fun}(\mathfrak{g})^G$ (with small support), the following holds in $\hat{U}(\mathfrak{g})$:

$$\iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_{x+y}^2 e^{x+y} = \iint_{\mathfrak{g} \times \mathfrak{g}} \phi(x)\psi(y)\omega_x^2 \omega_y^2 e^x e^y. \quad (\text{shhh, this is Duflo})$$


Convolutions statement (Kashiwara-Vergne). Convolutions of invariant functions on a Lie group agree with convolutions of invariant functions on its Lie algebra. More accurately, let G be a finite dimensional Lie group and let \mathfrak{g} be its Lie algebra, let $j : \mathfrak{g} \rightarrow \mathbb{R}$ be the Jacobian of the exponential map $\exp : \mathfrak{g} \rightarrow G$, and let $\Phi : \text{Fun}(G) \rightarrow \text{Fun}(\mathfrak{g})$ be given by $\Phi(f)(x) := j^{1/2}(x)f(\exp x)$. Then if $f, g \in \text{Fun}(G)$ are Ad-invariant and supported near the identity, then

$$\Phi(f) \star \Phi(g) = \Phi(f \star g).$$

From wTT to \mathcal{A}^w . $\text{gr}_m \text{wTT} := \{m\text{-cubes}\} / \{(m+1)\text{-cubes}\}$:



w-Jacobi diagrams and \mathcal{A} . $\mathcal{A}^w(Y \uparrow) \cong \mathcal{A}^w(\uparrow\uparrow\uparrow)$ is



Diagrammatic to Algebraic. With (x_i) and (φ^j) dual bases of \mathfrak{g} and \mathfrak{g}^* and with $[x_i, x_j] = \sum b_{ij}^k x_k$, we have $\mathcal{A}^w \rightarrow \mathcal{U}$ via

Unitary \iff Algebraic. The key is to interpret $\hat{U}(I\mathfrak{g})$ as tangential differential operators on $\text{Fun}(\mathfrak{g})$:

- $\varphi \in \mathfrak{g}^*$ becomes a multiplication operator.
- $x \in \mathfrak{g}$ becomes a tangential derivation, in the direction of the action of $\text{ad } x$: $(x\varphi)(y) := \varphi([x, y])$.
- $c : \hat{U}(I\mathfrak{g}) \rightarrow \hat{U}(I\mathfrak{g})/\hat{U}(\mathfrak{g}) = \hat{S}(\mathfrak{g}^*)$ is “the constant term”.

Unitary \implies Group-Algebra. $\iint \omega_{x+y}^2 e^{x+y} \phi(x)\psi(y) = \langle \omega_{x+y}, \omega_{x+y} e^{x+y} \phi(x)\psi(y) \rangle = \langle V\omega_{x+y}, V e^{x+y} \phi(x)\psi(y)\omega_{x+y} \rangle = \langle \omega_x \omega_y, e^x e^y V \phi(x)\psi(y)\omega_{x+y} \rangle = \langle \omega_x \omega_y, e^x e^y \phi(x)\psi(y)\omega_x \omega_y \rangle = \iint \omega_x^2 \omega_y^2 e^x e^y \phi(x)\psi(y).$



Convolutions and Group Algebras (ignoring all Jacobians). If G is finite, A is an algebra, $\tau : G \rightarrow A$ is multiplicative then $(\text{Fun}(G), \star) \cong (A, \cdot)$ via $L : f \mapsto \sum f(a)\tau(a)$. For Lie (G, \mathfrak{g}) ,

$$\begin{array}{ccc} (\mathfrak{g}, +) \ni x \xrightarrow{\tau_0 = \exp_S} e^x \in \hat{S}(\mathfrak{g}) & & \text{Fun}(\mathfrak{g}) \xrightarrow{L_0} \hat{S}(\mathfrak{g}) \\ \downarrow \exp_G & \searrow \exp_U & \downarrow \chi \\ (G, \cdot) \ni e^x \xrightarrow{\tau_1} e^x \in \hat{U}(\mathfrak{g}) & & \text{Fun}(G) \xrightarrow{L_1} \hat{U}(\mathfrak{g}) \end{array} \quad \text{so} \quad \begin{array}{ccc} & & \downarrow \Phi^{-1} \\ & & \downarrow \chi \end{array}$$

with $L_0\psi = \int \psi(x)e^x dx \in \hat{S}(\mathfrak{g})$ and $L_1\Phi^{-1}\psi = \int \psi(x)e^x \in \hat{U}(\mathfrak{g})$. Given $\psi_i \in \text{Fun}(\mathfrak{g})$ compare $\Phi^{-1}(\psi_1) \star \Phi^{-1}(\psi_2)$ and $\Phi^{-1}(\psi_1 \star \psi_2)$ in $\hat{U}(\mathfrak{g})$: (shhh, $L_{0/1}$ are “Laplace transforms”)

$$\star \text{ in } G : \iint \psi_1(x)\psi_2(y)e^x e^y \quad \star \text{ in } \mathfrak{g} : \iint \psi_1(x)\psi_2(y)e^{x+y}$$

- We skipped...**
- The Alexander polynomial and Milnor numbers.
 - v-Knots, quantum groups and Etingof-Kazhdan.
 - u-Knots, Alekseev-Torossian, and BF theory and the successful and Drinfel'd associators.
 - The simplest problem hyperbolic geometry solves.



Draft

The Orbit Method. By Fourier analysis, the characters of $(\text{Fun}(\mathfrak{g})^G, \star)$ correspond to coadjoint orbits in \mathfrak{g}^* . By averaging representation matrices and using Schur's lemma to replace intertwiners by scalars, to every irreducible representation of G we can assign a character of $(\text{Fun}(G)^G, \star)$.

Measure theoretic statement. Ignoring all ω 's, there exists a measure preserving and orbit preserving transformation $T : \mathfrak{g}_x \times \mathfrak{g}_y \rightarrow \mathfrak{g}_x \times \mathfrak{g}_y$ for which $e^{x+y} \circ T = e^x e^y$.

Alekseev-Torossian statement. There is an element $F \in \text{TAut}_2$ with

$$F(x + y) = \log e^x e^y$$

and $j(F) \in \text{im } \tilde{\delta} \subset \text{tr}_2$, where for $a \in \text{tr}_1$, $\tilde{\delta}(a) := a(x) + a(y) - a(\log e^x e^y)$.

Free Lie statement (Kashiwara-Vergne). There exist convergent Lie series F and G so that with $z = \log e^x e^y$

$$x + y - \log e^y e^x = (1 - e^{-\text{ad } x})F + (e^{\text{ad } y} - 1)G$$

$$\begin{aligned} \text{tr}(\text{ad } x)\partial_x F + \text{tr}(\text{ad } y)\partial_y G = \\ \frac{1}{2} \text{tr} \left(\frac{\text{ad } x}{e^{\text{ad } x} - 1} + \frac{\text{ad } y}{e^{\text{ad } y} - 1} - \frac{\text{ad } z}{e^{\text{ad } z} - 1} - 1 \right) \end{aligned}$$

Δ acts by double and sum, S by reverse and negate.