



Plot [ $\omega = e^{it}$ ;  $u = \text{Re}[\omega^{1/2}]$ ;  $v = \text{Re}[\omega]$ ;  $-(\text{MatrixSignature}[A] - \text{Writhe}[K]) / 2$ ,  $\{t, 0, 2\pi\}$ ]




Plot [Bed[Knot[8, 2],  $e^{it}$ ],  $\{t, 0, 2\pi\}$ ]



### Kashaev for Mathematicians.


For a knot  $K$  and a complex unit  $\omega$  set  $u = \Re(\omega^{1/2})$ ,  $v = \Re(\omega)$ , make an  $F \times F$  matrix  $A$  with contributions



and output  $\frac{1}{2}(\sigma(A) - w(K))$ .

### Bedlewo for Mathematicians.

For a knot  $K$  and a complex unit  $\omega$  set  $t = 1 - \omega$ ,  $r = 2\Re(t)$ , make an  $F \times F$  matrix  $A$  with contributions



(conjugate if going against the flow) and output  $\sigma(A)$ .

### Why are they equal?


I dunno, yet note that

- ▶ Kashaev is over the  $\mathbb{R}$ eals, Bedlewo is over the  $\mathbb{C}$ omplex numbers.
- ▶ There's a factor of 2 between them, and a shift.

...so it's not merely a matrix manipulation.

**Theorem.** The Bedlewo program computes the Levine-Tristram signature of  $K$  at  $\omega$ .

(Easy) **Proof.** Levine and Tristram tell us to look at  $\sigma((1 - \omega)L + (1 - \omega^*)L^T)$ , where  $L$  is the linking matrix for a Seifert surface  $S$  for  $K$ :  $L_{ij} = \text{lk}(\gamma_i, \gamma_j^+)$  where  $\gamma_i$  run over a basis of  $H_1(S)$  and  $\gamma_i^+$  is the pushout of  $\gamma_i$ . But signatures don't change if you run over an over-determined basis, and the faces make such an over-determined basis whose linking numbers are controlled by the crossings. The rest is details.



Art by Emily Redelmeier

Thank You!