## What happens to a quantum particle on a pendulum at $T = \frac{\pi}{2}$ ?

Abstract. This subject is the best one-hour introduction I know for the mathematical techniques that appear in quantum mechanics — in one short lecture we start with a meaningful question, visit Schrödinger's equation, operators and exponentiation of operators, Fourier analysis, path integrals, the least action principle, and Gaussian integration, and at the end we land with a meaningful and interesting answer.

Based a lecture given by the author in the "trivial notions" seminar in Harvard on April 29, 1989. This edition, January 10, 2014.

## 1. The Question

Let the complex valued function  $\psi = \psi(t, x)$  be a solution of the Schrödinger equation

$$\frac{\partial \psi}{\partial t} = -i\left(-\frac{1}{2}\Delta_x + \frac{1}{2}x^2\right)\psi$$
 with  $\psi|_{t=0} = \psi_0.$ 

What is  $\psi|_{t=T=\frac{\pi}{2}}$ ?

In fact, the major part of our discussion will work just as well for the general Schrödinger equation,

$$\frac{\partial \psi}{\partial t} = -iH\psi, \qquad H = -\frac{1}{2}\Delta_x + V(x)$$
$$\psi|_{t=0} = \psi_0, \qquad \text{arbitrary } T,$$

where,

- $\psi$  is the "wave function", with  $|\psi(t, x)|^2$  representing the probability of finding our particle at time t in position x.
- *H* is the "energy", or the "Hamiltonian".
- $-\frac{1}{2}\Delta_x$  is the "kinetic energy".
- V(x) is the "potential energy at x".

## 2. The Solution

The equation  $\frac{\partial \psi}{\partial t} = -iH\psi$  with  $\psi|_{t=0} = \psi_0$  formally implies

$$\psi(T,x) = \left(e^{-iTH}\psi_0\right)(x) = \left(e^{i\frac{T}{2}\Delta - iTV}\psi_0\right)(x).$$

By Lemma 3.1 with  $n = 10^{58} + 17$  and setting  $x_n = x$ we find that  $\psi(T, x)$  is

$$\left(e^{i\frac{T}{2n}\Delta}e^{-i\frac{T}{n}V}e^{i\frac{T}{2n}\Delta}e^{-i\frac{T}{n}V}\dots e^{i\frac{T}{2n}\Delta}e^{-i\frac{T}{n}V}\psi_0\right)(x_n)$$

Now using Lemmas 3.2 and 3.3 we find that this is:  $(c \text{ denotes the ever-changing universal fixed numer$  $ical constant})$ 

$$c \int dx_{n-1} e^{i \frac{(x_n - x_{n-1})^2}{2T/n}} e^{-i \frac{T}{N} V(x_{n-1})} \dots$$
$$\int dx_1 e^{i \frac{(x_2 - x_1)^2}{2T/n}} e^{-i \frac{T}{N} V(x_1)}$$
$$\int dx_0 e^{i \frac{(x_1 - x_0)^2}{2T/n}} e^{-i \frac{T}{N} V(x_0)} \psi_0(x_0)$$

Repackaging, we get

$$c \int dx_0 \dots dx_{n-1} \\ \exp\left(i\frac{T}{2n}\sum_{k=1}^n \left(\frac{x_k - x_{k-1}}{T/n}\right)^2 - i\frac{T}{n}\sum_{k=0}^{n-1} V(x_k)\right) \\ \psi_0(x_0).$$

Now comes the novelty. keeping in mind the picture  $x_{h}$ 



and replacing Riemann sums by integrals, we can write

$$\psi(T,x) = c \int dx_0 \int_{W_{x_0x_n}} \mathcal{D}x$$
$$\exp\left(i \int_0^T dt \left(\frac{1}{2}\dot{x}^2(t) - V(x(t))\right)\right) \psi_0(x_0),$$

where  $W_{x_0x_n}$  denotes the space of paths that begin at  $x_0$  and end at  $x_n$ ,

$$W_{x_0x_n} = \{ x : [0,T] \to \mathbb{R} : x(0) = x_0, x(T) = x_n \},\$$

and  $\mathcal{D}x$  is the formal "path integral measure". This is a good time to introduce the "action"  $\mathcal{L}$ :

$$\mathcal{L}(x) := \int_0^T dt \left( \frac{1}{2} \dot{x}^2(t) - V(x(t)) \right).$$

With this notation,

$$\psi(T,x) = c \int dx_0 \psi_0(x_0) \int_{W_{x_0} x_n} \mathcal{D}x e^{i\mathcal{L}(x)}.$$

Video and more at http://drorbn.net/?title=AKT-14 (Jan 10 and Jan 17 classes)