

# Meta-Groups, Meta-Bicrossed-Products, and the Alexander Polynomial, 1



Dror Bar-Natan at Sheffield, February 2013.

<http://www.math.toronto.edu/~drorbn/Talks/Sheffield-130206/>



**Abstract.** I will define “meta-groups” and explain how one specific meta-group, which in itself is a “meta-bicrossed-product”, gives rise to an “ultimate Alexander invariant” of tangles, that contains the Alexander polynomial (multivariable, if you wish), has extremely good composition properties, is evaluated in a topologically meaningful way, and is least-wasteful in a computational sense. If you believe in categorification, that’s a wonderful playground.

This work is closely related to work by Le Dimet (Comment. Math. Helv. **67** (1992) 306–315), Kirk, Livingston and Wang (arXiv:math/9806035) and Cimasoni and Turaev (arXiv:math.GT/0406269).



**A Standard Alexander Formula.** Label the arcs  $1$  through  $(n+1) = 1$ , make an  $n \times n$  matrix as below, delete one row and one column, and compute the determinant:

$$\begin{array}{c} \begin{array}{ccc} \begin{array}{c} \nearrow \\ \searrow \end{array} & \begin{array}{c} b \\ a \end{array} \\ \begin{array}{c} \nwarrow \\ \nearrow \end{array} & \begin{array}{c} c \\ a \end{array} \end{array} \rightarrow \begin{array}{c|cc} & a & b & c \\ \hline c & -1 & 1-X & X \end{array}$$

$$\begin{array}{c} \begin{array}{ccc} \begin{array}{c} \nwarrow \\ \nearrow \end{array} & \begin{array}{c} b \\ a \end{array} \\ \begin{array}{c} \nearrow \\ \nwarrow \end{array} & \begin{array}{c} c \\ a \end{array} \end{array} \rightarrow \begin{array}{c|cc} & a & b & c \\ \hline c & -X & X-1 & 1 \end{array}$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & x-1 & 0 & -x \\ -1 & x & 0 & 0 & 0 & 0 & 1-x & 0 \\ 0 & -1 & x & 0 & 1-x & 0 & 0 & 0 \\ x-1 & 0 & -x & 1 & 0 & 0 & 0 & 0 \\ 0 & 1-x & 0 & -1 & x & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x & 1 & 0 & x-1 \\ 0 & 0 & 1-x & 0 & 0 & -1 & x & 0 \\ 0 & 0 & 0 & x-1 & 0 & 0 & -x & 1 \end{pmatrix} \quad [[1 ;; 7, 1 ;; 7]] \quad // \text{Det}$$

$$-1 + 4x - 8x^2 + 11x^3 - 8x^4 + 4x^5 - x^6$$

## Alexander Issues.

- Quick to compute, but computation departs from topology.
- Extends to tangles, but at an exponential cost.
- Hard to categorify.


**Idea.** Given a group  $G$  and two “YB” pairs  $R^\pm = (g_o^\pm, g_u^\pm) \in G^2$ , map them to xings and “multiply along”, so that

$$\begin{array}{c} \text{Xing} \end{array} \xrightarrow{Z} \begin{array}{c} \text{Xing} \end{array} \begin{array}{c} g_o^\pm \\ g_u^\pm \end{array}$$

$$\begin{array}{c} \text{Xing} \end{array} \xrightarrow{Z} \begin{pmatrix} g_o^+ g_u^+ g_o^+ g_u^- g_o^- g_u^+ g_o^- g_u^+ \\ g_u^- g_o^- \end{pmatrix}$$

**This Fails!** R2 implies that  $g_o^\pm g_o^\mp = e = g_u^\pm g_u^\mp$  and then R3 implies that  $g_o^+$  and  $g_u^+$  commute, so the result is a simple counting invariant.

**A Group Computer.** Given  $G$ , can store group elements and perform operations on them:



Also has  $S_x$  for inversion,  $e_x$  for unit insertion,  $d_x$  for register deletion,  $\Delta_{xy}^z$  for element cloning,  $\rho_y^x$  for renamings, and  $(D_1, D_2) \mapsto D_1 \cup D_2$  for merging, and many obvious composition axioms relating those.

$$P = \{x : g_1, y : g_2\} \Rightarrow P = \{d_y P\} \cup \{d_x P\}$$

**A Meta-Group.** Is a similar “computer”, only its internal structure is unknown to us. Namely it is a collection of sets  $\{G_\gamma\}$  indexed by all finite sets  $\gamma$ , and a collection of operations  $m_z^{xy}$ ,  $S_x$ ,  $e_x$ ,  $d_x$ ,  $\Delta_{xy}^z$  (sometimes),  $\rho_y^x$ , and  $\cup$ , satisfying the exact same *linear* properties.

**Example 0.** The non-meta example,  $G_\gamma := G^\gamma$ .

**Example 1.**  $G_\gamma := M_{\gamma \times \gamma}(\mathbb{Z})$ , with simultaneous row and column operations, and “block diagonal” merges. Here if  $P = \begin{pmatrix} x & a & b \\ y & c & d \end{pmatrix}$  then  $d_y P = (x : a)$  and  $d_x P = (y : d)$  so  $\{d_y P\} \cup \{d_x P\} = \begin{pmatrix} x & a & 0 \\ y & 0 & d \end{pmatrix} \neq P$ . So this  $G$  is truly meta.

**Claim.** From a meta-group  $G$  and YB elements  $R^\pm \in G_2$  we can construct a knot/tangle invariant.

**Bicrossed Products.** If  $G = HT$  is a group presented as a product of two of its subgroups, with  $H \cap T = \{e\}$ , then also  $G = TH$  and  $G$  is determined by  $H$ ,  $T$ , and the “swap” map  $sw^{th} : (t, h) \mapsto (h', t')$  defined by  $th = h't'$ . The map  $sw$  satisfies (1) and (2) below; conversely, if  $sw : T \times H \rightarrow H \times T$  satisfies (1) and (2) (+ lesser conditions), then (3) defines a group structure on  $H \times T$ , the “bicrossed product”.

