MAT 347

The symmetric and the alternating groups October 10, 2014

Recall some definitions.:

- A permutation of n elements is an element of the group S_n .
- A cycle of length m is a permutation that can be written as $(a_1 \cdots a_m)$.
- A transposition is a permutation of length 2.
- The type of a permutation is the set of lengths of the cycles in its decomposition as product of disjoint cycles. For example the type of (12345)(67)(89) in S_{11} is (5, 2, 2, 1, 1).
- 1. In general, for an arbitrary group G, the *conjugacy class* of $g \in G$ is the orbit of g in the action of G on G by conjugation. Find a description of the conjugacy classes of S_n .

Hint: Fix your favourite $\sigma \in S_n$. Then for various $\tau \in S_n$ compute $\tau \sigma \tau^{-1}$. Can you find a formula for $\tau \sigma \tau^{-1}$? Can you describe the conjugacy class of σ ?

2. List all the conjugacy classes of S_5 and the size of each class.

Hint: You know the sum of the sizes of all the conjugacy classes should be 120, so you can check your final answer.

- 3. Which of the following sets are generators of S_n ?
 - (a) The set of all cycles.
 - (b) The set of all transpositions.
 - (c) The set $\{(12), (23), (34), \dots, (n-1, n)\}.$
 - (d) The set $\{(12), (13), (14), \dots, (1n)\}$.

4.

- (a) Write the permutation (123) as product of transpositions. This can be done in more than one way. Try to write (123) as product of N transpositions, for different values of N. Not all values of N are possible. Which ones are?
- (b) Repeat the same question with the permutation (1234).

Note: At this point, you can probably make a conjecture for which values of N are not possible, but most likely you won't be able to prove it. For that, we need to introduce some sophistication.

Building the alternating group

Let us fix a positive integer n. Let R be the set of polynomials in the n variables $X_1, \ldots X_n$. We can define an action of S_n on R as follows:

$$\sigma \cdot p(x_1, \dots, x_n) := p(x_{\sigma(1)}, \dots, x_{\sigma(n)})$$

Make sure you understand what this notation means before continuing. Convince yourself that it is, indeed, an action. You will work further with this action in HW 5 (both to understand this proof, and because it will be relevant in Galois Theory at the end of the course.) We define the following polynomial:

$$\Delta := \prod_{1 \le i < j \le n} (X_i - X_j)$$

For example, if n = 3, then $\Delta = (X_1 - X_2)(X_1 - X_3)(X_2 - X_3)$.

- 5. Prove that for every $\sigma \in S_n$ there exists a number $\varepsilon_{\sigma} \in \{1, -1\}$ such that $\sigma \cdot \Delta = \varepsilon_{\sigma} \Delta$
- 6. Prove that the map $\varepsilon: S_n \to \{1, -1\}$ is a group homomorphism!

We say that a permutation σ is *even* when $\varepsilon_{\sigma} = 1$ and it is *odd* when $\varepsilon_{\sigma} = -1$. When we mention the *parity* of a permutation, we are referring to whether it is odd or even. We define A_n to be the set of all even permutations.

- 7. Complete: "A cycle of length m is an even permutation iff m is"
- 8. Go back to the conjecture you made in Question 4. Now you can prove it!
- 9. Prove that A_n is a normal subgroup of S_n .
- 10. What is $|A_n|$?

Hint: Use the first isomorphism theorem.

11. Review your answer to Question 2. Which of the conjugacy classes are in A_5 ? Do their sizes add up to the right number?

The platonic solids

12. Each one of the five platonic solids has a group of rotations that is isomorphic to either some S_n or some A_n . Find them all (with proof).