

## 5. Some Problems in Heaven

Unfortunately,  $\dim \mathcal{A}(\mathcal{X}, X) = \dim \Lambda(\mathcal{X}, X) = 4^{|\mathcal{X}|}$  is big. Fortunately, we have the following theorem, a version of one of the main results in Halacheva's thesis, [Ha1, Ha2]:

**Theorem.** Working in  $\Lambda(\mathcal{X} \cup X)$ , if  $w = \omega e^\lambda$  is a balanced Gaussian (namely, a scalar  $\omega$  times the exponential of a quadratic  $\lambda = \sum_{\zeta \in \mathcal{X}, z \in X} \alpha_{\zeta, z} \zeta z$ ), then generically so is  $c_{x, \xi} e^\lambda$ .  
(This is great news! The space of balanced quadratics is only  $|\mathcal{X}||X|$ -dimensional!)

**Proof.** Recall that  $c_{x, \xi}: (1, \xi, x, x\xi)w' \mapsto (1, 0, 0, 1)w'$ , write  $\lambda = \mu + \eta x + \xi y + \alpha \xi x$ , and ponder  $e^\lambda =$

$$\dots + \frac{1}{k!} \underbrace{(\mu + \eta x + \xi y + \alpha \xi x)(\mu + \eta x + \xi y + \alpha \xi x) \cdots (\mu + \eta x + \xi y + \alpha \xi x)}_{k \text{ factors}} + \dots$$

Then  $c_{x, \xi} e^\lambda$  has three contributions:

- $e^\mu$ , from the term proportional to 1 (namely, independent of  $\xi$  and  $x$ ) in  $e^\lambda$
- $-\alpha e^\mu$ , from the term proportional to  $x\xi$ , where the  $x$  and the  $\xi$  come from the same factor above.
- $\eta y e^\mu$ , from the term proportional to  $x\xi$ , where the  $x$  and the  $\xi$  come from different factors above.

So  $c_{x, \xi} e^\lambda = e^\mu (1 - \alpha + \eta y) = (1 - \alpha) e^\mu (1 + \eta y / (1 - \alpha)) = (1 - \alpha) e^{\mu + \eta y / (1 - \alpha)}$ . □

## $\Gamma$ -calculus.

Thus we have an almost-always-defined “ $\Gamma$ -calculus”: a contraction algebra morphism  $\mathcal{T}(\mathcal{X}, X) \rightarrow R \times (\mathcal{X} \otimes_{R/R} X)$  whose behaviour under contractions is given by

$$c_{x, \xi}(\omega, \lambda = \mu + \eta x + \xi y + \alpha \xi x) = ((1 - \alpha)\omega, \mu + \eta y / (1 - \alpha)).$$

( $\Gamma$  is fully defined on pure tangles – tangles without closed components – and hence on long knots).

Multiplying and comparing  $\Gamma$  objects:

```

Γ /: Γ[is1_, os1_, cs1_, ω1_, λ1_] × Γ[is2_, os2_, cs2_, ω2_, λ2_] :=
  Γ[is1 ∪ is2, os1 ∪ os2, Join[cs1, cs2], ω1 ω2, λ1 + λ2]
Γ /: Γ[is1_, os1_, ω1_, λ1_] ≡ Γ[is2_, os2_, ω2_, λ2_] :=
  TrueQ[(Sort@is1 == Sort@is2) ∧ (Sort@os1 == Sort@os2) ∧
    Simplify[ω1 == ω2] ∧ CF@λ1 == CF@λ2]

```

No rules for linear operations!

The crossings and the point:

```

Γ[Xi_j, j, k, l, S_, T_] := Γ[{l, i}, {j, k}, <[xi_i → S, x_j → T, x_k → S, xi_l → T]>,
  T^{-1/2}, CF[{xi_l, xi_i} · (1 1 - T / 0 T) · {x_j, x_k}]];
Γ[Xi_j, j, k, l, S_, T_] := Γ[{i, j}, {k, l}, <[xi_i → S, xi_j → T, x_k → S, x_l → T]>,
  T^{1/2}, CF[{xi_i, xi_j} · (T^{-1} 0 / 1 - T^{-1} 1) · {x_k, x_l}]];
Γ[Xi_j, j, k, l, S_] := Γ[Xi_j, j, k, l, T_i, T_l];
Γ[Xi_j, j, k, l, S_] := Γ[Xi_j, j, k, l, T_i, T_j];
Γ[P_i, j, T_] := Γ[{i}, {j}, <[xi_i → T, x_j → T]>, 1, xi_i x_j];
Γ[P_i, j, S_] := Γ[P_i, j, T_i];

```

## 6. An Implementation of $\Gamma$ .

If I didn't implement I wouldn't believe myself.

Written in Mathematica [Wo], available as the notebook Gamma.nb at <http://drorbn.net/mo21/ap>. Code lines are highlighted in grey, demo lines are plain. We start with canonical forms for quadratics with rational function coefficients:

```

CCF[ξ_] := Factor[ξ];
CF[ξ_] := Module[{vs = Union@Cases[ξ, {ξ | x}_, ∞]},
  Total[(CCF[#][2]] (Times @@ vs^{#1})) & /@ CoefficientRules[ξ, vs]];

```

Contractions:

```

ch, t_ @ Γ[is_, os_, cs_, ω_, λ_] := Module[{α, η, y, μ},
  α = D[ξ_t, x_h λ]; μ = λ /. ξ_t | x_h → 0;
  η = D[x_h λ /. ξ_t → 0; y = D[ξ_t λ /. x_h → 0;
  Γ[
    DeleteCases[is, t], DeleteCases[os, h], KeyDrop[cs, {x_h, ξ_t}],
    CCF[(1 - α) ω], CF[μ + η y / (1 - α)]
  ] /. If[MatchQ[cs[ξ_t], t_], cs[ξ_t] → cs[x_h], cs[x_h] → cs[ξ_t]];
  c @ Γ[is_, os_, cs_, ω_, λ_] := Fold[c_{#2, #2}[#1] &, Γ[is, os, cs, ω, λ], is ∩ os]

```

Automatic intelligent contractions:

```

Γ[{γ_ T_}] := c[γ];
Γ[{γ1 T_, γ2 T_}] := Module[{γ2},
  γ2 = First@MaximalBy[{γs}, Length[γ1[[1]] ∩ #[[2]]] + Length[γ1[[2]] ∩ #[[1]]] &];
  Γ[Join[{c[γ1 γ2]}, DeleteCases[{γs}, γ2]]];
Γ[os_List] := Γ[Γ /@ os]

```