**So What?** If V is a representation, then  $V^{\otimes n}$  explodes as a function of *n*, while in **DoPeGDO** up to a fixed power of  $\epsilon$ , the ranks of mor( $A \rightarrow B$ ) grow polynomially as a function of |A| and |B|.

**Compositions.** In  $mor(A \rightarrow B)$ ,  $Q = \sum_{i \in A, j \in B} E_{ij} \zeta_i z_j + \frac{1}{2} \sum_{i, j \in A} F_{ij} \zeta_i \zeta_j + \frac{1}{2} \sum_{i, j \in B} G_{ij} z_i z_j,$ and so (remember,  $e^x = 1 + x + xx/2 + xxx/6 + ...)$ A  $\omega_1$ B $\omega_2$ A ω CE $E_2$  $E_1$ A.  $Q_1$  $O_2$ Q  $E_1E_2 + E_1F_2G_1E_2$ G G  $+E_1F_2G_1F_2G_1E_2$  $=\sum_{r=0}^{\infty} E_1 (F_2 G_1)^r E_2$ greek greek latin latin greek latin where •  $E = E_1(I - F_2G_1)^{-1}E_2$ . •  $F = F_1 + E_1 F_2 (I - G_1 F_2)^{-1} E_1^T$ . •  $G = G_2 + E_2^T G_1 (I - F_2 G_1)^{-1} E_2.$ dat(I  $E(C)^{-1}$ 

messy PDE or using "connected Feynman diagrams" (yet we're still in pure algebra!). Docility is preserved.

**DoPeGDO Footnotes.** Each variable has a "weight"  $\in \{0, 1, 2\}$ , and always wt  $z_i$  + wt  $\zeta_i$  = 2.

of a

- †1. Really, "weight-graded finite sets"  $A = A_0 \sqcup A_1 \sqcup A_2$ .
- $\dagger 2$ . Really, a power series in the weight-0 variables<sup> $\dagger 5$ </sup>.
- †3. The weight of Q must be 2, so it decomposes as Q = $Q_{20}+Q_{11}$ . The coefficients of  $Q_{20}$  are rational numbers while the coefficients of  $Q_{11}$  may be weight-0 power series<sup>†5</sup>.
- †4. Setting wt  $\epsilon = -2$ , the weight of P is  $\leq 2$  (so the powers of the weight-0 variables are not constrained)<sup> $\dagger 5$ </sup>.
- <sup>†5</sup>. In the knot-theoretic case, all weight-0 power series are rational functions of bounded degree in the exponentials of the weight-0 variables.
- †6. There's also an obvious product

$$\operatorname{mor}(A_1 \to B_1) \times \operatorname{mor}(A_2 \to B_2) \to \operatorname{mor}(A_1 \sqcup A_2 \to B_1 \sqcup B_2).$$

• A 1-1 phase over the ring of power series in the weight-0 variables, in which the weight-2 variables are spectators.

• A (slightly modified) 2-0 phase over  $\mathbb{Q}$ , in which the weight-1 variables are spectators.

Analog. Solve Ax = a, B(x)y = b

**Questions.** • Are there QFT precedents for "two-step Gaussian integration"?

• In QFT, one saves even more by considering "one-particleirreducible" diagrams and "effective actions". Does this mean anything here?

• Understanding Hom( $\mathbb{Q}[z_A] \to \mathbb{Q}[z_B]$ ) seems like a good cause. Can you find other applications for the technology here?

 $\mathcal{U}QU = \mathcal{U}_{\hbar}(sl_{2+}^{\epsilon}) = A\langle y, b, a, x \rangle \llbracket \hbar \rrbracket$  with  $[a, x] = x, [b, y] = -\epsilon y, [a, b] = 0, \gamma$  $[a, y] = -y, [b, x] = \epsilon x$ , and  $xy - qyx = (1 - AB)/\hbar$ , where  $q = e^{\hbar \epsilon}$ ,  $A = e^{-\hbar \epsilon a}$ , and  $B = e^{-\hbar b}$ . Also  $\Delta(y, b, a, x) = (y_1 + B_1 y_2, b_1 + b_2, a_1 + a_2, x_1 + A_1 x_2)$ ,  $S(y, b, a, x) = (-B^{-1}y, -b, -a, -A^{-1}x)$ , and  $R = \sum \hbar^{j+k} y^k b^j \otimes a^j x^k / j! [k]_q!$ .

**Theorem.** Everything of value regrading U = CU and/or its quantization U = QU is **DoPeGDO**:



also Cartan's  $\theta$ , the Dequantizator, and more, and all of their compositions.



There are lots of poly-time-computable well-**Conclusion.** behaved near-Alexander knot invariants: • They extend to tangles with appropriate multiplicative behaviour. • They have cabling and strand reversal formulas.  $\omega \epsilon \beta / akt$ The invariant for  $sl_{2\perp}^{\epsilon}/(\epsilon^2 = 0)$  (prior art:  $\omega \epsilon \beta / Ov$ ) attains 2,883 distinct values on the 2,978 prime knots with  $\leq$  12 crossings. HOMFLY-PT and Khovanov homology together attain only 2,786 distinct values.

knot	$n_{L}^{t}$ Alexander's $\omega^{+}$ genus / ribbon	knot	$n_{t}^{t}$ Alexander's $\omega^{+}$ gen	us / ribbon	knot	$n_{l}^{t}$ Alexander's $\omega^{+}$	genus / ribbon	
diag	$(\rho'_1)^+$ unknotting # / amphi?	diag	$(\hat{\rho}'_1)^+$ unknotting	#/amphi?	diag	$(\hat{\rho}'_1)^+$ unknow	otting # / amphi?	
	$(\rho'_2)^+$		$(\rho'_2)^+$			$(\rho'_2)^+$		
$\bigcirc$	$0_1^a$ 1 0 / $\checkmark$		$3_1^a T - 1$	1 / X	$(\Omega)$	$4_1^a  3-T$	1 / 🗙	
	0 0 / 🗸	P	Т	1 / 🗙	× ey	0	1 / 🖌	
_	$0$ $3T^3 - 12T^2 + 26T - 38$				$T^4 - 3T^3 - 15T^2 + 74T - 110$			
A	$5^a_1 T^2 - T + 1$ 2 / X	0	$5^{a}_{2}$ 2T-3	1 / 🗙	0	$6^{a}_{1}$ 5-2T	1 / 🗸	
8J	$2T^3+3T$ 2/X		5T - 4	1 / 🗙	62	T-4	1 / 🗙	
$5T^7 - 20T^6 + 55T^5 - 120T^4 + 217T^3 - 338T^2 + 450T - 510$			$-10T^4 + 120T^3 - 487T^2 + 1054T - 1362$			$14T^4 - 16T^3 - 293T^2 + 1098T - 1598$		
Æ	$6_2^a - T^2 + 3T - 3$ 2 / X	<i>A</i>	$6^a_3 T^2 - 3T + 5$	2/×	A.	$7^a_1$ $T^3 - T^2 + T - 1$	3 / 🗙	
88	$T^{3}-4T^{2}+4T-4$ 1 / X	S	0	1 / 🖌	ЪS	$3T^5 + 5T^3 + 6T$	3 / 🗙	
$3T^8 - 21T^7 + 49T^6 + 15T^5 - 433T^4 + 1543T^3 - 3431T^2 + 5482T - 6410$			$4T^8 - 33T^7 + 121T^6 - 203T^5 - 111T^4 + 1499T^3 - 4210T^2 + 7186T - 8510$			$7T^{11} - 28T^{10} + 77T^9 - 168T^8 + 322T^7 - 560T^6 + 891T^5 - 1310T^4 + \\$		
					$1777T^3 - 2238T^2 + 2604T - 2772$			

Video and more at http://www.math.toronto.edu/~drorbn/Talks/Columbia-191125/

Full DoPeGDO. Compute compositions in two phases: