

2. The “ $z_i \rightarrow z_j$ variable rename map $\sigma_j^i: S(z_i) \rightarrow S(z_j)$ becomes ${}^t\sigma_j^i = \oplus_{\zeta_i}^{\zeta_j}$, and it’s easy to rename several variables simultaneously.
3. The “archetypal multiplication map $m_k^{ij}: S(z_i, z_j) \rightarrow S(z_k)$ ” has ${}^t m = \oplus_{\zeta_k}^{\zeta_i + \zeta_j}$.
4. The “archetypal coproduct $\Delta_{jk}^i: S(z_i) \rightarrow S(z_j, z_k)$ ”, given by $z_i \rightarrow z_j + z_k$ or $\Delta z = z \otimes 1 + 1 \otimes z$, has ${}^t \Delta = \oplus_{\zeta_j + \zeta_k}^{\zeta_i}$.
5. R -matrices tend to have terms of the form $\oplus_q^{\hbar y_1 x_2} \in \mathcal{U}_q \otimes \mathcal{U}_q$. The “baby R -matrix” is ${}^t R = \oplus^{\hbar y x} \in S(y, x)$.


Proposition. If $F: S(B) \rightarrow S(B')$ is linear and “continuous”, then ${}^t F = \exp(\sum_{z_i \in B} \zeta_i z_i) // F$.

The Heisenberg Example. The “Weyl form of the canonical commutation relations” states that if $[y, x] = t$ and t is central, then $\oplus^{\xi x} \oplus^{\eta y} = \oplus^{\eta y} \oplus^{\xi x} \oplus^{-\eta \xi t}$. Thus with

$$SW_{xy} \begin{array}{c} \textcircled{S}(t, y, x) \\ \xrightarrow{\quad \oplus_{xy} \quad} \\ \textcircled{U}(t, y, x) \\ \xleftarrow{\quad \oplus_{yx} \quad} \end{array}$$

we have ${}^t SW_{xy} = \oplus^{\tau t + \eta y + \xi x - \eta \xi t}$.

The Zipping Issue (between unbound and bound lies half-zipped).



Zipping. If $P(\zeta^j, z_i)$ is a polynomial, or whenever otherwise convergent, set

$$\langle P(\zeta^j, z_i) \rangle_{(\zeta^j)} = P(\partial_{z_j}, z_i) \Big|_{z_i=0}.$$

(E.g., if $P = \sum a_{nm} \zeta^n z^m$ then $\langle P \rangle_\zeta = \sum n! a_{nn}$).

The Zipping / Contraction Theorem. If P has a finite ζ -degree and the y 's and the q 's are “small” then

$$\langle P(z_i, \zeta^j) e^{\eta^i z_i + y_j \zeta^j} \rangle_{(\zeta^j)} = \langle P(z_i + y_i, \zeta^j) e^{\eta^i (z_i + y_i)} \rangle_{(\zeta^j)},$$

(proof: replace $y_j \rightarrow \hbar y_j$ and test at $\hbar = 0$ and at ∂_\hbar , and

$$\begin{aligned} & \langle P(z_i, \zeta^j) \oplus^{c+\eta^i z_i + y_j \zeta^j + q_j^i z_i \zeta^j} \rangle_{(\zeta^j)} \\ &= \det(\tilde{q}) \langle P(\tilde{q}_i^k (z_k + y_k), \zeta^j) \oplus^{c+\eta^i \tilde{q}_i^k (z_k + y_k)} \rangle_{(\zeta^j)} \end{aligned}$$

where \tilde{q} is the inverse matrix of $1 - q$: $(\delta_j^i - q_j^i) \tilde{q}_k^j = \delta_k^i$ (proof: replace $q_j^i \rightarrow \hbar q_j^i$ and test at $\hbar = 0$ and at ∂_\hbar).

Implementation. [omegaBeta/ZipBindDemo](#)

```
Kδ /: Kδ[i == j, 1, 0];
{z*, x*, y*} = {ξ, ε, η}; {ξ*, ε*, η*} = {z, x, y};
(u_i)* := (u*)_i;
Zip[P_] := P;
Zip[ξ, ss__][P_] :=
(Expand[P // Zip[ss]] /. f_. ξ^d_. :> ∂{ξ^*, d}f) /. ξ* → 0
Zip[ξ][(a ξ^6 + ξ + 3) (z^5 e^z + 7 z) + 99 b]
7 + 720 a + 99 b
Zip[ξ, η][ξ^3 η^3 e^a x + b y + c x y]
a^3 b^3 + 9 a^2 b^2 c + 18 a b c^2 + 6 c^3
(* IE [Q,P] means e^Q P *)
IE /: Zip[ss_List]@IE[Q_, P_] :=
Module[{ξ, z, zs, c, ys, ηs, qt, zrule, Q1, Q2},
zs = Table[ξ^*, {ξ, ss}];
c = Q /. Alternatives @@ (ss ∪ zs) → 0;
ys = Table[∂ξ(Q /. Alternatives @@ zs → 0), {ξ, ss}];
ηs = Table[∂z(Q /. Alternatives @@ ss → 0), {z, zs}];
qt = Inverse@Table[Kδ[z, ξ^*] - ∂z, ξ Q, {ξ, ss}, {z, zs}];
zrule = Thread[zs → qt.(zs + ys)];
Q1 = c + ηs.zs /. zrule;
Q2 = Q1 /. Alternatives @@ zs → 0;
Simplify /@ IE[Q2, Det[qt] e^-Q2 Zip[ss[e^Q1 (P /. zrule)]]];
```

$$Eh = \mathbb{E} \left[h \sum_{i=1}^3 \sum_{j=1}^3 a_{10} \xi_{i+j} x_i \xi_j, \sum_{i=1}^3 f_i[x_1, x_2, x_3] \xi_i \right];$$

$$E1 = Eh / . h \rightarrow 1$$

$$\begin{aligned} & \mathbb{E} [a_{11} x_1 \xi_1 + a_{21} x_2 \xi_1 + a_{31} x_3 \xi_1 + a_{12} x_1 \xi_2 + \\ & a_{22} x_2 \xi_2 + a_{32} x_3 \xi_2 + a_{13} x_1 \xi_3 + a_{23} x_2 \xi_3 + a_{33} x_3 \xi_3, \\ & \xi_1 f_1[x_1, x_2, x_3] + \xi_2 f_2[x_1, x_2, x_3] + \xi_3 f_3[x_1, x_2, x_3]] \end{aligned}$$

$$Short[lhs = Zip[\xi_1, \xi_2] @ E1, 5]$$

$$\begin{aligned} & \mathbb{E} \left[((a_{13} ((-1 + a_{22}) a_{31} - a_{21} a_{32}) + a_{12} (-a_{23} a_{31} + a_{21} a_{33}) + \right. \\ & \left. (-1 + a_{11}) (a_{23} a_{32} - (-1 + a_{22}) a_{33})) x_3 \xi_3) / \right. \\ & \left. \frac{(-1 + a_{12} a_{21} - a_{11} (-1 + a_{22}) + a_{22})}{(-1 + a_{12} a_{21} - a_{11} (-1 + a_{22}) + a_{22})^2} \right] \\ Lhs &== Zip[\xi_1] @ Zip[\xi_2] @ E1 == Zip[\xi_2] @ Zip[\xi_1] @ E1 \end{aligned}$$

True

Short[

$$lhs = Normal[Eh /. IE[Q_, P_] :> Series[P e^Q, {h, 0, 3}]] // Zip[\xi_1, \xi_2], 5]$$

$$\begin{aligned} & h a_{13} \xi_3 f_1[0, 0, x_3] + 2 h^2 a_{11} a_{13} \xi_3 f_1[0, 0, x_3] + \\ & 3 h^3 a_{11}^2 a_{13} \xi_3 f_1[0, 0, x_3] + 2 h^3 a_{12} a_{13} a_{21} \xi_3 f_1[0, 0, x_3] + \\ & h^2 a_{13} a_{22} \xi_3 f_1[0, 0, x_3] + <> 337 >> + \\ & \frac{1}{6} h^3 a_{31}^3 x_3^3 \xi_3 f_3^{(3, 0, 0)}[0, 0, x_3] + \frac{1}{2} h^3 a_{31}^2 a_{32} x_3^3 f_1^{(3, 1, 0)}[0, 0, x_3] + \\ & \frac{1}{6} h^3 a_{31}^3 x_3^3 f_2^{(3, 1, 0)}[0, 0, x_3] + \frac{1}{6} h^3 a_{31}^3 x_3^3 f_1^{(4, 0, 0)}[0, 0, x_3] \end{aligned}$$

rhs =

$$Normal[Zip[\xi_1, \xi_2] @ Eh /. IE[Q_, P_] :> Series[P e^Q, {h, 0, 3}]];$$

Simplify[lhs == rhs]

True

$$E /: E[Q1_, P1_] E[Q2_, P2_] := E[Q1 + Q2, P1 * P2];$$

```
Bind[ss_List][L_E, R_E] := Module[{n, hideξs, hidezs},
  hideξs = Table[ξs[[i]] → ξs[[i], {i, Length@ξs}]];
  hidezs = Table[ξs[[i]]* → ξs[[i], {i, Length@ξs}]];
  Zip[ξs, hideξs] [(L /. hideξs) (R /. hidezs)]];
```

Bind[ξs][IE[ξ(x1 + x2), 1], IE[ξ2(x2 + x3), 1]]

IE[ξ(x1 + x2 + x3), 1]

Bind[ξs][IE[(ξ2 + ξ3)x2, 1], IE[(ξ1 + ξ2)x, 1]]

IE[x(ξ1 + ξ2 + ξ3), 1]

The 2D Lie Algebra. Clever people know* that if $[a, x] = \gamma x$ then $\oplus^{\xi x} \oplus^{aa} = \oplus^{aa} \oplus^{e^{-\gamma a} \xi x}$. Ergo with

$$SW_{ax} \begin{array}{c} \textcircled{S}(a, x) \\ \xrightarrow{\quad \oplus_{ax} \quad} \\ \textcircled{U}(a, x) \\ \xleftarrow{\quad \oplus_{xa} \quad} \end{array}$$

we have ${}^t SW_{ax} = \oplus^{aa + e^{-\gamma a} \xi x}$.

* Indeed $xa = (a - \gamma)x$ thus $xa^n = (a - \gamma)^n x$ thus $x \oplus^{aa} = e^{a(a-\gamma)} x = e^{-\gamma a} e^{aa} x$ thus $x^n \oplus^{aa} = e^{aa} (e^{-\gamma a})^n x^n$ thus $e^{\xi x} \oplus^{aa} = e^{aa} \oplus^{e^{-\gamma a} \xi x}$.

The Real Thing. In $QU/(\epsilon^2 = 0)$ over $\mathbb{Q}[[\hbar]]$ using the yax order, $T = \oplus^{\hbar t}$, $\bar{T} = T^{-1}$, $\mathcal{A} = \oplus^{\gamma a}$, and $\bar{\mathcal{A}} = \mathcal{A}^{-1}$, we have

$$R_{ij} = e^{\hbar(y_i x_j - t_i a_j)/\gamma} (1 + \epsilon \hbar (a_i a_j / \gamma - \gamma \hbar^2 y_i^2 x_j^2 / 4))$$

in $S(B_i, B_j)$, and in $S(B_1^*, B_2^*, B)$ we have

$$t_m = e^{(a_1 + a_2)a + \eta_2 \xi_1 (1-T)/\hbar + (\xi_1 \bar{\mathcal{A}}_2 + \xi_2 \mathcal{A}_2)x + (\eta_1 + \eta_2 \bar{\mathcal{A}}_1)y} (1 + \epsilon \lambda_m),$$

where $\lambda_m = 2a\eta_2 \xi_1 T + \frac{1}{4}\gamma\eta_2^2 \xi_1^2 (3T^2 - 4T + 1)/\hbar - \frac{1}{2}\gamma\eta_2 \xi_1^2 (3T - 1)x \bar{\mathcal{A}}_2 - \frac{1}{2}\gamma\eta_2^2 \xi_1 (3T - 1)y \bar{\mathcal{A}}_1 + \gamma\eta_2 \xi_1 xy \hbar \bar{\mathcal{A}}_1 \bar{\mathcal{A}}_2$. Similar formulas delight us for ${}^t \Delta$ and ${}^t S$.

A generic morphism.

$$\begin{array}{ccc} L: & \begin{array}{c} t \quad t \\ | \qquad | \\ \textcircled{S} \\ | \qquad | \\ a \quad a \end{array} & Q: \begin{array}{c} x \quad x \quad x \\ \swarrow \quad \searrow \quad \swarrow \\ \textcircled{S} \\ \searrow \quad \swarrow \quad \searrow \\ \alpha \quad \alpha \quad \alpha \end{array} \\ & \xrightarrow{\quad \oplus_{ax} \quad} & \\ & \begin{array}{c} x \quad a \quad x \\ | \qquad | \qquad | \\ \textcircled{U} \\ | \qquad | \qquad | \\ a \quad x \quad a \end{array} & P: \begin{array}{c} x \quad a \quad x \\ | \qquad | \qquad | \\ \textcircled{U} \\ | \qquad | \qquad | \\ a \quad x \quad \xi \end{array} \end{array}$$