


| | |
|--|--|
| <p>Dror Bar-Natan: Talks: Greece-1607: oeβ:=http://drorbn.net/Greece-1607/</p> <p>Work in Progress! The Brute and the Hidden Paradise</p> <p>Local Algebra (with van der Veen) Much can be reformulated as (non-standard) “quantum algebra” for the 4D Lie algebra $\mathfrak{g} = \langle b, c, u, w \rangle$ over $\mathbb{Q}[\epsilon]/(\epsilon^2 = 0)$, with b central and $[w, c] = w$, $[c, u] = u$, and $[u, w] = b - 2\epsilon c$. The key: $a_{ij} = (b_i - \epsilon c_i)c_j + u_i w_j$ in $\mathcal{U}(\mathfrak{g})^{\otimes \{i,j\}}$.</p> <p></p> <p>van der Veen</p> | <p>(bas // TG_{1,2} // TG_{1,3}) - (bas // TG_{1,3} // TG_{1,2}) ... OC</p> <p>$\{0, -f[t_1, t_2, t_3] u_1 u_2 w_3 + f[t_1, t_2, t_3] t_1 u_1 u_2 w_3 + f[t_1, t_2, t_3] u_1 u_3 w_3 - f[t_1, t_2, t_3] t_1 u_1 u_3 w_3, -f[t_1, t_2, t_3] u_1 u_2 w_2 + f[t_1, t_2, t_3] t_1 u_1 u_2 w_2 + f[t_1, t_2, t_3] u_1 u_3 w_2 - f[t_1, t_2, t_3] t_1 u_1 u_3 w_2, 0, 0, 0, 0, 0, 0\}$</p> <p>$\eta /: \eta[i] = 0; \eta /: \eta[i] \eta[j] = 0;$ Turbo-Burau (new!)</p> |
| <p>Some (new) representations of the (v-)braid groups. oeβ/Reps</p> <p>B_{i,j} [ε] := ε /. v_j ↪ (1 - t) v_i + t v_j Burau (old)</p> <p>Column@{lhs = {v₁, v₂, v₃} // B_{1,2} // B_{1,3} // B_{2,3}, rhs = {v₁, v₂, v₃} // B_{2,3} // B_{1,3} // B_{1,2}, lhs - rhs // Expand}</p> <p>{v₁, (1 - t) v₁ + t v₂, (1 - t) v₁ + t ((1 - t) v₂ + t v₃)}</p> <p>{v₁, (1 - t) v₁ + t v₂, (1 - t) ((1 - t) v₁ + t v₂) + t ((1 - t) v₁ + t v₃)}</p> <p>{0, 0, 0}</p> <p>G_{i,j} [ε] := ε /. v_j ↪ (1 - t_i) v_i + t_i v_j Gassner (old)</p> <p>... Overcrossings Commute (OC):</p> | <p>TB_{i,j} [ε] :=</p> <p>Expand[ε /. f_k ↪ Plus[f_{v_k} /. v_j → (1 - t - η[i]) v_i + (t + η[i]) v_j, (t - 1) (Coefficient[f, η[i]] - Coefficient[f, η[j]]) * (u_k /. u_j → (1 - t) u_i + t u_j) * u_i w_j, Kδ_{k,i} (f /. _η → 0) (u_j - u_i) u_i w_j, u_j → (1 - t) u_i + t u_j, w_i → w_i + (1 - t⁻¹) w_j, w_j → t⁻¹ w_j]];</p> <p>ff = f₀ + f₁ η[1] + f₂ η[2] + f₃ η[3];</p> <p>bas = {ff v₁, ff v₂, ff v₃, u₁^{2 w₁, u₁² w₂, u₁, u₂, u₃, w₁, w₂, w₃};}</p> <p>(bas // TB_{1,2} // TB_{1,3}) - (bas // TB_{1,3} // TB_{1,2}) ... OC</p> <p>{0, -f₀ u₁ u₂ w₃ + t f₀ u₁ u₂ w₃ + f₀ u₁ u₃ w₃ - t f₀ u₁ u₃ w₃, -f₀ u₁ u₂ w₂ + t f₀ u₁ u₂ w₂ + f₀ u₁ u₃ w₂ - t f₀ u₁ u₃ w₂, 0, 0, 0, 0, 0, 0, 0}</p> |
| <p>Column@{lhs = {v₁, v₂, v₃} // G_{1,2} // G_{1,3}, Expand[lhs - ({v₁, v₂, v₃} // G_{1,3} // G_{1,2})]}]</p> <p>{v₁, (1 - t₁) v₁ + t₁ v₂, (1 - t₁) v₁ + t₁ v₃}</p> <p>{0, 0, 0}</p> <p>... Undercrossings Commute (UC):</p> | <p>Flower Surgery Theorem. A knot is ribbon iff it is the result of n-petal flower surgery (from thin petals to wide petals) on an n-component unlink, for some n.</p> <p> Colin, you happy?</p> |
| <p>Column@{lhs = {v₁, v₂, v₃} // G_{1,3} // G_{2,3}, rhs = {v₁, v₂, v₃} // G_{2,3} // G_{1,3}, lhs - rhs // Expand}]</p> <p>{v₁, v₂, (1 - t₁) v₁ + t₁ ((1 - t₂) v₂ + t₂ v₃)}</p> <p>{v₁, v₂, (1 - t₂) v₂ + t₂ ((1 - t₁) v₁ + t₁ v₃)}</p> <p>{0, 0, v₁ - t₁ v₁ - t₂ v₁ + t₁ t₂ v₁ - v₂ + t₁ v₂ + t₂ v₂ - t₁ t₂ v₂}</p> <p>Gassner Plus (new?)</p> <p>GP_{i,j} [ε] := Expand[ε /. {u_j ↪ (1 - t_i) u_i + t_i u_j, f_{v_k} ↪ f (1 - t_i) v_i + f t_i v_j + (t_i - 1) (t_i ∂_{t_i} f - t_j ∂_{t_j} f) u_i + f t_i u_i }];</p> <p>bas = {f[t₁, t₂, t₃] v₁, f[t₁, t₂, t₃] v₂, f[t₁, t₂, t₃] v₃, u₁, u₂, u₃};</p> <p>Short[lhs = bas // GP_{1,2} // GP_{1,3} // GP_{2,3}, 2] ... R3 (left)</p> <p>{f[t₁, t₂, t₃] v₁, f[t₁, t₂, t₃] t₁ u₁ + f[t₁, t₂, t₃] v₁ - f[t₁, t₂, t₃] t₁ v₁ + <<6>> + t₁² u₁ f^(1,0,0) [t₁, t₂, t₃], <<1>> + <<19>> + <<1>>, <<1>>, u₁ - t₁ u₁ + t₁ u₂, u₁ - t₁ u₁ + t₁ u₂ - t₁ t₂ u₂ + t₁ t₂ u₃}</p> <p>(bas // GP_{2,3} // GP_{1,3} // GP_{1,2}) - lhs ... R3 (rest)</p> <p>{0, 0, 0, 0, 0, 0}</p> <p>(bas // GP_{1,2} // GP_{1,3}) - (bas // GP_{1,3} // GP_{1,2}) ... OC</p> <p>{0, 0, 0, 0, 0, 0}</p> <p>Question. Does Gassner Plus factor through Gassner?</p> | <p>References.</p> <p>[BN] D. Bar-Natan, <i>Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant</i>, oeβ/KBH, arXiv: 1308.1721.</p> <p>[BND] D. Bar-Natan and Z. Dancso, <i>Finite Type Invariants of W-Knotted Objects I, II, IV</i>, oeβ/WKO1, oeβ/WKO2, oeβ/WKO4, arXiv:1405.1956, arXiv: 1405.1955, arXiv:1511.05624.</p> <p>[BNG] D. Bar-Natan and S. Garoufalidis, <i>On the Melvin-Morton-Rozansky conjecture</i>, Invent. Math. 125 (1996) 103–133.</p> <p>[BNS] D. Bar-Natan and S. Selmani, <i>Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial</i>, J. of Knot Theory and its Ramifications 22-10 (2013), arXiv:1302.5689.</p> <p>[En] B. Enriquez, <i>A Cohomological Construction of Quantization Functors of Lie Bialgebras</i>, Adv. in Math. 197-2 (2005) 430-479, arXiv:math/0212325.</p> <p>[EK] P. Etingof and D. Kazhdan, <i>Quantization of Lie Bialgebras, I</i>, Selecta Mathematica 2 (1996) 1–41, arXiv:q-alg/9506005.</p> <p>[GPV] M. Goussarov, M. Polyak, and O. Viro, <i>Finite type invariants of classical and virtual knots</i>, Topology 39 (2000) 1045–1068, arXiv: math.GT/9810073.</p> <p>[Ha] A. Haviv, <i>Towards a diagrammatic analogue of the Reshetikhin-Turaev link invariants</i>, Hebrew University PhD thesis, Sep. 2002, arXiv: math.QA/0211031.</p> <p>[MM] P. M. Melvin and H. R. Morton, <i>The coloured Jones function</i>, Commun. Math. Phys. 169 (1995) 501–520.</p> <p>[PV] M. Polyak and O. Viro, <i>Gauss Diagram Formulas for Vassiliev Invariants</i>, Inter. Math. Res. Notices 11 (1994) 445–453.</p> <p>[Ro] L. Rozansky, <i>A contribution of the trivial flat connection to the Jones polynomial and Witten’s invariant of 3d manifolds, I</i>, Comm. Math. Phys. 175-2 (1996) 275–296, arXiv:hep-th/9401061.</p> <p>[Se] P. Ševera, <i>Quantization of Lie Bialgebras Revisited</i>, Sel. Math., NS, to appear, arXiv:1401.6164.</p> <p> “God created the knots, all else in topology is the work of mortals.” Leopold Kronecker (modified)</p> <p>www.katlas.org </p> |

Video and more at <http://www.math.toronto.edu/~drorbn/Talks/Greece-1607/>