

(u, v, and w knots) x (topology, combinatorics, low algebra, and high algebra)

Dror Bar-Natan, Kansas State April 7 2009, <http://www.math.toronto.edu/~drorbn/Talks/KSU-090407>

"God created the knots, all else in topology is the work of mortals."
Leopold Kronecker (modified)

1	<p>u-knots</p> <p>u-knots are usual knots:</p> <p>=PA $\langle \text{R123} \rangle_0$ legs Reidemeister</p> <p>"Knots in \mathbb{R}^3"</p>	1+1	<p>v-knots</p> <p>v-knots are virtual knots:</p> <p>=PA $\langle \text{R123} \rangle_0$</p> <p>=CA $\langle \text{R123} \rangle_0$</p> <p>= Knots on surfaces, modulo stabilization:</p> <p>Kauffman</p>	onto	<p>w-knots</p> <p>w is for welded, weakly v, and warmup:</p> <p>$\{w\text{-knots}\} = \{v\text{-knots}\} / (\text{OC})$</p> <p>where OC is Overcrossings Commute:</p> <p>yet \neq</p> <p>Related to "movies of flying rings" to knotted tubes in 4-space, and to "basis conjugating automorphisms of free groups".</p> <p>McCool Goldsmith Fenn Rimanyi Rourke Satoh Brendle Hatcher</p>
topology	<p>Extend any $V : \{u\text{-knots}\} \rightarrow \mathcal{A}$ to "singular u-knots" using $V(\bowtie) := V(\times) - V(\times)$, and think "differentiation".</p> <p>Declare "V is of type m" iff $V^{(m+1)} \equiv 0$, think "polynomial of degree m".</p> <p>$W = V^{(m)}$ roughly determines V; $W \in \mathcal{A}_m^* = (\mathcal{K}_m / \mathcal{K}_{m+1})^*$ with</p> <p>$\mathcal{A}_m := \left\{ \begin{array}{c} \text{diagram with } m \text{ chords} \end{array} \right\} / \sim$</p> <p>Need an expansion $Z : \{u\text{-knots}\} \rightarrow \mathcal{A} = \bigoplus \mathcal{A}_m$.</p> <p>Vassiliev Goussarov</p>	combinatorics	<p>All the same, except</p> <p>$V(\bowtie) := V(\times) - V(\times)$</p> <p>$V(\bowtie) := V(\times) - V(\times)$</p> <p>$\mathcal{A}^v := \{ \text{"arrow diagrams"} \} / 6T$</p> <p>Need a $Z : \{v\text{-knots}\} \rightarrow \mathcal{A}^v$.</p> <p>The 6T Relation (and a hidden 4T):</p>	6	<p>All the same, except</p> <p>$\mathcal{A}^w := \mathcal{A}^v / TC$</p> <p>Need a $Z : \{w\text{-knots}\} \rightarrow \mathcal{A}^w$.</p> <p>"Tails Commute (TC)":</p>
low algebra	<p>Similar</p> <p>with metrized Lie algebras replacing arbitrary Lie algebras</p> <p>Penrose Cvitanovic Vogel</p>	10	<p>Similar</p> <p>with Lie bi-algebras replacing arbitrary Lie algebras</p> <p>Haviv Leung</p>	9	<p>Theorem. $\mathcal{A}^w \cong \mathcal{A}^{wt} :=$</p> <p>&TC</p> <p>This screams, if you speak the language, LIE ALGEBRAS. And indeed we have</p> <p>Theorem. Given a finite dimensional Lie algebra \mathfrak{g}, there is $T : \mathcal{A}^w \rightarrow \mathcal{U}(I\mathfrak{g}) := \mathcal{U}(\mathfrak{g} \ltimes \mathfrak{g}_{ab}^*)$.</p>
high algebra	<p>Knots are the wrong objects to study in knot theory! They are not finitely generated and they carry no interesting operations.</p> <p>11</p> <p>Knotted Trivalent Graphs</p> <p>Theorem (~). A homomorphic Z is the same as a "Drinfel'd Associator".</p> <p>Drinfel'd</p>	11	<p>13</p> <p>Z is a Quantum Group?</p> <p>More precisely, a homomorphic Z ought to be equivalent to the Etingof-Kazhdan theory of deformation quantization of Lie bialgebras.</p> <p>Etingof Kazhdan</p> <p>Dror's Dream: Straighten and fatten this column.</p> <p>An Idle Question.</p> <p>Is there physics in this column?</p>	13	<p>Switch to w-knotted trivalent tangles, 12</p> <p>wKTT := $CA \langle \bowtie, \times, Y \rangle$.</p> <p>Theorem (~). A homomorphic Z is equivalent to proving the Kashiwara-Vergne statement.</p> <p>Statement (~, KV, 1978) (proven Alekseev-Meinrenken, 2006). Convolutions of invariant functions on a group match with convolutions of invariant functions on its Lie algebra: for any finite dim. Lie group G with Lie algebra \mathfrak{g},</p> <p>$(\text{Fun}(G)^{\text{Ad } G}, \star) \cong (\text{Fun}(\mathfrak{g})^{\text{Ad } G}, \star)$.</p> <p>(Closely related to the "orbit method" of representation theory).</p> <p>Alekseev Torossian</p>