

Dror Bar-Natan: Talks: Hanoi-0708: **Following Lin: Expansions for Groups**



Riverside, April 2000



Kyoto, September 2001

See Lin's "Power Series Expansions and Invariants of Links", 1993 Georgia International Topology Conference, AMS/IP Studies in Adv. Math. **2** (1997) 184-202.

**Vaughan's Hierarchy**  
(generalized, unauthorized)

- ☺ Computation
- ☺ Formula
- ☹ Proof
- ☹ Theory
- ☹ Dream

### The Magnus and Exponential Expansions

$$Z_{1,2} : G_n = \left( \begin{array}{c} \text{free group} \\ \text{on} \\ X_1, \dots, X_n \end{array} \right) \rightarrow \hat{A}_n = \left( \begin{array}{c} \text{completed free} \\ \text{associative} \\ \text{algebra on} \\ x_1, \dots, x_n \end{array} \right)$$

by  $X_i \mapsto 1 + x_i$  or  $e^{x_i}$

$$X_i^{-1} \mapsto 1 - x_i + x_i^2 - \dots \text{ or } e^{-x_i}.$$

**What's "An Expansion"?** A filtration-preserving isomorphism  $Z : C(G) \rightarrow \mathcal{A}(G)$  where

$$I := \{ \sum a_i g_i : \sum a_i = 0 \} \subset \mathbb{C}G$$

$$\mathbb{C}G = I^0 \supset I^1 \supset I^2 \supset I^3 \supset \dots$$

$$C(G) := \varprojlim_k \mathbb{C}G/I^k \rightarrow \dots \rightarrow \mathbb{C}G/I^2 \rightarrow \mathbb{C}G/I \rightarrow 0$$

is filtered by  $F_m C(G) := \varprojlim_{k>m} I^m/I^k$  and  $\mathcal{A}(G) := \text{gr } C(G) = \bigoplus I^m/I^{m+1}$ .

So all  
expansions  
are  
"equivalent"

**Think duals!**  $C(G)^*$  are "finite type invariants".  
 $\mathcal{A}(G)^*$  are "weight systems".  
 $Z$  is a "universal finite type invariant".

**$Z_{1,2}$  are Expansions.** With  $Z^0 = Z_1$  or  $Z^0 = Z_2$ :

- $\iota$  is automatic.
- $\rho$  is well-defined.
- $Z^0|_{I^m} \subset F_m \mathcal{A}_n$ .
- $Z^0$  descends to  $Z^1$ .
- Define  $Z^2$ .
- $\rho$  is surjective.
- $\text{gr } Z^2$  is the identity.
- $Z^2$  is an isomorphism.
- $\rho$  is an isomorphism.
- Everything generalizes, step 2 sometimes becomes tricky.

$$\begin{array}{ccc} G_n & \xrightarrow{Z^0} & \hat{A}_n \\ \downarrow \iota & \nearrow Z^1 & \downarrow \rho \\ C(G_n) & \xrightarrow{Z^2} & \mathcal{A}(G_n) \end{array}$$

$x_i \downarrow x_i^{-1}$

### The Kontsevich Integral for Braids

$$\mathcal{A} := \left\langle \begin{array}{c} \text{crossings} \\ \text{Reidemeister moves} \end{array} \right\rangle \xrightarrow{Z_K} \left\langle \begin{array}{c} \text{crossings} \\ \text{Reidemeister moves} \end{array} \right\rangle$$

modulo loc and "6T":

$$\text{loc: } \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array}$$

$$4T: \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{crossing} \\ \hline \end{array}$$



M. Kontsevich


Which other groups / groupoids / categories have expansions?



### Dror's Dream / Obsession:

"Unify" quantum groups – find one object that contains all.

**Example:** One invariant to rule them all:



Easy! Universal! A Morphism! Unique! An Isomorphism!

**What is a "Quantum Group"?** For now, a "deformation of the trivial" solution in  $\mathcal{U}(\mathfrak{g})^{\otimes*}[[\hbar]]$  of the major equations:

$$(\Delta \otimes 1)\Delta = (1 \otimes \Delta)\Delta \quad R^{-1}\Delta R = \Delta^{op}$$

$$(\Delta \otimes 1)R = R^{23}R^{13} \quad (1 \otimes \Delta)R = R^{12}R^{13}$$

(as well as a few minor equations).

**Dror's Guess:** A unified object exists; we'll need:

- Expansions as in Lin / universal finite type invariants.
- Naturality / functoriality.
- Knotted graphs, especially trivalent.
- Associators following Drinfel'd.
- The work of Etingof and Kazhdan on bialgebras.
- Virtual braids / knots / knotted graphs.
- Polyak (LMP 54) & Haviv (arXiv:math/0211031) on arrow diagrams.  
(and when construction ends, we'll dump the scaffolding)

**Why care?**  
Quantum groups  
computable  
invariants  
make!  
  
**Visit!**  
katlas.org  
**Edit!**

### (Quasi?) Natural Expansions

$G \mapsto C(G)$  and  $G \mapsto \mathcal{A}(G)$  are functors. Can you choose a ((quasi?) natural)  $Z$  satisfying

$$\begin{array}{ccc} C(G_1) & \xrightarrow{C(\Delta)} & C(G_2) \\ \downarrow Z(G_1) & & \downarrow Z(G_2) \\ \mathcal{A}(G_1) & \xrightarrow{\mathcal{A}(\Delta)} & \mathcal{A}(G_2) \end{array}$$

$\left( \begin{array}{l} \text{quasi: } \mathcal{A}'(\Delta) = \\ J_{\Delta}^{-1} \mathcal{A}(\Delta) J_{\Delta} ? \\ D_{\Delta} \mathcal{A}(\Delta) ? \end{array} \right)$

Perhaps just on a subcategory of **Groups**? Perhaps **Braids** with strands addition, deletion and doubling:

$$\begin{array}{c} \text{strand} \\ \text{addition} \end{array} \Rightarrow \begin{array}{c} \text{strand} \\ \text{deletion} \end{array} \quad \text{or} \quad \begin{array}{c} \text{strand} \\ \text{doubling} \end{array} \quad \text{or} \quad \begin{array}{c} \text{strand} \\ \text{doubling} \end{array}$$

Note the relation:

$$\begin{array}{c} \text{crossing} \\ \text{relation} \end{array} = \begin{array}{c} \text{crossing} \\ \text{relation} \end{array} \cdot \begin{array}{c} \text{crossing} \\ \text{relation} \end{array}$$

### Virtual Braids

crossings are real, strands go virtual

**Definition.**

Crossings,

modulo

Reidemeister

moves,

but the linkages between

crossings are "virtual":

two diagrams

of  $v_{21}$

L. Kauffman

M. Polyak

T. Ohtsuki

Drinfel'd

Etingof

Kazhdan

Haviv

Polyak's  $\vec{\mathcal{A}}$ .

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

modulo loc and "6T":

**Lie bialgebras.**

The  $\mathfrak{g}$  in a sum  $\mathfrak{g} \oplus \mathfrak{g}^*$  which in itself is a Lie algebra with subalgebras  $\mathfrak{g}$  and  $\mathfrak{g}^*$ , and in which the tautological metric is invariant.

**Why bother?**

Their deformations are quantum groups, and their diagrammatic universalization is  $\vec{\mathcal{A}}$ .

Drinfel'd

Etingof


Kazhdan

Haviv

Leopold Kronecker (modified)

**Question** Can you interpret quantum groups as (quasi?)-natural expansions on virtual braids?

**Dror's Guess:** No, but the effort will be worthwhile.



"God created the knots, all else in topology is the work of mortals"

Leopold Kronecker (modified)

<http://www.math.toronto.edu/~drorbn/Talks/Hanoi-0708/>; thanks to Jana Comstock, Peter Lee, Scott Morrison, Dylan Thurston.