



From Stonehenge to Witten Skipping all the Details


Oporto Meeting on Geometry, Topology and Physics, July 2004

Dror Bar-Natan, University of Toronto



It is well known that when the Sun rises on midsummer's morning over the "Heel Stone" at Stonehenge, its first rays shine right through the open arms of the horseshoe arrangement. Thus astrological lineups, one of the pillars of modern thought, are much older than the famed Gaussian linking number of two knots.

Recall that the latter is itself an astrological construct: one of the standard ways to compute the Gaussian linking number is to place the two knots in space and then count (with signs) the number of shade points cast on one of the knots by the other knot, with the only lighting coming from some fixed distant star.




The Gaussian linking number

$$lk(\bigcirc, \bigcirc) = \frac{1}{2} \sum_{\text{vertical chopsticks}} (\text{signs})$$



Carl Friedrich Gauss



$$lk=2$$

Thus we consider the generating function of all stellar coincidences:

$$Z(K) := \lim_{N \rightarrow \infty} \sum_{3\text{-valent } D} \frac{1}{2^c c! \binom{N}{e}} \langle D, K \rangle_{\overline{W}} D \cdot \left(\begin{array}{c} \text{framing-} \\ \text{dependent} \\ \text{counter-term} \end{array} \right) \in \mathcal{A}(\odot)$$

$N := \# \text{ of stars}$
 $c := \# \text{ of chopsticks}$
 $e := \# \text{ of edges of } D$

$$\mathcal{A}(\odot)$$

$:= \text{Span} \left\langle \begin{array}{c} \text{oriented vertices} \\ \text{AS: } \begin{array}{c} \text{Y} + \text{Y} = 0 \\ \text{\& more relations} \end{array} \end{array} \right\rangle$



Dylan Thurston

Theorem. Modulo Relations, $Z(K)$ is a knot invariant!

When deforming, catastrophes occur when:

A plane moves over an intersection point –
 Solution: Impose IHX,

$$I = H - X$$

(see below)

An intersection line cuts through the knot –
 Solution: Impose STU,

$$Y = U - V$$


(similar argument)

The Gauss curve slides over a star –
 Solution: Multiply by a framing-dependent counter-term.

(not shown here)

The IHX Relation

the red star is your eye.



V : vector space
 dV : Lebesgue's measure on V .
 Q : A quadratic form on V ;
 $Q(V) = \langle L^*V, V \rangle$ where
 $L: V \rightarrow V^*$ is linear
Compute $I = \int_V dV e^{\pm Q + P}$
 $= \sum_{n=0}^{\infty} \frac{1}{n!} \int_V dV P^n e^{Q/2}$
 $\sim \sum_{n=0}^{\infty} \frac{1}{n!} P^n(\partial_V) e^{-Q/2} / \int_V dV$
 $\stackrel{\pi=2\pi i}{=} \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n n! n!} P^n(a) (Q^{-1})^n / \int_V dV$

The Fourier Transform:

$$(F: V \rightarrow \mathbb{C}) \Rightarrow (F: V^* \rightarrow \mathbb{C})$$

Via $F(V) = \int_V F(V) e^{-i \langle V, V \rangle} dV$.

Simple Facts:

- $F(0) = \int_V F(V) dV$.
- $\frac{\partial}{\partial V} F \sim \widehat{V} F$.
- $(e^{Q/2}) \sim e^{-Q'/2}$
 where $Q'(V) = \langle V, L^{-1}V \rangle$
 (That's the heart of the Fourier Inversion Formula).

Differentiation and Pairings:

$$\frac{\partial^3}{\partial x^2 \partial y} x^3 y^2 = 3! 2! j \text{ indeed,}$$


$$\left\{ \begin{array}{c} \frac{\partial}{\partial x} \frac{\partial}{\partial x} \frac{\partial}{\partial x} \frac{\partial}{\partial y} \frac{\partial}{\partial y} \\ \text{X X X Y Y} \end{array} \right\}$$

$$(\lambda_{ijk} \partial_i \partial_j \partial_k)^2 (\lambda^{pqr} \psi_p \psi_q \psi_r)^3 \text{ is}$$

(2 possible)


In our case,

- Q is d , so Q^{-1} is an integral operator.
- P is $\sum A_i A_i$
- H is the holonomy, itself a sum of integrals along the knot K ,



& when the dust settles, we get $Z(K)$!

So $\int_V H(V) e^{\pm Q + P} dV$
 $\sim H(a) e^{H(a)} e^{-Q'(a)/2} / \int_V dV$
 is $\sum \begin{array}{c} \text{H} \\ \text{P} \end{array}$
 $= \sum_{\text{Diagrams}} c(D) \left(\begin{array}{c} \text{products of} \\ Q^{-1}\text{'s, } P\text{'s} \\ \text{and one } H \end{array} \right)$



Richard Feynman

It all is perturbative Chern-Simons-Witten theory:

$$\int_{\text{g-connections}} \mathcal{D}A \text{ hol}_K(A) \exp \left[\frac{ik}{4\pi} \int_{\mathbb{R}^3} \text{tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right]$$

$$\rightarrow \sum_{D: \text{Feynman diagram}} W_g(D) \oint \mathcal{E}(D) \rightarrow \sum_{D: \text{Feynman diagram}} D \oint \mathcal{E}(D)$$



Shiing-shen Chern



James H. Simons

"God created the knots,
 all else in topology
 is the work of man."



Leopold Kronecker

(modified)

This handout is at <http://www.math.toronto.edu/~drorbn/Talks/Oporto-0407>

More at <http://www.math.toronto.edu/~drorbn/Talks/Oporto-0407/>