

THE FULTON-MACPHERSON COMPACTIFICATION

DROR BAR-NATAN

Let M be a manifold and let A be a finite set.

Definition 1. The open configuration space of A in M is

$$C_A^o(M) := \{\text{injections } \iota : A \rightarrow M\}.$$

Definition 2. The compactified configuration space of A in M is

$$C_A(M) := \coprod_{\substack{\{A_1, \dots, A_k\} \\ A = \cup A_\alpha}} \left\{ \left(p_\alpha \in M, c_\alpha \in \tilde{C}_{A_\alpha}(T_{p_\alpha} M) \right)_{\alpha=1}^k : p_\alpha \neq p_\beta \text{ for } \alpha \neq \beta \right\}$$

where if V is a vector space and $|A| \geq 2$,

$$\tilde{C}_A(V) := \coprod_{\substack{\{A_1, \dots, A_k\} \\ A = \cup A_\alpha; k \geq 2}} \left\{ \left(v_\alpha \in V, c_\alpha \in \tilde{C}_{A_\alpha}(T_{v_\alpha} V) \right)_{\alpha=1}^k : v_\alpha \neq v_\beta \text{ for } \alpha \neq \beta \right\} / \text{translations and dilations.}$$

while if A is a singleton, $\tilde{C}_A(V) := \{\text{a point}\}$.

Theorem 1. (1) $C_A(M)$ is a manifold with corners, and if M is compact, so is $C_A(M)$.

- (2) If A is a singleton, $C_A(M) = M$. If A is a doubleton, then $C_A(M)$ is isomorphic to $M \times M$ minus a tubular neighborhood of the diagonal $\Delta \subset M \times M$. That is, $C_A(M) = M \times M - V(\Delta)$.
- (3) If $B \subset A$ then there is a natural map $C_A(M) \rightarrow C_B(M)$. In particular, for every $i, j \in A$ there is a map $\phi_{ij} : C_A(\mathbb{R}^3) \rightarrow C_{\{i,j\}}(\mathbb{R}^3) \sim S^2$.
- (4) If $f : M \rightarrow N$ is a smooth embedding, then there's a natural $f_* : C_A(M) \rightarrow C_A(N)$. □

Now let D be a graph whose set of vertices is A . If two different vertices $a_{0,1} \in A$ are connected by an edge in D , we write $a_0 \overset{D}{-} a_1$. Likewise, if $A_{0,1} \subset A$ are disjoint subsets, we write $A_0 \overset{D}{-} A_1$ if $a_0 \overset{D}{-} a_1$ for some $a_0 \in A_0$ and $a_1 \in A_1$. For any subset A_0 of A we let $D(A_0)$ be the restriction of D to A_0 .

Definition 3. The open configuration space of D in M is

$$C_D^o(M) := \{\iota : A \rightarrow M : \iota(a_0) \neq \iota(a_1) \text{ whenever } a_0 \overset{D}{-} a_1\}.$$

Definition 4. The compactified configuration space of D in M is

$$C_D(M) := \coprod_{\substack{\{A_1, \dots, A_k\} \\ A = \cup A_\alpha \\ \forall \alpha D(A_\alpha) \text{ connected}}} \left\{ \left(p_\alpha \in M, c_\alpha \in \tilde{C}_{D(A_\alpha)}(T_{p_\alpha} M) \right)_{\alpha=1}^k : p_\alpha \neq p_\beta \text{ whenever } A_\alpha \overset{D}{-} A_\beta \right\}$$

where if V is a vector space and $|A| \geq 2$,

$$\tilde{C}_D(V) := \coprod_{\substack{\{A_1, \dots, A_k\} \\ A = \cup A_\alpha; k \geq 2 \\ \forall \alpha D(A_\alpha) \text{ connected}}} \left\{ \left(v_\alpha \in V, c_\alpha \in \tilde{C}_{D(A_\alpha)}(T_{v_\alpha} V) \right)_{\alpha=1}^k : v_\alpha \neq v_\beta \text{ whenever } A_\alpha \overset{D}{-} A_\beta \right\} / \text{translations and dilations.}$$

while if A is a singleton, $\tilde{C}_D(V) := \{\text{a point}\}$.

Theorem 2. The obvious parallel of the previous theorem holds. □

Date: December 9, 2002.

This handout is available from <http://www.ma.huji.ac.il/~drorbn/classes/0102/FeynmanDiagrams/>.