

The Kauffman Bracket: $\langle \emptyset \rangle = 1$; $\langle \bigcirc L \rangle = (q + q^{-1})\langle L \rangle$; $\langle \times \rangle = \langle \text{0-smoothing} \rangle - q \langle \text{1-smoothing} \rangle$.

The Jones Polynomial: $\hat{J}(L) = (-1)^{n_+ - n_-} q^{n_+ - 2n_-} \langle L \rangle$, where (n_+, n_-) count (\times, \times) crossings.

Khovanov's construction: $\llbracket L \rrbracket$ — a chain complex of graded \mathbb{Z} -modules;

$$\llbracket \emptyset \rrbracket = 0 \rightarrow \begin{matrix} \mathbb{Z} \\ \text{height 0} \end{matrix} \rightarrow 0; \quad \llbracket \bigcirc L \rrbracket = V \otimes \llbracket L \rrbracket; \quad \llbracket \times \rrbracket = \text{Flatten} \left(0 \rightarrow \begin{matrix} \mathbb{Z} \\ \text{height 0} \end{matrix} \rightarrow \begin{matrix} \mathbb{Z} \\ \text{height 1} \end{matrix} \right);$$

$$\mathcal{H}(L) = \mathcal{H}(\mathcal{C}(L)) = \llbracket L \rrbracket[-n_-]\{n_+ - 2n_-\}$$


$$V = \text{span}\langle v_+, v_- \rangle; \quad \deg v_\pm = \pm 1; \quad q\dim V = q + q^{-1} \quad \text{with} \quad q\dim \mathcal{O} := \sum_m q^m \dim \mathcal{O}_m;$$

$$\mathcal{O}\{l\}_m := \mathcal{O}_{m-l} \quad \text{so} \quad q\dim \mathcal{O}\{l\} = q^l q\dim \mathcal{O}; \quad \cdot[s] : \quad \text{height shift by } s;$$

| | |
|---|--|
| $(\bigcirc \bigcirc \xrightarrow{\text{smoothing}} \bigcirc \times)$ $\longrightarrow (V \otimes V \xrightarrow{m} V)$ | $m : \begin{cases} v_+ \otimes v_- \mapsto v_- & v_+ \otimes v_+ \mapsto v_+ \\ v_- \otimes v_+ \mapsto v_- & v_- \otimes v_- \mapsto 0 \end{cases}$ |
| $(\times \times \xrightarrow{\text{smoothing}} \bigcirc \bigcirc)$ $\longrightarrow (V \xrightarrow{\Delta} V \otimes V)$ | $\Delta : \begin{cases} v_+ \mapsto v_+ \otimes v_- + v_- \otimes v_+ \\ v_- \mapsto v_- \otimes v_- \end{cases}$ |

Example:

$$\vdash q^{-2} + 1 + q^2 - q^6 \xrightarrow[\text{(with } (n_+, n_-) = (3, 0)\text{)}]{\cdot(-1)^{n_+ - n_-}} q + q^3 + q^5 - q^9.$$



(here $(-1)^\xi := (-1)^{\sum_i \xi_i}$ if $\xi_j = \star$)

$$= \llbracket \circledast \rrbracket \xrightarrow[\text{(with } (n_+, n_-) = (3, 0)\text{)}]{\cdot[-n_-]\{n_+ - 2n_-\}} \mathcal{C}(\circledast).$$

Theorem 1. The graded Euler characteristic of $\mathcal{C}(L)$ is $\hat{J}(L)$.

Theorem 2. The homology $\mathcal{H}(L)$ is a link invariant and thus so is $Kh_{\mathbb{F}}(L) := \sum_r t^r q\dim \mathcal{H}_{\mathbb{F}}^r(\mathcal{C}(L))$ over any field \mathbb{F} .

Theorem 3. $\mathcal{H}(\mathcal{C}(L))$ is strictly stronger than $\hat{J}(L)$: $\mathcal{H}(\mathcal{C}(\bar{5}_1)) \neq \mathcal{H}(\mathcal{C}(10_{132}))$ whereas $\hat{J}(\bar{5}_1) = \hat{J}(10_{132})$.

Conjecture 1. $Kh_{\mathbb{Q}}(L) = q^{s-1} (1 + q^2 + (1 + tq^4)Kh')$ and $Kh_{\mathbb{F}_2}(L) = q^{s-1} (1 + q^2) (1 + (1 + tq^2)Kh')$ for even $s = s(L)$ and non-negative-coefficients laurent polynomial $Kh' = Kh'(L)$.

Conjecture 2. For alternating knots s is the signature and Kh' depends only on tq^2 .

References. Khovanov's arXiv:math.QA/9908171 and arXiv:math.QA/0103190 and DBN's