

DEPARTMENT OF MATHEMATICS  
University of Toronto

**Practice Exam in Algebra (3 hours)**

1. (a) State the class equation for the finite group  $G$ , explaining briefly the meaning of the terms used.  
(b) If  $G$  is a finite group of order  $p^n$ , where  $p$  is a prime and  $n \geq 1$ , show that the center  $Z(G)$  of  $G$  is non-trivial.
2.  $G$  is a group of order 100. Prove that  $G$  is not simple. Must  $G$  be solvable? Must  $G$  be nilpotent? (Justify your answer.)
3.  $F$  is a free abelian group with basis  $x_1, x_2, x_3, x_4$  and  $H$  is the subgroup generated by the subset  $\{x_1 + x_2 - 2x_3 - x_4, 2x_1 - 2x_3, 2x_2 + 4x_3 + 4x_4\}$ . If  $A = F/H$ , find the free rank and the invariant factors of  $A$ .
4.  $R$  is a ring, and  $M$  is an  $R$ -module.
  - (a) If  $S$  is a subset of  $M$ , explain what it means to say  $M$  is a free  $R$ -module with basis  $S$ .
  - (b) If  $A$  and  $B$  are free  $R$ -modules, prove that  $A \oplus B$  is a free  $R$ -module.
  - (c) If  $M = A \oplus B$ , where  $A$  and  $B$  are  $R$ -modules, and  $M$  is free, need  $A$  and  $B$  be free? (Justify your answer.)
5.  $R$  is a commutative ring.
  - (a) Define what is meant by a prime ideal of  $R$ .
  - (b) If  $J$  is a prime ideal of  $R$ , show that the quotient ring  $R/J$  is an integral domain. Is the converse true?
  - (c) Give an example of a non-zero prime ideal of  $\mathbb{Z}[x]$  which is not a maximal ideal.
6.  $R$  is an integral domain, and  $N: R^\# \rightarrow \mathbb{Z}^+$  is a function (from the non-zero elements of  $R$  to the positive integers) such that  $N(\alpha\beta) = N(\alpha)N(\beta)$  for all  $\alpha, \beta \in R^\#$ , and  $N(\alpha) = 1$  if and only if  $\alpha$  is a unit of  $R$ .
  - (a) Define what is meant by an irreducible element of  $R$ .

- (b) If  $N(\beta) > 1$ , prove that  $\beta$  can be written as a product of  $(\geq 1)$  irreducibles.
  - (c) State an example of such an  $R$  which is not a unique factorisation domain.
7. Let  $p$  be a prime, and let  $K$  be the subfield of  $\mathbb{C}$  which is the splitting field over  $\mathbb{Q}$  of  $x^p - 3$ .
- (a) Show  $x^p - 3$  is irreducible over  $\mathbb{Q}$ .
  - (b) If  $\alpha$  is a primitive  $p$ th root of unity in  $\mathbb{C}$ , explain why  $\mathbb{Q}(\alpha)$  is a subfield of  $K$ .
  - (c) Let  $G$  be the Galois group of  $K$  over  $\mathbb{Q}$ . Show that  $G$  has order  $p(p-1)$ , and that  $G$  has a normal subgroup  $H$  of order  $p$ .
8. (a) Find the order of the abelian group

$$(\mathbb{Z}_2 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_6) \otimes (\mathbb{Z}_{10} \oplus \mathbb{Z}_{12}) .$$

- (b) Prove that  $\mathbb{Q}/\mathbb{Z} \otimes \mathbb{Q}/\mathbb{Z} = \{0\}$ .
9.  $R$  is a ring.
- (a) Give a definition of the Jacobson Radical  $J(R)$  of  $R$ , in terms of the annihilators  $A(M)$  of simple  $R$  modules  $M$ .
  - (b) For  $y \in R$ , show  $y \in J(R)$  if and only if  $1 - xy$  has a left inverse, for all  $x \in R$ .
  - (c) If  $F$  is a field, find  $J(F[x_1, \dots, x_n])$ .

N.B. Each ring  $R$  is required to have an identity element, and modules and ring homomorphisms are required to be unitary.