

DEPARTMENT OF MATHEMATICS
University of Toronto

Topology Exam (3 hours)

January 1997


No aids.

Do all questions.

Each question is worth 20 marks.

1. a) Let $f: S^2 \times I \rightarrow \mathbb{R}$ be continuous. Suppose that $\min_{p \in S^2} f(p, 0) = 0$, $\min_{p \in S^2} f(p, 1) = 1$.
Prove that for every $0 < m < 1$, $\exists 0 < t < 1$ such that $\min_{p \in S^2} f(p, t) = m$.
b) Does the result in part (a) hold if S^2 is replaced by \mathbb{R} ? Explain your answer.
2. Let $X_1 \subset X_2 \subset \cdots$ be a sequence of Hausdorff spaces where X_i is a closed subspace of X_{i+1} for each i . Let $X = \bigcup_{i=1}^{\infty} X_i$. Define the coherent topology on X by $U \subset X$ is open $\Leftrightarrow U \cap X_i$ is open in $X_i \forall i$.
 - a) Verify that this is a topology.
 - b) Show that X_i is a subspace of X in this topology.
 - c) Suppose that each X_i is normal; state Tietze's extension theorem and use it to show that X is normal.
3. Let M be a compact connected manifold. (Recall that a manifold is Hausdorff and locally Euclidean, that is, every point has a neighbourhood homeomorphic to an open set in \mathbb{R}^n for some n .) Let $\pi: P \rightarrow M$ be the universal cover. Show that P is compact $\Leftrightarrow \pi_1(M)$ is finite.

4. Let X be the outline of the tetrahedron; that is, $X = \bigcup_{i=1}^6 L_i \cup \bigcup_{i=1}^4 P_i$



where L_i are the edges and P_i are the vertices. Calculate $H_*(X; \mathbb{Z})$.

5. a) Calculate $\pi_1(S^2 \times S^1)$.
 b) Calculate $\pi_1(S^2 \times \Delta)$ where $\Delta = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$.
 c) Notice that $S^2 \times S^1 \underset{\text{homeo}}{\cong} S^2 \times \partial\Delta$. Show that $S^2 \times S^1$ is not a retract of $S^2 \times \Delta$.