

3.2 Vector bundles

Definition 25. A smooth real vector bundle of rank k over the base manifold M is a manifold E (called the total space), together with a smooth surjection $\pi : E \rightarrow M$ (called the bundle projection), such that

- $\forall p \in M, \pi^{-1}(p) = E_p$ has the structure of k -dimensional vector space,
- Each $p \in M$ has a neighbourhood U and a diffeomorphism $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$ (called a local trivialization of E over U) such that $\pi_1(\Phi(\pi^{-1}(x))) = x$, where $\pi_1 : U \times \mathbb{R}^k \rightarrow U$ is the first projection, and also that $\Phi : \pi^{-1}(x) \rightarrow \{x\} \times \mathbb{R}^k$ is a linear map, for all $x \in M$.

Given two local trivializations $\Phi_i : \pi^{-1}(U_i) \rightarrow U_i \times \mathbb{R}^k$ and $\Phi_j : \pi^{-1}(U_j) \rightarrow U_j \times \mathbb{R}^k$, we obtain a smooth gluing map $\Phi_j \circ \Phi_i^{-1} : U_{ij} \times \mathbb{R}^k \rightarrow U_{ij} \times \mathbb{R}^k$, where $U_{ij} = U_i \cap U_j$. This map preserves images to M , and hence it sends (x, v) to $(x, g_{ji}(v))$, where g_{ji} is an invertible $k \times k$ matrix smoothly depending on x . That is, the gluing map is uniquely specified by a smooth map

$$g_{ji} : U_{ij} \rightarrow GL(k, \mathbb{R}).$$

These are called transition functions of the bundle, and since they come from $\Phi_j \circ \Phi_i^{-1}$, they clearly satisfy $g_{ij} = g_{ji}^{-1}$ as well as the ‘‘cocycle condition’’

$$g_{ij}g_{jk}g_{ki} = \text{Id}|_{U_i \cap U_j \cap U_k}.$$

Example 3.11. To build a vector bundle, choose an open cover $\{U_i\}$ and form the pieces $\{U_i \times \mathbb{R}^k\}$. Then glue these together on double overlaps $\{U_{ij}\}$ via functions $g_{ij} : U_{ij} \rightarrow GL(k, \mathbb{R})$. As long as g_{ij} satisfy $g_{ij} = g_{ji}^{-1}$ as well as the cocycle condition, the resulting space has a vector bundle structure.

Example 3.12. Let $S^2 = U_0 \sqcup U_1$ for $U_i = \mathbb{R}^2$, as before. Then on $U_{01} = \mathbb{R}^2 \setminus \{0\} = \mathbb{C} \setminus \{0\}$, define

$$g_{01}(z) = [z^k], \quad k \in \mathbb{Z}.$$

In real coordinates $z = re^{i\theta}$, $g_{01}(r, \theta) = r^k \begin{pmatrix} \cos(k\theta) & -\sin(k\theta) \\ \sin(k\theta) & \cos(k\theta) \end{pmatrix}$. This defines a vector bundle $E_k \rightarrow S^2$ of rank 2 for each $k \in \mathbb{Z}$ (or a complex vector bundle of rank 1, since $g_{01} : U_{01} \rightarrow GL(1, \mathbb{C})$). Actually, since the map g_{01} is actually holomorphic as a function of z , we have defined holomorphic vector bundles on $\mathbb{C}P^1$.

Example 3.13 (The tangent bundle). The tangent bundle TM is indeed a vector bundle, of rank $\dim M$. For any chart (U, φ) of M , there is an associated local trivialization $(\pi^{-1}(U), \Phi)$ of TM , and the transition function $g_{ji} : U_{ij} \rightarrow GL(n, \mathbb{R})$ between two trivializations obtained from $(U_i, \varphi_i), (U_j, \varphi_j)$ is simply the Jacobian matrix

$$g_{ji} : p \mapsto D(\varphi_j \circ \varphi_i^{-1})(p).$$

Just as for the tangent bundle, we can define the analog of a vector-valued function, where the function has values in a vector bundle:

Definition 26. A smooth section of the vector bundle $E \xrightarrow{\pi} M$ is a smooth map $s : M \rightarrow E$ such that $\pi \circ s = \text{Id}_M$. The set of all smooth sections, denoted $\Gamma^\infty(M, E)$, is an infinite-dimensional real vector space, and is also a module over the ring $C^\infty(M, \mathbb{R})$.

Having introduced vector bundles, we must define the notion of *morphism* between vector bundles, so as to form a category.

Definition 27. A smooth bundle map between the bundles $E \xrightarrow{\pi} M$ and $E' \xrightarrow{\pi'} M'$ is a pair (f, F) of smooth maps $f : M \rightarrow M'$ and $F : E \rightarrow E'$ such that $\pi' \circ F = f \circ \pi$ and such that $F : E_p \rightarrow E'_{f(p)}$ is a linear map for all p .

Example 3.14. I claim that the bundles $E_k \xrightarrow{\pi} S^2$ are all non-isomorphic, except that E_k is isomorphic to E_{-k} over the antipodal map $S^2 \rightarrow S^2$.

Example 3.15. Suppose $f : M \rightarrow N$ is a smooth map. Then $f_* : TM \rightarrow TN$ is a bundle map covering f , i.e. (f_*, f) defines a bundle map.

Example 3.16 (Pullback bundle). if $f : M \rightarrow N$ is a smooth map and $E \xrightarrow{\pi} N$ is a vector bundle over N , then we may form the fiber product $M_f \times_{\pi} E$, which then is a bundle over M with local trivializations $(f^{-1}(U_i), f^*g_{ij})$, where (U_i, g_{ij}) is the local transition data for E over N . This bundle is called the pullback bundle and is denoted by f^*E . The natural projection to E defines a vector bundle map back to E :

$$\begin{array}{ccc} f^*E & \xrightarrow{p_2} & E \\ p_1 \downarrow & & \downarrow \pi \\ M & \xrightarrow{f} & N \end{array}$$

There is also a natural pullback map on sections: given a section $s \in \Gamma^\infty(N, E)$, the composition $s \circ f$ gives a map $M \rightarrow E$. This then determines a smooth map $f^*s : M \rightarrow f^*E$ by the universal property of the fiber product. We therefore obtain a pullback map

$$f^* : \Gamma^\infty(N, E) \rightarrow \Gamma^\infty(M, f^*E).$$

Example 3.17. If $f : M \rightarrow N$ is an embedding, then so is the bundle map $f_* : TM \rightarrow TN$. By the universal property of the fiber product we obtain a bundle map, also denoted f_* , from TM to f^*TN . This is a vector bundle inclusion and $f^*TN/f_*TM = NM$ is a vector bundle over M called the normal bundle of M . Note: we haven't covered subbundles and quotient bundles in detail. I'll leave this as an exercise.

3.3 Associated bundles

We now describe a functorial construction of vector bundles, using functors from vector spaces. Consider the category $\mathbf{Vect}_{\mathbb{R}}$ of finite-dimensional real vector spaces and linear maps. We will describe several functors from $\mathbf{Vect}_{\mathbb{R}}$ to itself.

Example 3.18. If $V \in \mathbf{Vect}_{\mathbb{R}}$, then $V^* \in \mathbf{Vect}_{\mathbb{R}}$, and if $f : V \rightarrow W$ then $f^* : W^* \rightarrow V^*$. Since the composition of duals is the dual of the composition, duality defines a contravariant functor $*$: $\mathbf{Vect}_{\mathbb{R}} \rightarrow \mathbf{Vect}_{\mathbb{R}}$.

Example 3.19. If $V, W \in \mathbf{Vect}_{\mathbb{R}}$, then $V \oplus W \in \mathbf{Vect}_{\mathbb{R}}$, and this defines a covariant functor $\mathbf{Vect}_{\mathbb{R}} \times \mathbf{Vect}_{\mathbb{R}} \rightarrow \mathbf{Vect}_{\mathbb{R}}$.

Example 3.20. If $V, W \in \mathbf{Vect}_{\mathbb{R}}$, then $V \otimes W \in \mathbf{Vect}_{\mathbb{R}}$ and this again defines a covariant functor $\mathbf{Vect}_{\mathbb{R}} \times \mathbf{Vect}_{\mathbb{R}} \rightarrow \mathbf{Vect}_{\mathbb{R}}$.

Example 3.21. If $V \in \mathbf{Vect}_{\mathbb{R}}$, then

$$\otimes^\bullet V = \mathbb{R} \oplus V \oplus (V \otimes V) \oplus \dots \oplus (\otimes^k V) \oplus \dots$$

is an infinite-dimensional vector space, with a product $a \otimes b$. Quotienting by the double-sided ideal $I = \langle v \otimes v : v \in V \rangle$, we obtain the exterior algebra

$$\wedge^\bullet V = \mathbb{R} \oplus V \oplus \wedge^2 V \oplus \dots \oplus \wedge^n V,$$

with $n = \dim V$. The product is customarily denoted $(a, b) \mapsto a \wedge b$. The direct sum decompositions above, where $\wedge^k V$ or $\otimes^k V$ is labeled by the integer k , are called \mathbb{Z} -gradings, and since the product takes $\wedge^k \times \wedge^l \rightarrow \wedge^{k+l}$, these algebras are called \mathbb{Z} -graded algebras.

If (v_1, \dots, v_n) is a basis for V , then $v_{i_1} \wedge \dots \wedge v_{i_k}$ for $i_1 < \dots < i_k$ form a basis for $\wedge^k V$. This space then has dimension $\binom{n}{k}$, hence the algebra $\wedge^\bullet V$ has dimension 2^n .

Note in particular that $\wedge^n V$ has dimension 1, is also called the determinant line $\det V$, and a choice of nonzero element in $\det V$ is called an “orientation” on the vector space V .

Recall that if $f : V \rightarrow W$ is a linear map, then $\wedge^k f : \wedge^k V \rightarrow \wedge^k W$ is defined on monomials via

$$\wedge^k f(a_1 \wedge \dots \wedge a_k) = f(a_1) \wedge \dots \wedge f(a_k).$$

In particular, if $A : V \rightarrow V$ is a linear map, then for $n = \dim V$, the top exterior power $\wedge^n A : \wedge^n V \rightarrow \wedge^n V$ is a linear map of a 1-dimensional space onto itself, and is hence given by a number, called $\det A$, the determinant of A .

We may now apply any of these functors to vector bundles. The main observation is that if F is a vector space functor as above, we may apply it to any vector bundle $E \xrightarrow{\pi} M$ to obtain a new vector bundle

$$F(E) = \sqcup_{p \in M} F(E_p).$$

If (U_i) is an atlas for M and E has local trivializations $(U_i \times \mathbb{R}^k)$, glued together via $g_{ji} : U_{ij} \rightarrow GL(k, \mathbb{R})$, then $F(E)$ may be given the local trivialization $(U_i \times F(\mathbb{R}^k))$, glued together via $F(g_{ji})$. This new vector bundle $F(E)$ is called the “associated” vector bundle to E , given by the functor F .

Example 3.22. If $E \rightarrow M$ is a vector bundle, then $E^* \rightarrow M$ is the dual vector bundle. If E, F are vector bundles then $E \oplus F$ is called the direct or “Whitney” sum, and has rank $\text{rk } E + \text{rk } F$. $E \otimes F$ is the tensor product bundle, which has rank $\text{rk } E \cdot \text{rk } F$.

Example 3.23. If $E \rightarrow M$ is a vector bundle of rank n , then $\otimes^k E$ and $\wedge^k E$ are its tensor power bundles, of rank n^k and $\binom{n}{k}$, respectively. The top exterior power $\wedge^n E$ has rank 1, and is hence a line bundle. If this line bundle is trivial (i.e. isomorphic to $M \times \mathbb{R}$) then E is said to be an orientable bundle.

Example 3.24. Starting with the tangent bundle $TM \rightarrow M$, we may form the cotangent bundle T^*M , the bundle of tensors of type (r, s) , $\otimes^r TM \otimes \otimes^s T^*M$.

We may also form the bundle of multivectors $\wedge^k TM$, which has sections $\Gamma^\infty(M, \wedge^k TM)$ called multivector fields.

Finally, we may form the bundle of k -forms, $\wedge^k T^*M$, whose sections $\Gamma^\infty(M, \wedge^k T^*M) = \Omega^k(M)$ are called differential k -forms, and will occupy us for some time.

We have now produced several vector bundles by applying functors to the tangent bundle. We are familiar with vector fields, which are sections of TM , and we know that a vector field is written locally in coordinates (x^1, \dots, x^n) as

$$X = \sum_i a^i \frac{\partial}{\partial x^i},$$

with coefficients a^i smooth functions.

There is an easy way to produce examples of 1-forms in $\Omega^1(M)$, using smooth functions f . We note that the action $X \mapsto X(f)$ defines a dual vector at each point of M , since $(X(f))_p$ depends only on the vector X_p and not the behaviour of X away from p . Recall that $X(f) = Df_2(X)$.

Definition 28. The exterior derivative of a function f , denoted df , is the section of T^*M given by the fiber projection Df_2 .

Since $dx^i(\frac{\partial}{\partial x^j}) = \delta_j^i$, we see that (dx^1, \dots, dx^n) is the dual basis to $(\frac{\partial}{\partial x^1}, \dots, \frac{\partial}{\partial x^n})$. Therefore, a section of T^*M has local expression

$$\xi = \sum_i \xi_i dx^i,$$

for ξ_i smooth functions, given by $\xi_i = \xi(\frac{\partial}{\partial x^i})$. In particular, the exterior derivative of a function df can be written

$$df = \sum_i \frac{\partial f}{\partial x^i} dx^i.$$

A section of the tensor bundle $\otimes^r TM \otimes \otimes^s T^*M$ can be written as

$$\Theta = \sum_{\substack{i_1, \dots, i_r \\ j_1, \dots, j_s}} a_{j_1 \dots j_s}^{i_1 \dots i_r} \frac{\partial}{\partial x^{i_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{i_r}} \otimes dx^{j_1} \otimes \dots \otimes dx^{j_s},$$

where $a_{j_1 \dots j_s}^{i_1 \dots i_r}$ are n^{r+s} smooth functions.

A general differential form $\rho \in \Omega^k(M)$ can be written

$$\rho = \sum_{i_1 < \dots < i_k} \rho_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

3.4 Differential forms

There are several properties of differential forms which make them indispensable: first, the k -forms are intended to give a notion of k -dimensional volume (this is why they are multilinear and skew-symmetric, like the determinant) and in a way compatible with the boundary map (this leads to the exterior derivative, which we define below). Second, they behave well functorially, as we see now.

Given a smooth map $f : M \rightarrow N$, we obtain bundle maps $f_* : TM \rightarrow TN$ and hence $f^* := \wedge^k(f_*)^* : \wedge^k T^*N \rightarrow \wedge^k T^*M$. Hence we have the diagram

$$\begin{array}{ccc} \wedge^k T^*M & \xleftarrow{f^*} & \wedge^k T^*N \\ \pi_M \downarrow & & \downarrow \pi_N \\ M & \xrightarrow{f} & N \end{array}$$

The interesting thing is that if $\rho \in \Omega^k(N)$ is a differential form on N , then it is a section of π_N . Composing with f, f^* , we obtain a section $f^*\rho := f^* \circ \rho \circ f$ of π_M . Hence we obtain a natural map

$$\Omega^k(N) \xrightarrow{f^*} \Omega^k(M).$$

Such a natural map does not exist (in either direction) for multivector fields, for instance.

Suppose that $\rho \in \Omega^k(N)$ is given in a coordinate chart by $\rho = \sum \rho_{i_1 \dots i_k} dy^{i_1} \wedge \dots \wedge dy^{i_k}$. Now choose a coordinate chart for M with coordinates x^1, \dots, x^m . What is the local expression for $f^*\rho$? We need only compute f^*dy_i . We use a notation where f^k denotes the k^{th} component of f in the coordinates (y^1, \dots, y^n) , i.e. $f^k = y^k \circ f$.

$$f^*dy_i(\frac{\partial}{\partial x^j}) = dy_i(f_* \frac{\partial}{\partial x^j}) \quad (23)$$

$$= dy_i(\sum_k \frac{\partial f^k}{\partial x^j} \frac{\partial}{\partial y_k}) \quad (24)$$

$$= \frac{\partial f^i}{\partial x^j}. \quad (25)$$

Hence we conclude that

$$f^*dy_i = \sum_j \frac{\partial f^i}{\partial x^j} dx^j.$$

Finally we compute

$$f^*\rho = \sum_{i_1 < \dots < i_k} f^* \rho_{i_1 \dots i_k} f^*(dy^{i_1}) \wedge \dots \wedge f^*(dy^{i_k}) \quad (26)$$

$$= \sum_{i_1 < \dots < i_k} (\rho_{i_1 \dots i_k} \circ f) \sum_{j_1} \dots \sum_{j_k} \frac{\partial f^{i_1}}{\partial x^{j_1}} \dots \frac{\partial f^{i_k}}{\partial x^{j_k}} dx^{j_1} \wedge \dots \wedge dx^{j_k}. \quad (27)$$