Reading: Woit Chapter 22

Exercise 1. Recall that the 1-dimensional harmonic oscillator is the quantum system where $\mathcal{H} = L^2(\mathbb{R})$ and $H = \frac{1}{2}(P^2 + Q^2)$. We saw that H has a 1-dimensional space of ground states (lowest-eigenvalue states) generated by the eigenvector

$$|0\rangle = \pi^{-1/4} e^{-q^2/2}$$

with eigenvalue 1/2, and that by applying the raising operator $a^* = 2^{-1/2}(Q - iP)$ we produce a sequence of states

$$|n+1\rangle = (n+1)^{-1/2}a^* |n\rangle, n = 0, 1, 2, \dots$$

such that $|n\rangle$ has eigenvalue $E_n = n + 1/2$.

Prove that any eigenstate of H must lie in one of the eigenspaces listed above.

Exercise 2. Consider the 3-d harmonic oscillator, with $\mathcal{H} = L^2(\mathbb{R}^3)$ and Hamiltonian

$$H = \frac{1}{2}(P_1^2 + P_2^2 + P_3^2) + \frac{1}{2}(Q_1^2 + Q_2^2 + Q_3^2).$$

- 1. What are the eigenvalues of H?
- 2. Describe an explicit basis of eigenvectors for the eigenspaces corresponding to the three lowest eigenvalues.
- 3. The rotation group SO(3) acts on $L^2(\mathbb{R}^3)$ and the Lie algebra generators l_1, l_2, l_3 are sent by this representation to

$$\pi'(l_1) = -(q_2\frac{\partial}{\partial q_3} - q_3\frac{\partial}{\partial q_2}) \qquad \pi'(l_2) = -(q_3\frac{\partial}{\partial q_1} - q_1\frac{\partial}{\partial q_3}) \qquad \pi'(l_3) = -(q_1\frac{\partial}{\partial q_2} - q_2\frac{\partial}{\partial q_1})$$

As a result, we may express the corresponding self-adjoint operators, known as the angular momentum operators, $L_i = i\pi'(l_i)$ in terms of the linear momenta and position operators, that is,

$$L_1 = Q_2 P_3 - Q_3 P_2$$
 $L_2 = Q_3 P_1 - Q_1 P_3$ $L_3 = Q_1 P_2 - Q_2 P_1$

Determine whether the angular momentum obserables are conserved in this system.

4. What does the previous result imply about the action of SO(3) on the states found in question 2. Which irreducible representations occur?

Exercise 3. The perturbative series $\hat{F} \in \mathbb{C}[[\epsilon]]$ which describes the function

$$F(\epsilon) = \frac{\int \exp(-\frac{1}{2}ax^2 + \epsilon x^3/3!)}{\int \exp(-\frac{1}{2}ax^2)}$$

may be described as follows:

$$\hat{F}(\epsilon) = \sum_{\Gamma} \frac{1}{|\operatorname{Aut}(\Gamma)|} a^{-e} \epsilon^{v},$$

where the sum is over all possible trivalent graphs Γ with e edges and v vertices, and Aut(Γ) is the symmetry group of the graph. Also, the empty graph is assigned 1.

To be precise: divide each edge of the graph into two half-edges. Then the graph may be viewed (in fact is defined) as a collection of half-edges together with two partitions: the partition into sets of size 2 called edges, and the partition into sets of size 3 called vertices. A symmetry of the graph is a bijection from the half-edges to themselves which preserves the two partitions.

Note that there are no trivalent graphs with only 1 vertex. So the power series must begin with $1 + c\epsilon^2 + \cdots$ for some constant c. We saw in class that there are two trivalent graphs with two vertices: the theta-graph (shaped like the letter θ) and the dumbbell graph (shaped like O—O). These graphs have symmetry group of size 12 (permute 3 edges or 2 vertices) and 8 (flip each of the 3 edges independently), respectively. So we see that up to the ϵ^2 term we have

$$\hat{F}(\epsilon) = 1 + \frac{1}{12}a^{-3}\epsilon^2 + \frac{1}{8}a^{-3}\epsilon^2 + \cdots$$

Compute $\hat{F}(\epsilon)$ to the next order in perturbation theory – the coefficient of ϵ^3 is zero, so determine the coefficient of ϵ^4 . In particular, you must find all trivalent graphs with exactly four vertices and study their automorphisms.

Mega Bonus: make a numerical comparison between $F(\epsilon)$ and its asymptotic expansion (for a = 1, say): you can define $F(\epsilon)$ (for ϵ chosen generically, not necessarily on the real axis) by integrating along the path of steepest descent of $\operatorname{Re}(-x^2/2 + \epsilon x^3/3!)$, or in other words along the path given by $\operatorname{Im}(-x^2/2 + \epsilon x^3/3!) = 0$.