


**So What?** If  $V$  is a representation, then  $V^{\otimes n}$  explodes as a function of  $n$ , while in **DoPeGDO** up to a fixed power of  $\epsilon$ , the ranks of  $\text{mor}(A \rightarrow B)$  grow polynomially as a function of  $|A|$  and  $|B|$ .


**Compositions.** In  $\text{mor}(A \rightarrow B)$ ,

$$Q = \sum_{i \in A, j \in B} E_{ij} \zeta_i \zeta_j + \frac{1}{2} \sum_{i, j \in A} F_{ij} \zeta_i \zeta_j + \frac{1}{2} \sum_{i, j \in B} G_{ij} \zeta_i \zeta_j,$$

and so (remember,  $e^x = 1 + x + xx/2 + xxx/6 + \dots$ )



where  $\bullet E = E_1(I - F_2G_1)^{-1}E_2$ .  
 $\bullet F = F_1 + E_1F_2(I - G_1F_2)^{-1}E_1^T$ .  
 $\bullet G = G_2 + E_2^T G_1(I - F_2G_1)^{-1}E_2$ .  
 $\bullet \omega = \omega_1\omega_2 \det(I - F_2G_1)^{-1}$ .  
 $\bullet P$  is computed as the solution of a messy PDE or using “connected Feynman diagrams” (yet we’re still in pure algebra!). Docility is preserved.




**DoPeGDO Footnotes.** Each variable has a “weight”  $\in \{0, 1, 2\}$ , and always  $\text{wt } z_i + \text{wt } \zeta_i = 2$ .

- †1. Really, “weight-graded finite sets”  $A = A_0 \sqcup A_1 \sqcup A_2$ .
- †2. Really, a power series in the weight-0 variables<sup>†5</sup>.
- †3. The weight of  $Q$  must be 2, so it decomposes as  $Q = Q_{20} + Q_{11}$ . The coefficients of  $Q_{20}$  are rational numbers while the coefficients of  $Q_{11}$  may be weight-0 power series<sup>†5</sup>.
- †4. Setting  $\text{wt } \epsilon = -2$ , the weight of  $P$  is  $\leq 2$  (so the powers of the weight-0 variables are not constrained)<sup>†5</sup>.
- †5. In the knot-theoretic case, all weight-0 power series are rational functions of bounded degree in the exponentials of the weight-0 variables.
- †6. There’s also an obvious product  $\text{mor}(A_1 \rightarrow B_1) \times \text{mor}(A_2 \rightarrow B_2) \rightarrow \text{mor}(A_1 \sqcup A_2 \rightarrow B_1 \sqcup B_2)$ .

**Full DoPeGDO.** Compute compositions in two phases:

- A 1-1 phase over the ring of power series in the weight-0 variables, in which the weight-2 variables are spectators.
- A (slightly modified) 2-0 phase over  $\mathbb{Q}$ , in which the weight-1 variables are spectators.



knot diag	$n_k^+$ $(\rho_1^+)^+$	Alexander's $\omega^+$	genus / ribbon unknotting # / amphi?	knot diag	$n_k^+$ $(\rho_1^+)^+$	Alexander's $\omega^+$	genus / ribbon unknotting # / amphi?	knot diag	$n_k^+$ $(\rho_1^+)^+$	Alexander's $\omega^+$	genus / ribbon unknotting # / amphi?
	$0_1^+$ 0	1	0 / ✓ 0 / ✓		$3_1^+$ $T$	$T-1$	1 / ✗ 1 / ✗		$4_1^+$ 0	$3-T$	1 / ✗ 1 / ✓
	$5_1^+$ $2T^3+3T$	$T^2-T+1$	2 / ✗ 2 / ✗		$5_2^+$ $5T-4$	$2T-3$	1 / ✗ 1 / ✗		$6_1^+$ $T-4$	$5-2T$	1 / ✓ 1 / ✗
	$6_2^+$ $T^3-4T^2+4T-4$	$-T^2+3T-3$	2 / ✗ 1 / ✗		$6_3^+$ 0	$T^2-3T+5$	2 / ✗ 1 / ✓		$7_1^+$ $3T^5+5T^3+6T$	$T^3-T^2+T-1$	3 / ✗ 3 / ✗

**Questions.** • Are there QFT precedents for “two-step Gaussian integration”?


• In QFT, one saves even more by considering “one-particle-irreducible” diagrams and “effective actions”. Does this mean anything here?

• Understanding  $\text{Hom}(\mathbb{Q}[z_A] \rightarrow \mathbb{Q}[z_B])$  seems like a good cause. Can you find other applications for the technology here?

$$QU = \mathcal{U}_h(sl_{2+}^\epsilon) = A(y, b, a, x) \llbracket \hbar \rrbracket \text{ with } [a, x] = x, [b, y] = -\epsilon y, [a, b] = 0, [a, y] = -y, [b, x] = \epsilon x, \text{ and } xy - qyx = (1-AB)/\hbar, \text{ where } q = e^{\hbar\epsilon}, A = e^{-\hbar\epsilon a}, \text{ and } B = e^{-\hbar\epsilon b}.$$

Also  $\Delta(y, b, a, x) = (y_1 + B_1 y_2, b_1 + b_2, a_1 + a_2, x_1 + A_1 x_2)$ ,  $S(y, b, a, x) = (-B^{-1}y, -b, -a, -A^{-1}x)$ , and  $R = \sum \hbar^{j+k} y^j b^k \otimes a^j x^k / j! k! q^j$ .



**Theorem.** Everything of value regrading  $U = CU$  and/or its quantization  $U = QU$  is **DoPeGDO**:



also Cartan’s  $\theta$ , the Dequantizer, and more, and all of their compositions.

**Solvable Approximation.** In  $sl_n$ , half is enough! Indeed  $sl_n \oplus a_{n-1} = \mathcal{D}(\nabla, b, \delta)$ . Now define  $sl_{n+}^\epsilon := \mathcal{D}(\nabla, b, \epsilon\delta)$ . Schematically, this is  $[\nabla, \nabla] = \nabla$ ,  $[\Delta, \Delta] = \epsilon\Delta$ , and  $[\nabla, \Delta] = \Delta + \epsilon\nabla$ . The same process works for all semi-simple Lie algebras, and at  $\epsilon^{k+1} = 0$  always yields a solvable Lie algebra.

**4D Metrized Lie Algebras**



**Conclusion.** There are lots of poly-time-computable well-behaved near-Alexander knot invariants: • They extend to tangles with appropriate multiplicative behaviour. • They have cabling and strand reversal formulas.

The invariant for  $sl_{2+}^\epsilon / (\epsilon^2 = 0)$  (prior art:  $\omega\epsilon\beta/Ov$ ) attains 2,883 distinct values on the 2,978 prime knots with  $\leq 12$  crossings. HOMFLY-PT and Khovanov homology together attain only 2,786 distinct values.