

Day 3: Chern-Simons, Gaussian Integration, Feynman Diagrams

Cosmic Coincidences

Recall. $\mathcal{K} = \{\text{knots}\}$, $\mathcal{A} := \text{gr}\mathcal{A} = \mathcal{D}/\text{rels} =$

$$\begin{array}{c} \text{AS: } \text{Y} + \text{Y} = \text{O} \\ \text{STU: } \text{Y} = \text{V} - \text{X} \\ \text{IHX: } \text{I} = \text{H} - \text{X} \end{array}$$



Seek $Z: \mathcal{K} \rightarrow \hat{\mathcal{A}}$ such that if K is n -singular, $Z(K) = D_k + \dots$

$$\mathcal{K} \xrightarrow[\substack{\text{Z: high algebra} \\ \text{solving finitely many equations in finitely many unknowns}}]{} \mathcal{A} := \text{gr}\mathcal{K} \xrightarrow[\substack{\text{given a "Lie" algebra } \mathfrak{g} \\ \text{low algebra: pictures represent formulas}}]{} \text{"U}(\mathfrak{g})"$$

$\langle D, K \rangle_{\overline{\square}} := (\text{The signed Stonehenge}):$

$$D = \text{circle} \quad K = \text{dotted square} \quad \Rightarrow \quad \text{signed Stonehenge}$$

The Gaussian linking number $lk(\text{circle}) = \sum_{\text{vertical chopsticks}} (\text{signs})$



$$D = \text{circle} \Rightarrow C_D(\mathbb{R}^3, \gamma) = \text{circle} \left(\mathbb{R}^3, \gamma \right)$$

Theorem. Given a parametrized knot γ in \mathbb{R}^3 , up to renormalizing the “framing anomaly”,

$$Z(\gamma) = \sum_{D \in \mathcal{D}} \frac{C(D)D}{|\text{Aut}(D)|} \int_{C_D(\mathbb{R}^3, \gamma)} \bigwedge_{e \in E(D)} \phi_e^* \omega \in \mathcal{A}$$

is an expansion. Here \mathcal{D} is the set of all “Feynman diagrams”, $E(D)$ is the set of internal edges (and chords) of D , $C_D(\mathbb{R}^3, \gamma)$ is the configuration space of placements of D on/around γ , $\phi: C_D(\mathbb{R}^3, \gamma) \rightarrow (S^2)^{E(D)}$ is the “direction of the edges” map, and ω is a volume form on S^2 .

Gaussian Integration. (λ_{ij}) is a symmetric positive definite matrix and (λ^{ij}) is its inverse, and (λ_{ijk}) are the coefficients of some cubic form. Denote by $(x^i)_{i=1}^n$ the coordinates of \mathbb{R}^n , let $(t_i)_{i=1}^n$ be a set of “dual” variables, and let ∂^i denote $\frac{\partial}{\partial t_i}$. Also let $C := \frac{(2\pi)^{n/2}}{\det(\lambda_{ij})}$. Then

$$\int_{\mathbb{R}^n} e^{-\frac{1}{2}\lambda_{ij}x^i x^j + \frac{1}{6}\lambda_{ijk}x^i x^j x^k} = \sum_{m \geq 0} \frac{\epsilon^m}{6^m m!} \int_{\mathbb{R}^n} (\lambda_{ijk}x^i x^j x^k)^m e^{-\frac{1}{2}\lambda_{ij}x^i x^j} \\ = \sum_{m \geq 0} \frac{C\epsilon^m}{6^m m!} (\lambda_{ijk}\partial^i \partial^j \partial^k)^m e^{\frac{1}{2}\lambda^{\alpha\beta} t_\alpha t_\beta} \Big|_{t_\alpha=0} = \sum_{\substack{m, l \geq 0 \\ 3m=2l}} \frac{C\epsilon^m}{6^m m! 2^l l!} (\lambda_{ijk}\partial^i \partial^j \partial^k)^m (\lambda^{\alpha\beta} t_\alpha t_\beta)^l$$



$$= \sum_{\substack{m, l \geq 0 \\ 3m=2l}} \frac{C\epsilon^m}{6^m m! 2^l l!} \left[\begin{array}{cccc} \lambda^{\alpha_1 \beta_1} & \lambda^{\alpha_2 \beta_2} & \lambda^{\alpha_3 \beta_3} & \dots \\ \Delta t_{\alpha_1} t_{\beta_1} \Delta & \Delta t_{\alpha_2} t_{\beta_2} \Delta & \Delta t_{\alpha_3} t_{\beta_3} \Delta & \dots \\ \uparrow \partial^{i_1} & \uparrow \partial^{i_2} & \uparrow \partial^{i_3} & \dots \\ \dots \text{sum over all pairings} \dots & & & \\ \downarrow \partial^{i_m} & \downarrow \partial^{j_1} & \downarrow \partial^{j_2} & \dots \\ \lambda_{i_1 j_1 k_1} & \lambda_{i_2 j_2 k_2} & \lambda_{i_3 j_3 k_3} & \dots \\ & & & \end{array} \right]$$

$$= \sum_{\substack{m, l \geq 0 \\ 3m=2l}} \frac{C\epsilon^m}{6^m m! 2^l l!} \sum_{\substack{\text{m-vertex fully marked} \\ \text{Feynman diagrams } D}} \mathcal{E}(D) \\ = C \sum_{\substack{\text{unmarked Feynman} \\ \text{diagrams } D}} \frac{\epsilon^m |\mathcal{E}(D)|}{|\text{Aut}(D)|}.$$

Claim. The number of pairings that produce a given unmarked Feynman diagram D is $\frac{6^m m! 2^l l!}{|\text{Aut}(D)|}$.

Proof of the Claim. The group $G_{m,l} := [(S_3)^m \rtimes S_m] \times [(S_2)^l \rtimes S_l]$ acts on the set of pairings, the action is transitive on the set of pairings P that produce a given D , and the stabilizer of any given P is $\text{Aut}(D)$. \square


The generating function of all cosmic coincidences:

$$Z(K) := \lim_{N \rightarrow \infty} \sum_{\substack{\text{3-valent } D \\ \text{}}} \frac{\langle D, K \rangle_{\overline{\square}} D}{2^c c! \binom{N}{e}} \in \mathcal{A}$$



Claim. It all comes from the Chern-Simons-Witten theory,

$$\int_{A \in \Omega^1(\mathbb{R}^3, \mathfrak{g})} \mathcal{D}A \text{tr}_R \text{hol}_\gamma(A) \exp \left[\frac{ik}{4\pi} \int_{\mathbb{R}^3} \text{tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right],$$



where $\Omega^1(\mathbb{R}^3, \mathfrak{g})$ is the space of all \mathfrak{g} -valued 1-forms on \mathbb{R}^3 (really, connections), k is some large constant, R is some representation of \mathfrak{g} and tr_R is trace in R , and $\text{hol}_\gamma(A)$ is the holonomy of A along γ .

References. Witten’s *Quantum field theory and the Jones polynomial*, Axelrod-Singer’s *Chern-Simons perturbation theory I-II*, D. Thurston’s arXiv:math.QA/9901110, Polyak’s arXiv:math.GT/0406251, and my videotaped 2014 class ω/AKT .

The Fourier Transform.

$$(F: V \rightarrow \mathbb{C}) \Rightarrow (\tilde{f}: V^* \rightarrow \mathbb{C})$$

via $\tilde{F}(\varphi) := \int_V f(v) e^{-i\langle \varphi, v \rangle} dv$. Some facts:

- $\tilde{f}(0) = \int_V f(v) dv$.
- $\frac{\partial}{\partial \varphi_i} \tilde{f} \sim \widetilde{v^i f}$.
- $\widetilde{(e^{Q/2})} \sim e^{Q^{-1}/2}$, where Q is quadratic, $Q(v) = \langle Lv, v \rangle$ for $L: V \rightarrow V^*$, and $Q^{-1}(\varphi) := \langle \varphi, L^{-1}\varphi \rangle$. (This is the key point in the proof of the Fourier inversion formula!)

Examples.

$$\begin{array}{ccc} \text{Feynman diagram} & | \text{Aut}(D)| = 12 & | \text{Aut}(D)| = 8 \end{array}$$

Monsters left to Slay.

- Convergence.
- Proof of invariance.
- The framing anomaly.
- Universality.
- d^{-1} doesn’t really exist, Faddeev-Popov, determinants, ghosts, Berezin integration.
- Assembly.

