MAT 347 Semidirect products October 30, 2018

Semidirect products

Recall that if if A, B are subsets of a group G, we write $AB = \{ab : a \in A, b \in B\} \subset G$. The first two problems are a review from Homework 5!

- 1. Suppose that H, K are subgroups of G. Suppose that $H \cap K = \{1\}$. Prove that every element of HK can be written uniquely as hk for $h \in H, k \in K$.
- 2. Suppose that H, K are normal subgroups of G and $H \cap K = \{1\}$. Explain how to multiply h_1k_1 with h_2k_2 . Prove that HK is isomorphic to $H \times K$.
- 3. Prove that $D_{4n} \cong D_{2n} \times \mathbb{Z}/2\mathbb{Z}$ if n is odd.
- 4. Suppose that N is normal in G, but K is not. (We will write N instead of H, to remember that N is normal.) Explain how to multiply n_1k_1 with n_2k_2 , expressing you answer as nk for some $n \in N, k \in K$.
- 5. Suppose now that N, K are two abstract groups (i.e. not embedded as subgroups of a third group). Suppose that we are given a homomorphism φ : K → Aut N, so for each element k ∈ K, we are given an automorphism φ(k) : N → N of N. Explain how we can use this to define a new group structure on the set N × K, motivated by your computation in Question 4. The set N × K with this group structure will be denoted N ⋈_φ K and is called the semidirect product of N and K with respect to φ.
- 6. Show that N, K are both subgroups of $N \rtimes_{\varphi} K$ and that N is a normal subgroup.
- 7. Show that D_{2n} is isomorphic to a semidirect product of $\mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z}$.
- 8. Let F be a field. Consider $N = F, K = F^{\times}$. Define a natural map $K \to \text{Aut } N$ (not the trivial one with kernel K) and form the semidirect product $N \rtimes K$. How can you think about this group?

Remark: we write $N \bowtie_{\varphi} K$ (as opposed to $K \bowtie_{\varphi} N$) to remember that $N \triangleleft G$.

Isometries

Definition: An isometry of the plane is a map $f : \mathbb{R}^2 \to \mathbb{R}^2$ such that |f(x) - f(y)| = |x - y| (where $|\cdot|$ denotes the length of a vector). The set of isometries of the plane forms a group $\text{Isom}(\mathbb{R}^2)$ under composition.

- 1. Show that any translation is an isometry.
- 2. Show that any orthogonal linear operator on \mathbb{R}^2 is an isometry.
- 3. Show that any isometry is the composition of a translation and an orthogonal linear map. (You may use the following fact without proof: if f is an isometry such that f(0) = 0, then f is an orthogonal linear map.)
- 4. What can you say about the subgroup of translations inside Isom(\mathbb{R}^2)?
- 5. Express $\text{Isom}(\mathbb{R}^2)$ as a semidirect product.