

# Linear Algebraic Groups

Florian Herzig

Notes by Joshua Seaton

Spring 2013

# Contents

<b>Introduction</b>	<b>3</b>
<b>0. Algebraic geometry (review).</b>	<b>4</b>
0.1 Zariski topology on $k^n$ . . . . .	4
0.2 Nullstellensatz. . . . .	4
0.3 Some topology. . . . .	4
0.4 Functions on closed subsets of $k^n$ . . . . .	5
0.5 Products. . . . .	6
0.6 Subvarieties. . . . .	6
0.7 Projective varieties. . . . .	7
0.8 Dimension. . . . .	7
0.9 Constructible sets. . . . .	8
0.10 Other examples. . . . .	8
<b>1. Algebraic groups: beginnings.</b>	<b>9</b>
1.1 Preliminaries. . . . .	9
1.2 Subgroups. . . . .	10
1.3 Commutators. . . . .	12
1.4 $G$ -spaces. . . . .	13
1.5 Jordan Decomposition. . . . .	15
<b>2. Diagonalisable and elementary unipotent groups.</b>	<b>19</b>
2.1 Unipotent and semisimple subsets. . . . .	19
2.2 Diagonalisable groups and tori. . . . .	20
2.3 Elementary unipotent groups. . . . .	25
<b>3. Lie algebras.</b>	<b>27</b>
3.1 Tangent Spaces. . . . .	27
3.2 Lie algebras. . . . .	28
3.3 Adjoint representation. . . . .	32
3.4 Some derivatives. . . . .	33
3.5 Separable morphisms. . . . .	35
3.6 Fibres of morphisms. . . . .	36
3.7 Semisimple automorphisms. . . . .	37
<b>4. Quotients.</b>	<b>40</b>
4.1 Existence and uniqueness as a variety. . . . .	40
4.2 Quotient algebraic groups. . . . .	45

<b>5. Parabolic and Borel subgroups.</b>	<b>48</b>
5.1 Complete varieties. . . . .	48
5.2 Borel subgroups. . . . .	49
5.3 Structure of solvable groups. . . . .	52
5.4 Cartan subgroups. . . . .	57
5.5 Conjugacy of parabolic and Borel subgroups. . . . .	62
<b>6. Reductive groups.</b>	<b>65</b>
6.1 Semisimple and reductive groups. . . . .	65
6.2 Overview of the rest. . . . .	73
6.3 Reductive groups of rank 1. . . . .	75
6.4 Reductive groups of semisimple rank 1. . . . .	77
6.5 Root data. . . . .	81
<b>Appendix. An example: the symplectic group</b>	<b>85</b>

# Introduction.

**Algebraic group:** a group that is also an algebraic variety such that the group operations are maps of varieties.

*Example.*  $G = \mathrm{GL}_n(k)$ ,  $k = \bar{k}$

**Goal:** to understand the structure of reductive/semisimple affine algebraic groups over algebraically closed fields  $k$  (not necessarily of characteristic 0). Roughly, they are classified by their Dynkin diagrams, which are associated graphs.

Within  $G$  are maximal, connected, solvable subgroups, called the Borel subgroups.

*Example.* In  $G = \mathrm{GL}_n(k)$ , a Borel subgroup  $B$  is given by the upper triangular matrices.

A fundamental fact is that the Borels are conjugate in  $G$ , and much of the structure of  $G$  is grounded in those of the  $B$ . (Thus, it is important to study solvable algebraic groups).  $B$  decomposes as

$$B = T \ltimes U$$

where  $T \cong \mathbf{G}_m^n$  is a maximal torus and  $U$  is unipotent.

*Example.* With  $G = \mathrm{GL}_n(k)$ , we can take  $T$  consisting of all diagonal matrices with  $U$  the upper triangular matrices with 1's along the diagonal.

$G$  acts on its Lie algebra  $\mathfrak{g} = T_1G$ . This action restricts to a semisimple action of  $T$  on  $\mathfrak{g}$ . From the nontrivial eigenspaces, we get characters  $T \rightarrow k^\times$  called the roots. The roots give a root system, which allows us to define the Dynkin diagrams.

*Example.*  $G = \mathrm{GL}_n(k)$ .  $\mathfrak{g} = M_n(k)$  and the action of  $G$  on  $\mathfrak{g}$  is by conjugation. The roots are given by

$$\mathrm{diag}(x_1, \dots, x_n) \mapsto x_i x_j^{-1}$$

for  $1 \leq i \neq j \leq n$ .

Main References:

- Springer's *Linear Algebraic Groups*, second edition
- Polo's course notes at [www.math.jussieu.fr/~polo/M2](http://www.math.jussieu.fr/~polo/M2)
- Borel's *Linear Algebraic Groups*

# 0. Algebraic geometry (review).

$k = \bar{k}$ .

## 0.1 Zariski topology on $k^n$ .

If  $I \subset k[x_1, \dots, x_n]$  is an ideal, then  $V(I) := \{x \in k^n \mid f(x) = 0 \ \forall f \in I\}$ . Closed subsets are defined to be the  $V(I)$ . We have

$$\begin{aligned}\bigcap_{\alpha} V(I_{\alpha}) &= V(\sum I_{\alpha}) \\ V(I) \cup V(J) &= V(I \cap J)\end{aligned}$$

Note: this topology is not  $T_2$  (i.e., Hausdorff). For example, when  $n = 1$  this is the finite complement topology.

## 0.2 Nullstellensatz.

**Theorem 1** (Nullstellensatz).

(i)

$$\{\text{radical ideals } I \text{ in } k[x_1, \dots, x_n]\} \xrightleftharpoons[I]{V} \{\text{closed subsets in } k^n\}$$

are inverse bijections, where  $I(X) = \{f \in k[x_1, \dots, x_n] \mid f(x) = 0 \ \forall x \in X\}$

(ii)  $I, V$  are inclusion-reversing

(iii) If  $I \leftrightarrow X$ , then  $I$  prime  $\iff X$  irreducible.

It follows that the maximal ideals of  $k[x_1, \dots, x_n]$  are of the form

$$\mathfrak{m}_a = I(\{a\}) = (x_1 - a_1, \dots, x_n - a_n), \quad a \in k.$$

## 0.3 Some topology.

$X$  is a topological space.

$X$  is **irreducible** if  $X = C_1 \cup C_2$ , for closed sets  $C_1, C_2$  implies that  $C_i = X$  for some  $i$ .

$\iff$  any two non-empty open sets intersect

$\iff$  any non-empty open set is dense

*Facts.*

- $X$  irreducible  $\implies X$  connected.
- If  $Y \subset X$ , then  $Y$  irreducible  $\iff \bar{Y}$  irreducible.

$X$  is **noetherian** if any chain of closed subsets  $C_1 \supset C_2 \supset \dots$  stabilises. If  $X$  is noetherian, any irreducible subset is contained in a maximal irreducible subset (which is automatically closed), an **irreducible component**.  $X$  is the union of its finitely many irreducible components:

$$X = X_1 \cup \dots \cup X_n$$

*Fact.* The Zariski topology on  $k^n$  is noetherian and compact (a consequence of Nullstellansatz).

## 0.4 Functions on closed subsets of $k^n$

$X \subset k^n$  is a closed subset.

$$X = \{a \in k^n \mid \{a\} \subset X\} \iff \mathfrak{m}_a \supset I(X) \iff \{\text{maximal ideals in } k[x_1, \dots, x_n]/I(X)\}$$

Define the **coordinate ring** of  $X$  to be  $k[X] := k[x_1, \dots, x_n]/I(X)$ . The coordinate ring is a reduced, finitely-generated  $k$ -algebra and can be regarded as the restriction of polynomial functions on  $k^n$  to  $X$ .

- $X$  irreducible  $\iff k[X]$  integral domain
- The closed subsets of  $X$  are in bijection with the radical ideals of  $k[X]$ .

**Definition 2.** For a non-empty open  $U \subset X$ , define

$$\begin{aligned} \mathcal{O}_X(U) := \{f : U \rightarrow k \mid \forall x \in U, \exists x \in V \subset U, V \text{ open, and } \exists p, q \in k[x_1, \dots, x_n] \\ \text{such that } f(y) = \frac{p(y)}{q(y)} \quad \forall y \in V\} \end{aligned}$$

$\mathcal{O}_X$  is a sheaf of  $k$ -valued functions on  $X$ :

- $U \subset V$ , then  $f \in \mathcal{O}_X(V) \implies f|_U \in \mathcal{O}_X(U)$ ;
- if  $U = \bigcup U_\alpha$ ,  $f : U \rightarrow k$  function, then  $f|_{U_\alpha} \in \mathcal{O}_X(U_\alpha) \quad \forall \alpha \implies f \in \mathcal{O}_X(U)$ .

*Facts.*

- $\mathcal{O}_X(X) \cong k[X]$
- If  $f \in \mathcal{O}_X(X)$ ,  $D(f) := \{x \in X \mid f(x) \neq 0\}$  is open and these sets form a basis for the topology.  $\mathcal{O}_X(D(f)) \cong k[X]_f$ .

**Definitions 3.** A **ringed space** is a pair  $(X, \mathcal{F}_X)$  of a topological space  $X$  and a sheaf of  $k$ -valued functions on  $X$ . A **morphism**  $(X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$  of ringed spaces is a continuous map  $\phi : X \rightarrow Y$  such that

$$\forall V \subset Y \text{ open, } \forall f \in \mathcal{F}_Y(V), \quad f \circ \phi \in \mathcal{F}_X(\phi^{-1}(V))$$

An **affine variety** (over  $k$ ) is a pair  $(X, \mathcal{O}_X)$  for a closed subset  $X \subset k^n$  for some  $n$  (with  $\mathcal{O}_X$  as above). **Affine  $n$ -space** is defined as  $\mathbf{A}^n := (k^n, \mathcal{O}_{k^n})$ .

**Theorem 4.**  $X \mapsto k[X], \phi \mapsto \phi^*$  gives an equivalence of categories

$$\{\text{affine varieties over } k\}^{\text{op}} \rightarrow \{\text{reduced finitely-generated } k\text{-algebras}\}$$

If  $\phi : X \rightarrow Y$  is a morphism of varieties, then  $\phi^* : k[Y] \rightarrow k[X]$  here is  $f + I(Y) \mapsto f \circ \phi + I(X)$ . The inverse functor is given by mapping  $A$  to  $\text{m-Spec}(A)$ , the spectrum of maximal ideals of  $A$ , along with the appropriate topology and sheaf.

## 0.5 Products.

**Proposition 5.**  $A, B$  finitely-generated  $k$ -algebras. If  $A, B$  are reduced (resp. integral domains), then so is  $A \otimes_k B$ .

From the above theorem and proposition, we get that if  $X, Y$  are affine varieties, then  $\text{m-Spec}(k[X] \otimes_k k[Y])$  is a product of  $X$  and  $Y$  in the category of affine varieties.

**Remark 6.**  $X \times Y$  is the usual product as a set, but not as topological spaces (the topology is finer).

**Definition 7.** A **prevariety** is a ringed space  $(X, \mathcal{F}_X)$  such that  $X = U_1 \cup \dots \cup U_n$  with the  $U_i$  open and the  $(U_i, \mathcal{F}|_{U_i})$  isomorphic to affine varieties.  $X$  is compact and noetherian. (This is too general of a construction. Gluing two copies of  $\mathbf{A}^1$  along  $\mathbf{A}^1 - \{0\}$  (a pathological space) gives an example of a prevariety.)

Products in the category of prevarieties exist: if  $X = \bigcup_{i=1}^n U_i, Y = \bigcup_{j=1}^m V_j$  ( $U_i, V_j$  affine open), then  $X \times Y = \bigcup_{i,j}^{n,m} U_i \times V_j$ , where each  $U_i \times V_j$  is the product above. As before, this gives the usual products of sets but not topological spaces.

**Definition 8.** A prevariety is a **variety** if the diagonal  $\Delta_X \subset X \times X$  is a closed subset. (This is like being  $T_2$ !)

- Affine varieties are varieties;  $X, Y$  varieties  $\implies X \times Y$  variety.
- If  $Y$  is a variety, then the graph of a morphism  $X \rightarrow Y$  is closed in  $X \times Y$ .
- If  $Y$  is a variety,  $f, g : X \rightarrow Y$ , then  $f = g$  if  $f, g$  agree on a dense subset.

## 0.6 Subvarieties.

Let  $X$  be a variety and  $Y \subset X$  a **locally closed** subset (i.e.,  $Y$  is the intersection of a closed and an open set, or, equivalently,  $Y$  is open in  $\bar{Y}$ ). There is a unique sheaf  $\mathcal{O}_Y$  on  $Y$  such that  $(Y, \mathcal{O}_Y)$  is a prevariety and  $(Y, \mathcal{O}_Y) \rightarrow (X, \mathcal{O}_X)$  is a morphism such that

for all morphisms  $f : Z \rightarrow X$  such that  $f(Z) \subset Y$ ,  $f$  factors through the inclusion  $Y \rightarrow X$

Concretely,

$$\mathcal{O}_Y(V) = \{f : V \rightarrow k \mid \forall x \in V, \exists U \subset X, x \in U \text{ open, and } \exists \tilde{f} \in \mathcal{O}_X(U) \text{ such that } f|_{U \cap V} = \tilde{f}|_{U \cap V}\}$$

**Remarks 9.**  $Y, X$  as above.

- If  $Y \subset X$  is open, then  $\mathcal{O}_Y = \mathcal{O}_X|_Y$ .
- $Y$  is a variety ( $\Delta_Y = \Delta_X \cap (Y \times Y)$ )
- If  $X$  is affine and  $Y$  is closed, then  $Y$  is affine with  $k[Y] \cong k[X]/I(Y)$
- If  $X$  is affine and  $Y = D(f)$  is basic open, then  $Y$  is affine with  $k[Y] \cong k[X]_f$ . (Note that general open subsets of affine varieties need not be affine (e.g.,  $\mathbf{A}^2 - \{0\} \subset \mathbf{A}^2$ .)

**Theorem 10.** Let  $\phi : X \rightarrow Y$  be a morphism of affine varieties.

- (i)  $\phi^* : k[Y] \rightarrow k[X]$  surjective  $\iff \phi$  is a closed immersion (i.e., an isomorphism onto a closed subvariety)
- (ii)  $\phi^* : k[Y] \rightarrow k[X]$  is injective  $\iff \overline{\phi(X)} = Y$  (i.e.,  $\phi$  is **dominant**)

## 0.7 Projective varieties.

$\mathbf{P}^n = \frac{k^{n+1} - \{0\}}{k^\times}$  as a set. The Zariski topology on  $\mathbf{P}^n$  is given by defining, for all homogeneous ideals  $I$ ,  $V(I)$  to be a closed set. For  $U \subset \mathbf{P}^n$  open,

$$\mathcal{O}_{\mathbf{P}^n}(U) := \{f : U \rightarrow k \mid \forall x \in U \exists F, G \in k[x_0, \dots, x_n], \text{ homogeneous of the same degree} \\ \text{such that } f(y) = \frac{F(y)}{G(y)}, \text{ for all } y \text{ in a neighbourhood of } x.\}$$

Let  $U_i = \{(x_0 : \dots : x_n) \in \mathbf{P}^n \mid x_i \neq 0\} = \mathbf{P}^n - V((x_i))$ , which is open.  $\mathbf{A}^n \rightarrow U_i$  given by

$$x \mapsto (x_1 : \dots : x_{i-1} : 1 : x_i : \dots : x_n)$$

gives an isomorphism of ringed spaces, which implies that  $\mathbf{P}^n$  is a prevariety; in fact, it is an irreducible variety.

**Definitions 11.** A **projective variety** is a closed subvariety of  $\mathbf{P}^n$ . A **quasi-projective variety** is a locally closed subvariety of  $\mathbf{P}^n$ .

*Facts.*

- The natural map  $\mathbf{A}^{n+1} - \{0\} \rightarrow \mathbf{P}^n$  is a morphism
- $\mathcal{O}_{\mathbf{P}^n}(\mathbf{P}^n) = k$

## 0.8 Dimension.

$X$  here is an irreducible variety. The **function field** of  $X$  is  $k(X) := \varinjlim_{U \neq \emptyset \text{ open}} \mathcal{O}_X(U)$ , the *germs of regular functions*.

*Facts.*

- For  $U \subset X$  open,  $k(U) = k(X)$ .
- For  $U \subset X$  irreducible affine,  $k(U)$  is the fraction field of  $k[U]$ .
- $k(X)$  is a finitely-generated field extension of  $k$ .

**Definition 12.** The dimension of  $X$  is  $\dim X := \text{tr.deg}_k k(X)$ .

**Theorem 13.** If  $X$  is affine, then  $\dim X = \text{Krull dimension of } k[X]$  (which is the maximum length of chains of  $C_0 \subsetneq \cdots \subsetneq C_n$  of irreducible closed subsets).

*Facts.*

- $\dim \mathbf{A}^n = n = \dim \mathbf{P}^n$
- If  $Y \subsetneq X$  is closed and irreducible, then  $\dim Y < \dim X$
- $\dim(X \times Y) = \dim X + \dim Y$

For general varieties  $X$ , define  $\dim X := \max\{\dim Y \mid Y \text{ is an irreducible component}\}$ .

## 0.9 Constructible sets.

A subset  $A \subset X$  of a topological space is **constructible** if it is the union of finitely many locally closed subsets. Constructible sets are stable under finite unions and intersection, taking complements, and taking inverse images under continuous maps.

**Theorem 14** (Chevalley). Let  $\phi : X \rightarrow Y$  be a morphism of varieties.

- (i)  $\phi(X)$  contains a nonempty open subset of its closure.
- (ii)  $\phi(X)$  is constructible.

## 0.10 Other examples.

- A finite dimensional  $k$ -vector space is an affine variety: fix a basis to get an bijection  $V \xrightarrow{\sim} k^n$ , giving  $V$  the corresponding structure (which is actually independent of the basis chosen). Intrinsically, we can define the topology and functions using polynomials in linear forms of  $V$ , that is, from  $\text{Sym}(V^*) = \bigoplus_{n=0}^{\infty} \text{Sym}^n(V^*)$ :  $k[V] := \text{Sym}(V^*)$ .

- Similarly,  $\mathbf{P}V = \frac{V - \{0\}}{k^\times}$ . As above, use a linear isomorphism  $V \xrightarrow{\sim} k^{n+1}$  to get the structure of a projective space; or, intrinsically, use homogeneous elements of  $\text{Sym}(V^*)$ .

# 1. Algebraic groups: beginnings.

## 1.1 Preliminaries.

We will only consider the category of *affine* algebraic groups, a.k.a. **linear algebraic groups**. In future, by “algebraic group” we will mean “affine algebraic group”. There are three descriptions of the category:

(1)

**Objects:** affine varieties  $G$  over  $k$  with morphisms  $\mu : G \times G \rightarrow G$  (multiplication),  $i : G \rightarrow G$  (inversion), and  $\epsilon : \mathbf{A}^0 \rightarrow G$  (i.e., a distinguished point  $e \in G$ ) such that the *group axioms* hold, i.e., that the following diagrams commute.

$$\begin{array}{ccc}
 G \times G \times G & \xrightarrow{\mu \times \text{id}} & G \times G \\
 \text{id} \times \mu \downarrow & & \downarrow \mu \\
 G \times G & \xrightarrow{\mu} & G
 \end{array}
 \quad
 \begin{array}{ccc}
 G \times \mathbf{A}^0 & \xrightarrow{\text{id} \times \epsilon} & G \times G \xleftarrow{\epsilon \times \text{id}} \mathbf{A}^0 \times G \\
 & \searrow & \downarrow \mu \swarrow \\
 & & G
 \end{array}
 \quad
 \begin{array}{ccccc}
 G & \xrightarrow{(\text{id}, i)} & G \times G & \xleftarrow{(i, \text{id})} & G \\
 \downarrow & & \downarrow \mu & & \downarrow \\
 \mathbf{A}^0 & \xrightarrow{\epsilon} & G & \xleftarrow{\epsilon} & \mathbf{A}^0
 \end{array}$$

**Maps:** morphisms of varieties compatible with the above structure maps.

(2)

**Objects:** commutative Hopf  $k$ -algebras, which are reduced, commutative, finitely-generated  $k$ -algebras  $A$  with morphisms  $\Delta : A \rightarrow A \otimes A$  (comultiplication),  $i : A \rightarrow A$  (co-inverse), and  $\epsilon : A \rightarrow k$  (co-inverse) such that the *cogroup axioms* hold, i.e., that the following diagrams commute:

$$\begin{array}{ccc}
 A \otimes A \otimes A & \xleftarrow{\Delta \otimes \text{id}} & A \otimes A \\
 \text{id} \otimes \Delta \uparrow & & \uparrow \Delta \\
 A \otimes A & \xleftarrow{\Delta} & A
 \end{array}
 \quad
 \begin{array}{ccc}
 A \otimes k & \xleftarrow{\text{id} \otimes \epsilon} & A \otimes A \xrightarrow{\epsilon \otimes \text{id}} k \otimes A \\
 & \searrow & \uparrow \Delta \swarrow \\
 & & A
 \end{array}
 \quad
 \begin{array}{ccccc}
 A & \xleftarrow{(i, \text{id})} & A \otimes A & \xrightarrow{(\text{id}, i)} & A \\
 \uparrow & & \uparrow \Delta & & \uparrow \\
 k & \xleftarrow{\epsilon} & A & \xrightarrow{\epsilon} & k
 \end{array}$$

**Maps:**  $k$ -algebra morphisms compatible with the above structure maps.

(3)

**Objects:** representable functors

$$\left( \text{reduced finitely-generated } k\text{-algebras} \right) \rightarrow \left( \text{groups} \right)$$

**Maps:** natural transformations.

Here are the relationships:

- (1)  $\leftrightarrow$  (2):  $G \mapsto A = k[G]$  gives an equivalence of categories. Note that  $k[G \times G] = k[G] \otimes k[G]$ .  
 (2)  $\leftrightarrow$  (3):  $A \mapsto \text{Hom}_{\text{alg}}(A, -)$  gives an equivalence of categories by Yoneda's lemma.

*Examples.*

- $G = \mathbf{A}^1 =: \mathbf{G}_a$

In (1):  $\mu : (x, y) \mapsto x + y$  (sum of projections),  $i : x \mapsto -x$ ,  $\epsilon : * \mapsto 0$

In (2):  $A = k[T]$ ,  $\Delta(T) = T \otimes 1 + 1 \otimes T$ ,  $i(T) = -T$ ,  $\epsilon(T) = 0$

In (3): the functor  $\text{Hom}_{\text{alg}}(k[T], -)$  sends an algebra  $R$  to its additive group  $(R, +)$ .

- $G = \mathbf{A}^1 - \{0\} =: \mathbf{G}_m = \text{GL}_1$

In (1):  $\mu : (x, y) \mapsto xy$  (product of projections),  $i : x \mapsto x^{-1}$ ,  $\epsilon : * \mapsto 1$

In (2):  $A = k[T, T^{-1}]$ ,  $\Delta(T) = T \otimes T$ ,  $i(T) = T^{-1}$ ,  $\epsilon(T) = 1$

In (3): the functor  $\text{Hom}_{\text{alg}}(k[T, T^{-1}], -)$  sends an algebra  $R$  to its group of units  $(R, \times)$ .

- $G = \text{GL}_n$

In (1):  $\text{GL}_n(k) \subset M_n(k) \cong k^{n^2}$  with the usual operations is the basic open set given by  $\det \neq 0$

In (2):  $A = k[T_{ij}, \det(T_{ij})^{-1}]_{1 \leq i, j \leq n}$ ,  $\Delta(T_{ij}) = \sum_k T_{ik} \otimes T_{kj}$

In (3): the functor  $R \mapsto \text{GL}_n(R)$

- $G = V$  finite-dimensional  $k$ -vector space

Given by the functor  $R \mapsto (V \otimes_k R, +)$

- $G = \text{GL}(V)$ , for a finite-dimensional  $k$ -vector space  $V$

Given by the functor  $R \mapsto \text{GL}(V \otimes_k R)$

*Examples of morphisms.*

- For  $\lambda \in k^\times$ ,  $x \mapsto \lambda x$  is an automorphism of  $\mathbf{G}_a$

**Exercise.** Show that  $\text{Aut}(\mathbf{G}_a) \cong k^\times$ . Note that  $\text{End}(\mathbf{G}_a)$  can be larger, as we have the Frobenius  $x \mapsto x^p$  when  $\text{char } k = p > 0$ .

- For  $n \in \mathbf{Z}$ ,  $x \mapsto x^n$  gives an automorphism of  $\mathbf{G}_m$ .
- $g \mapsto \det g$  gives a morphism  $\text{GL}_n \rightarrow \mathbf{G}_m$ .

Note that if  $G, H$  are algebraic groups, then so is  $G \times H$  (in the obvious way).

## 1.2 Subgroups.

A *locally closed subgroup*  $H \leq G$  is a locally closed subvariety that is also a subgroup.  $H$  has a unique structure as an algebraic group such that the inclusion  $H \rightarrow G$  is a morphism (it is given by restricting the multiplication and inversion maps of  $G$ ).

*Examples.* Closed subgroups of  $\mathrm{GL}_n$ :

- $G = \mathrm{SL}_n$ , ( $\det = 1$ )
- $G = D_n$ , diagonal matrices ( $T_{ij} = 0 \ \forall i \neq j$ )
- $G = B_n$ , upper-triangular matrices ( $T_{ij} = 0 \ \forall i > j$ )
- $G = U_n$ , unipotent matrices (upper-triangular with 1's along the diagonal)
- $G = \mathrm{O}_n$  or  $\mathrm{Sp}_n$ , for a particular  $J \in \mathrm{GL}_n$  with  $J^t = \pm J$ , these are the matrices  $g$  with  $g^t J g = J$
- $G = \mathrm{SO}_n = \mathrm{O}_n \cap \mathrm{SL}_n$

**Exercise.**  $D_n \cong \mathbf{G}_m^n$ . Multiplication  $(d, n) \mapsto dn$  gives an isomorphism  $D_n \times U_n \rightarrow B_n$  as varieties. (Actually,  $B_n$  is a semidirect product of the two, with  $U_n \trianglelefteq B_n$ .)

**Remark 15.**  $\mathbf{G}_a$ ,  $\mathbf{G}_m$ , and  $\mathrm{GL}_n$  are irreducible (latter is dense in  $\mathbf{A}^{n^2}$ ).  $\mathrm{SL}_n$  is irreducible, as it is defined by the irreducible polynomial  $\det - 1$ . In fact,  $\mathrm{SO}_n, \mathrm{Sp}_n$  are also irreducible.

**Lemma 16.**

- (a) If  $H \leq G$  is an (abstract) subgroup, then  $\overline{H}$  is a (closed) subgroup.
- (b) If  $H \leq G$  is a locally closed subgroup, then  $H$  is closed.
- (c) If  $\phi : G \rightarrow H$  is a morphism of algebraic groups, then  $\ker \phi$ ,  $\mathrm{im} \phi$  are closed subgroups.

*Proof.*

(a). Multiplication by  $g$  is an isomorphism of varieties  $G \rightarrow G$ :  $g\overline{H} = \overline{gH}$  and  $\overline{H}g = \overline{Hg} \implies \overline{H} \cdot \overline{H} \subset \overline{H}$ . Inversion is an isomorphism of varieties  $G \rightarrow G$ :  $(\overline{H})^{-1} = \overline{H^{-1}} = \overline{H}$ .

(b).  $H \subset \overline{H}$  is open and  $\overline{H} \subset G$  is closed, so without loss of generality suppose that  $H \subset G$  is open. Since the complement of  $H$  is a union of cosets of  $H$ , which are open since  $H$  is, it follows that  $H$  is closed.

(c).  $\ker \phi$  is clearly a closed subgroup.  $\mathrm{im} \phi = \phi(G)$  contains a nonempty open subset  $U \subset \overline{\phi(G)}$  by Chevalley; hence,  $\phi(G) = \bigcup_{h \in \phi(G)} hU$  is open in  $\overline{\phi(G)}$  and so  $\phi(G)$  is closed by (b).  $\square$

**Lemma 17.** *The connected component  $G^0$  of the identity  $e \in G$  is irreducible. The irreducible and connected components of  $G^0$  coincide and they are the cosets of  $G^0$ .  $G^0$  is an open normal subgroup (and thus has finite index).*

*Proof.* Let  $X$  be an irreducible component containing  $e$  (which must be closed). Then  $X \cdot X^{-1} = \mu(X \times X^{-1})$  is irreducible and contains  $X$ ; hence,  $X = X \cdot X^{-1}$  is a subgroup as it is closed under inverse and multiplication. So  $G = \coprod_{gX \in G/X} gX$  gives a decomposition of  $G$  into its irreducible components. Since  $G$  has a finite number of irreducible components, it follows that  $(G : X) < \infty$  and  $X$  is open. Hence, the cosets  $gX$  are the connected components:  $X = G^0$ . Moreover,  $G^0$  is normal since  $gG^0g^{-1}$  is another connected component containing  $e$ .  $\square$

**Corollary 18.**  $G$  connected  $\iff G$  irreducible

**Exercise.**  $\phi : G \rightarrow H \implies \phi(G^0) = \phi(G)^0$

### 1.3 Commutators.

**Proposition 19.** *If  $H, K$  are closed, connected subgroups of  $G$ , then*

$$[H, K] = \langle [h, k] = hkh^{-1}k^{-1} \mid h \in H, k \in K \rangle$$

*is closed and connected. (Actually, we just need one of  $H, K$  to be connected. Moreover, without any of the connected hypotheses, Borel shows that  $[H, K]$  is closed.)*

**Lemma 20.** *Let  $\{X_\alpha\}_{\alpha \in I}$  be a collection of irreducible varieties and  $\{\phi_\alpha : X_\alpha \rightarrow G\}$  a collection of morphisms into  $G$  such that  $e \in Y_\alpha := \phi_\alpha(X_\alpha)$  for all  $\alpha$ . Then the subgroup  $H$  of  $G$  generated by the  $Y_\alpha$  is connected and closed. Furthermore,  $\exists \alpha_1, \dots, \alpha_n \in I, \epsilon_1, \dots, \epsilon_n \in \{\pm 1\}$  such that  $H = Y_{\alpha_1}^{\epsilon_1} \cdots Y_{\alpha_n}^{\epsilon_n}$ .*

*Proof of Lemma.* Without loss of generality suppose that  $\phi_\alpha^{-1} = i \circ \phi_\alpha : X_\alpha \rightarrow G$  is also among the maps for all  $\alpha$ . For  $n \geq 1$  and  $a \in I^n$ , write  $Y_a := Y_{\alpha_1} \cdots Y_{\alpha_n} \subset G$ .  $Y_a$  is irreducible, and so  $\overline{Y}_a$  is as well. Choose  $n, a$  such that  $\dim \overline{Y}_a$  is maximal. Then for all  $m, b \in I^m$ ,

$$\overline{Y}_a \subset \overline{Y}_a \cdot \overline{Y}_b \subset \overline{Y_a \cdot Y_b} = \overline{Y}_{(a,b)}$$

(second inclusion as in Lemma 1.(a)) which by maximality implies that  $\overline{Y}_a = \overline{Y_{(a,b)}}$  and  $\overline{Y}_b \subset \overline{Y}_a$ . In particular, this gives that

$$\overline{Y}_a \cdot \overline{Y}_a \subset \overline{Y_{(a,a)}} = \overline{Y}_a \quad \text{and} \quad \overline{Y}_a^{-1} \subset \overline{Y}_a$$

$\overline{Y}_a$  is a subgroup. By Chevalley, there is a nonempty  $U \subset Y_a$  open in  $\overline{Y}_a$ .

*Claim:*  $\overline{Y}_a = U \cdot U$  ( $\implies \overline{Y}_a = Y_a \cdot Y_a = Y_{(a,a)} \implies$  done.)

$$g \in \overline{Y}_a \implies gU^{-1} \cap U \neq \emptyset \implies g \in U \cdot U$$

□

*Proof of Proposition.* For  $k \in K$ , consider the morphisms  $\phi_k : H \rightarrow G, h \mapsto [h, k]$ . Note that  $\phi_k(e) = e$ . □

**Corollary 21.** *If  $\{H_\alpha\}$  are connected closed subgroups, then so is the subgroup generated by them.*

**Corollary 22.** *If  $G$  is connected, then its derived subgroup  $\mathfrak{D}G := [G, G]$  is closed and connected.*

**Definitions 23.** *Inductively define  $\mathfrak{D}^n G := \mathfrak{D}(\mathfrak{D}^{n-1} G) = [\mathfrak{D}^{n-1} G, \mathfrak{D}^{n-1} G]$  with  $\mathfrak{D}^0 G = G$ .*

$$G \supset \mathfrak{D}G \supset \mathfrak{D}^2 G \supset \cdots$$

*is the derived series of  $G$ , with each group a normal subgroup in the previous.  $G$  is solvable if  $\mathfrak{D}^n G = 1$  for some  $n \geq 0$ . Now, inductively define  $\mathcal{C}^n G := [G, \mathcal{C}^{n-1} G]$  with  $\mathcal{C}^0 G = G$ .*

$$G \supset \mathcal{C}G \supset \mathcal{C}^2 G \supset \cdots$$

*is the descending central series of  $G$ , with each group normal in the previous.  $G$  is nilpotent if  $\mathcal{C}^n G = 1$  for some  $n \geq 0$ .*

Recall the following facts of group theory:

- nilpotent  $\implies$  solvable
- $G$  solvable (resp. nilpotent)  $\implies$  subgroups, quotients of  $G$  are solvable (resp. nilpotent)
- If  $N \trianglelefteq G$ , then  $N$  and  $G/N$  solvable  $\implies G$  solvable.

*Examples.*

- $B_n$  is solvable. ( $\mathfrak{D}B_n = U_n$ )
- $U_n$  is nilpotent.

## 1.4 $G$ -spaces.

A  $G$ -space is a variety  $X$  with an action of  $G$  on  $X$  (as a set) such that  $G \times X \rightarrow X$  is a morphism of varieties. For each  $x \in X$  we have a morphism  $f_x : G \rightarrow X$  be given by  $g \mapsto gx$ , and for each  $g \in G$  we have an isomorphism  $t_g : X \rightarrow X$  given by  $x \mapsto gx$ .  $\text{Stab}_G(x) = f_x^{-1}(\{x\})$  is a closed subgroup.

*Examples.*

- $G$  acts on itself by  $g * x = gx$  or  $xg^{-1}$  or  $g x g^{-1}$ . (Note that in the case of the last action,  $\text{Stab}(x) = \mathcal{Z}_G(x)$  is closed and so the center  $\mathcal{Z}_G = \bigcap_{x \in G} \mathcal{Z}_G(x)$  is closed.)
- $\text{GL}(V) \times V \rightarrow V$ ,  $(g, x) \mapsto g(x)$
- $\text{GL}(V) \times \mathbf{P}V \rightarrow \mathbf{P}V$  (exercise)

### Proposition 24.

- Orbits are locally closed (so each orbit is a subvariety and is itself a  $G$ -space).
- There exists a closed orbit.

*Proof.*

(a). Let  $Gx$  be an orbit, which is the image of  $f_x$ . By Chevalley, there is a nonempty  $U \subset Gx$  open in  $\overline{Gx}$ . Then  $Gx = \bigcup_{g \in G} gU$  is open in  $\overline{Gx}$ .

(b). Since  $X$  is noetherian, we can choose an orbit  $Gx$  such that  $\overline{Gx}$  is minimal (with respect to inclusion). We will show that  $Gx$  is closed. Suppose otherwise. Then  $\overline{Gx} - Gx$  is nonempty, closed in  $\overline{Gx}$  by (a), and  $G$ -stable (by the usual argument); let  $y$  be an element in the difference. But then  $\overline{Gy} \subsetneq \overline{Gx}$ . Contradiction. Hence,  $Gx$  is closed.  $\square$

**Lemma 25.** *If  $G$  is irreducible, then  $G$  preserves all irreducible components of  $X$ .*

Exercise.

Suppose  $\theta : G \times X \rightarrow X$  gives an affine  $G$ -space. Then  $G$  acts linearly on  $k[X]$  by

$$(g \cdot f)(x) := f(g^{-1}x), \quad \text{i.e., } g \cdot f = t_{g^{-1}}^*(f)$$

**Definitions 26.** Suppose a group  $G$  acts linearly on a vector space  $W$ . Say the action is **locally finite** if  $W$  is the union of finite-dimensional  $G$ -stable subspaces. If  $G$  is an algebraic group, say the action is **locally algebraic** if it is locally finite and, for any finite-dimensional  $G$ -stable subspace  $V$ , the action  $\theta : G \times V \rightarrow V$  is a morphism.

**Proposition 27.** The action of  $G$  on  $k[X]$  is locally algebraic. Moreover, for all finite-dimensional  $G$ -stable  $V \subset k[X]$ , then  $\theta^*(V) \subset k[G] \otimes V$ .

*Proof.*  $t_{g^{-1}}$  factors as

$$\begin{aligned} t_{g^{-1}} : X &\rightarrow G \times X \xrightarrow{\theta} X \\ x &\mapsto (g^{-1}, x) \\ t_{g^{-1}}^* : k[X] &\xrightarrow{\theta^*} k[G] \otimes k[X] \xrightarrow{(\text{ev}_{g^{-1}}, \text{id})} k[X] \end{aligned}$$

Fix  $f \in k[X]$  and write  $\theta^*(f) = \sum_{i=1}^n h_i \otimes f_i$ , so

$$g \cdot f = t_{g^{-1}}^*(f) = \sum_{i=1}^n h_i(g^{-1})f_i$$

Hence, the  $G$ -orbit of  $f$  is contained in  $\sum_{i=1}^n k f_i$ , implying local finiteness.

Let  $V \subset k[X]$  be finite-dimensional and  $G$ -stable, and pick basis  $(e_i)_{i=1}^n$ . Extend the  $e_i$  to a basis  $\{e_i\}_i \cup \{e'_\alpha\}_\alpha$  of  $k[X]$ . Write

$$\begin{aligned} \theta^* e_i &= \sum_j h_{ij} \otimes e_j + \sum_\alpha h'_{i\alpha} \otimes e'_\alpha \\ \implies g \cdot e_i &= \sum_j h_{ij}(g^{-1})e_j + \sum_\alpha h'_{i\alpha}(g^{-1})e'_\alpha \in V \\ \implies h'_{i\alpha}(g^{-1}) &= 0 \quad \forall g, i, \alpha \\ \implies h'_{i\alpha} &= 0 \quad \forall i, \alpha \end{aligned}$$

Hence,  $\theta^*(V) \subset k[G] \otimes V$ . Moreover, we see that  $G \times V \rightarrow V$  is a morphism, as it is given by

$$(g, \sum_i \lambda_i e_i) \mapsto \sum_{i,j} \lambda_j h_{ij}(g^{-1})e_j$$

It follows that the action of  $G$  on  $k[X]$  is locally algebraic. □

**Theorem 28** (Analogue of Cayley's Theorem). Any algebraic group is isomorphic to a closed subgroup of some  $\text{GL}_n$ .

*Proof.*  $G$  acts on itself by right translation, so  $(g \cdot f)(\gamma) = f(\gamma g)$ . By Proposition 7 we know that this gives a locally algebraic action on  $k[G]$ . Let  $f_1, \dots, f_n$  be generators of  $k[G]$ . Without loss of generality, the  $f_i$  are linearly independent and  $V = \sum_{i=1}^n k f_i$  is  $G$ -stable. Write

$$g \cdot f_i = \sum_j h_{ji}(g^{-1})f_j = \sum_j h'_{ji}(g)f_j$$

where  $h_{ji} \in k[G]$  and  $h'_{ji} : g \mapsto h_{ji}(g^{-1})$ . It follows that  $\phi : G \rightarrow \text{GL}(V)$  given by  $g \mapsto (h'_{ij}(g))$  is a morphism of algebraic groups. It remains to show that  $\phi$  is a closed immersion.

We have  $h'_{ij} \in \text{im } \phi^*$  for all  $i, j$ , as they are the image of projections. Moreover,

$$f_i(g) = (g \cdot f_i)(e) = \sum_j h'_{ji}(g) f_j(e) \implies f_i \in \sum_j k h'_{ji} \subset \text{im } \phi^*$$

Since the  $f_i$  generate  $k[G]$ , it follows that  $\phi^*$  is surjective; that is,  $\phi$  is a closed immersion.  $\square$

## 1.5 Jordan Decomposition.

Let  $V$  be a finite-dimensional  $k$ -vector space.  $\alpha \in \text{GL}(V)$  is **semisimple** if it is diagonalisable, and is **unipotent** if 1 is its only eigenvalue. If  $\alpha, \beta$  commute then

$$\alpha \text{ and } \beta \text{ semisimple (resp. unipotent)} \implies \alpha\beta \text{ semisimple (resp. unipotent)}$$

**Proposition 29.**  $\alpha \in \text{GL}(V)$

- (i)  $\exists! \alpha_s$  (semisimple),  $\alpha_u$  (unipotent)  $\in \text{GL}(V)$  such that  $\alpha = \alpha_s \alpha_u = \alpha_u \alpha_s$ .
- (ii)  $\exists p_s(x), p_u(x) \in k[X]$  such that  $\alpha_s = p_s(\alpha)$ ,  $\alpha_u = p_u(\alpha)$ .
- (iii) If  $W \subset V$  is an  $\alpha$ -stable subspace, then

$$\begin{aligned} (\alpha|_W)_s &= \alpha_s|_W, & (\alpha|_{V/W})_s &= \alpha_s|_{V/W} \\ (\alpha|_W)_u &= \alpha_u|_W, & (\alpha|_{V/W})_u &= \alpha_u|_{V/W} \end{aligned}$$

- (iv) If  $f : V_1 \rightarrow V_2$  linear with  $\alpha_i \in \text{GL}(V_i)$  for  $i = 1, 2$ , then

$$f \circ \alpha_1 = \alpha_2 \circ f \implies \begin{cases} f \circ (\alpha_1)_s = (\alpha_2)_s \circ f \\ f \circ (\alpha_1)_u = (\alpha_2)_u \circ f \end{cases}$$

- (v) If  $\alpha_i \in \text{GL}(V_i)$  for  $i = 1, 2$ , then

$$\begin{aligned} (\alpha_1 \otimes \alpha_2)_s &= (\alpha_1)_s \otimes (\alpha_2)_s \\ (\alpha_1 \otimes \alpha_2)_u &= (\alpha_1)_u \otimes (\alpha_2)_u \end{aligned}$$

*Proof sketch.*

(i) - existence:

A Jordan block for an eigenvalue  $\lambda$  decomposes as

$$\begin{pmatrix} \lambda & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & \lambda \end{pmatrix} = \begin{pmatrix} \lambda & & & \\ & \ddots & & \\ & & \ddots & \\ & & & \lambda \end{pmatrix} \begin{pmatrix} 1 & \lambda^{-1} & & \\ & \ddots & \ddots & \\ & & \ddots & \lambda^{-1} \\ & & & 1 \end{pmatrix}$$

The left factor is semisimple and the right is unipotent, and so they both commute.

(i) - uniqueness:

If  $\alpha = \alpha_s \alpha_u = \alpha'_s \alpha'_u$ , then  $\alpha_s^{-1} \alpha'_s = \alpha_u^{-1} \alpha'_u$  is both unipotent and semisimple, and thus is the identity.

(ii): This follows from the Chinese Remainder Theorem.

(iii): Use (ii) + uniqueness.

(iv): Since  $f : V_1 \rightarrow \text{im } f \hookrightarrow V_2$ , it suffices to consider the cases where  $f$  is injective or surjective, in which we can invoke (iii).

(v): Exercise. □

**Definition 30.** An **(algebraic)  $G$ -representation** is a linear  $G$ -action on a finite-dimension  $k$ -vector space such that  $G \times V \rightarrow V$  is a morphism of varieties, which is equivalent to  $G \rightarrow \text{GL}(V)$  being a morphism of algebraic groups. Note that if  $G \rightarrow \text{GL}(V)$  is given by  $g \mapsto (h_{ij}(g))$ , then  $G \times V \rightarrow V$  is given by  $(g, \sum_i \lambda_i e_i) \mapsto \sum_j \lambda_i h_{ji}(g) e_j$ .

**Lemma 31.** There is a  $G$ -linear map  $\eta : V \hookrightarrow V_0 \otimes k[G]$ , where  $V_0$  is  $V$  with the trivial  $G$ -action and  $G$  acts on  $k[G]$  by right translation.

*Proof.* Define  $\eta$  by  $\eta(e_i) = \sum_j e_j \otimes h_{ji}$ . The diagram

$$\begin{array}{ccc} V & \xrightarrow{\eta} & V_0 \otimes k[G] \\ & \searrow g & \downarrow (\text{id}, \text{ev}_g) \\ & & V \end{array}$$

commutes and so “ $gv = \eta(v)(g)$ ”. □

**Proposition 32.** Suppose that for all algebraic  $G$ -representations  $V$ , there is a  $\alpha_V \in \text{GL}(V)$  such that

(i)  $\alpha_{k_0} = \text{id}_V$ , where  $k_0$  is the one-dimensional trivial representation.

(ii)  $\alpha_{V \otimes W} = \alpha_V \otimes \alpha_W$

(iii) If  $f : V \rightarrow W$  is a map of  $G$ -representations, then  $\alpha_W \circ f = f \circ \alpha_V$ .

Then  $\exists! g \in G$  such that  $\alpha_V = gv$  for all  $V$ .

*Proof.* From (iii), if  $W \hookrightarrow V$  is a  $G$ -stable subspace, then  $\alpha_V|_W = \alpha_W$ . If  $V$  is a local algebraic  $G$ -representation, then  $\exists! \alpha_V$  such that  $\alpha_V|_W = \alpha_W$  for all finite-dimensional  $G$ -stable  $W \subset V$ . Note that (ii), (iii) still hold for locally algebraic representations. Also note that from (iii) it follows that  $\alpha_{V \oplus W} = \alpha_V \oplus \alpha_W$ . Define  $\alpha = \alpha_{k[G]} \in \text{GL}(k[G])$ , where  $G$  acts on  $k[G]$  by  $(gf)(\lambda) = f(\lambda g)$ .

*Claim.*  $\alpha$  is a ring automorphism.

$m : k[G] \otimes k[G] \rightarrow k[G]$  is a map of locally algebraic  $G$ -representations:  $f_1(\cdot g) f_2(\cdot g) = (f_1 f_2)(\cdot g)$ . Thus, by (ii) and (iii),  $\alpha \circ m = m \circ (\alpha \otimes \alpha)$ , and so  $\alpha(f_1 f_2) = \alpha(f_1) \alpha(f_2)$ .

Therefore, the composition  $k[G] \xrightarrow{\alpha} k[G] \xrightarrow{\text{ev}_e} k$  is a ring homomorphism and is equal to  $\text{ev}_g$  for some unique  $g$ .

*Claim.*  $\alpha(f) = gf \quad \forall f$ , i.e.,  $\alpha = g_{k[G]}$ .

By above  $\alpha(f)(e) = f(g)$ . Also, if  $\ell(\lambda)(f) := f(\lambda^{-1} \cdot)$ , then  $\ell(\lambda) : k[G] \rightarrow k[G]$  is  $G$ -linear by (iii):

$$\alpha \circ \ell(\lambda) = \ell(\lambda) \circ \alpha \implies \alpha(f)(\lambda^{-1}) = f(\lambda^{-1}g) \implies \alpha(f) = gf$$

Now if  $V$  is a  $G$ -rep,  $\eta : V \hookrightarrow V_0 \otimes k[G]$  is  $G$ -linear, by Lemma 31, and so

$$\alpha_{V_0 \otimes k[G]} \circ \eta = \eta \circ \alpha_V$$

Since

$$\alpha_{V_0 \otimes k[G]} = \alpha_{V_0} \otimes \alpha_{k[G]} = \text{id}_{V_0} \otimes g_{k[G]} = g_{V_0 \otimes k[G]}$$

and

$$g_{V_0 \otimes k[G]} \circ \eta = \eta \circ g_V$$

and the fact that  $\eta$  is injective, it follows that  $\alpha_V = g_V$ . ( $g$  is unique, as  $G \rightarrow \text{GL}(k[G])$  is injective. Exercise!)  $\square$

**Theorem 33.** *Let  $G$  be an algebraic group.*

(i)  $\forall g \in G \quad \exists! g_s, g_u \in G$  such that for all representations  $\rho : G \rightarrow \text{GL}(V)$

$$\rho(g_s) = \rho(g)_s \quad \text{and} \quad \rho(g_u) = \rho(g)_u$$

$$\text{and } g = g_s g_u = g_u g_s.$$

(ii) For all  $\phi : G \rightarrow H$

$$\phi(g_s) = \phi(g)_s \quad \text{and} \quad \phi(g_u) = \phi(g)_u$$

*Proof.*

(i). Fix  $g \in G$ . For all  $G$ -representations  $V$ , let  $\alpha_V := (g_V)_s$ . If  $f : V \rightarrow W$  is  $G$ -linear, then  $f \circ g_V = g_W \circ f$  implies that  $f \circ \alpha_V = \alpha_W \circ f$  by Proposition 29. Also,  $\alpha_{k_0} = \text{id}_s = \text{id}$ , and

$$\alpha_{V \otimes W} = (g_{V \otimes W})_s = (g_V \otimes g_W)_s = \alpha_V \otimes \alpha_W$$

(the last equality following from Proposition 29). By Proposition 32, there is a unique  $g_s \in G$  such that  $\alpha_V = (g_s)_V$  for all  $V$ , i.e.,  $\rho_V(g_s) = \rho(g)_s$ . Similarly for  $g_u$ . From a closed immersion  $G \hookrightarrow \text{GL}(V)$ , from Theorem 28, we see that  $g = g_s g_u = g_u g_s$ .

(ii). Given  $\phi : G \rightarrow H$ , let  $\rho : H \rightarrow \text{GL}(V)$  be a closed immersion. Then

$$\rho(\phi(g_*)) = \rho(\phi(g))_* = \rho(\phi(g)_*)$$

where the first equality is by (i) for  $G$  (as  $\phi \circ \rho$  makes  $V$  into a  $G$ -representation) and the second equality is by (i) for  $H$ .  $\square$

**Exercise.** What is the Jordan decomposition in  $\mathbf{G}_a$ ? How about in a finite group?

**Remark 34.**  $F : (G\text{-representations}) \rightarrow (k\text{-vector spaces})$  denotes the forgetful functor, then Proposition 32 says that

$$G \cong \text{Aut}^{\otimes}(F)$$

where the left side is the group of natural isomorphisms  $F \rightarrow F$  respecting  $\otimes$ .

## 2. Diagonalisable and elementary unipotent groups.

### 2.1 Unipotent and semisimple subsets.

**Definitions 35.**

$$G_s := \{g \in G \mid g = g_s\}$$

$$G_u := \{g \in G \mid g = g_u\}$$

Note that  $G_s \cap G_u = \{e\}$  and  $G_u$  is a closed subset of  $G$  (embedding  $G$  into a  $\mathrm{GL}_n$ ,  $G_u$  is the closed consisting of  $g$  such that  $(g - I)^n = 0$ .  $G_s$ , however, need not be closed (as in the case  $G = B_2$ )).

**Corollary 36.** If  $gh = hg$  and  $g, h \in G_*$ , then  $gh, g^{-1} \in G_*$ , where  $*$  =  $s, u$ .

**Proposition 37.** If  $G$  is commutative, then  $G_s, G_u$  are closed subgroups and  $\mu : G_s \times G_u \rightarrow G$  is an isomorphism of algebraic groups.

*Proof.*  $G_s, G_u$  are subgroups by Corollary 36 and  $G_u$  is closed by a remark above. Without loss of generality,  $G \subset \mathrm{GL}(V)$  is a closed subgroup for some  $V$ . As  $G$  is commutative,  $V = \bigoplus_{\lambda: G_s \rightarrow k^\times} V_\lambda$  (a direct sum of eigenspaces for  $G_s$ ) and  $G$  preserves each  $V_\lambda$ . Hence, we can choose a basis for each  $V_\lambda$  such that the  $G$ -action is upper-triangular (commuting matrices are simultaneously upper-triangularisable), and so  $G \subset B_n$  and  $G_s = G \cap D_n$ . Then  $G \hookrightarrow B_n$  followed by projecting to the diagonal  $D_n$  gives a morphism  $G \rightarrow G_s$ ,  $g \mapsto g_s$ ; hence,  $g \mapsto (g_s, g_s^{-1}g)$  gives a morphism  $G \rightarrow G_s \times G_u$ , one inverse to  $\mu$ .  $\square$

**Definition 38.**  $G$  is unipotent if  $G = G_u$ .

*Example.*  $U_n$  is unipotent, and so is  $\mathbf{G}_a$  (as  $\mathbf{G}_a \cong U_2$ ).

**Proposition 39.** If  $G$  is unipotent and  $\phi : G \rightarrow \mathrm{GL}_n$ , then there is a  $\gamma \in \mathrm{GL}_n$  such that  $\mathrm{im}(\gamma\phi\gamma^{-1}) \subset U_n$ .

*Proof.* We prove this by induction on  $n$ . Suppose that this true for  $m < n$ , let  $V$  be an  $n$ -dimensional vector space, and  $\phi : G \rightarrow \mathrm{GL}(V)$ . Suppose that there is a  $G$ -invariant subspace  $0 \subsetneq W_1 \subsetneq V$ . Let  $W_2$  is complementary to  $W_1$ , so that  $V = W_1 \oplus W_2$ , and let  $\phi_i : G \rightarrow \mathrm{GL}(V_i)$  be the induces morphisms for  $i = 1, 2$ , so that  $\phi = \phi_1 \oplus \phi_2$ . Since  $n > \dim W_1, \dim W_2$ , there are  $\gamma_1, \gamma_2 \in \mathrm{GL}(V)$  such that  $\mathrm{im}(\gamma_i \phi_i \gamma_i^{-1})$  consists of unipotent elements for  $i = 1, 2$ . If  $\gamma = \gamma_1 \oplus \gamma_2$ , then it follows that  $\mathrm{im}(\gamma\phi\gamma^{-1})$  consists of unipotent elements as well.

Now, suppose that there does not exist such a  $W_1$ , so that  $V$  is irreducible. For  $g \in G$

$$\begin{aligned} \operatorname{tr}(\phi(g)) = n &\implies \forall h \in G \operatorname{tr}((\phi(g) - 1)\phi(h)) = \operatorname{tr}(\phi(gh)) - \operatorname{tr}(\phi(h)) = n - n = 0 \\ &\implies \forall x \in \operatorname{End}(V) \operatorname{tr}((\phi(g) - 1)x) = 0, \text{ by Burnside's theorem} \\ &\implies \phi(g) - 1 = 0 \\ &\implies \phi(g) = 1 \\ &\implies \operatorname{im} \phi = 1 \end{aligned}$$

(Recall that Burnside's Theorem says that  $G$  spans  $\operatorname{End}(V)$  as a vector space.) □

**Corollary 40.** *Any irreducible representation of a unipotent group is trivial.*

**Corollary 41.** *Any unipotent  $G$  is nilpotent.*

*Proof.*  $U_n$  is nilpotent. □

**Remark 42.** *The converse is not true; any torus is nilpotent (the definition of a torus to come immediately.)*

## 2.2 Diagonalisable groups and tori.

**Definitions 43.**  $G$  is **diagonalisable** if  $G$  is isomorphic to a closed subgroup of  $D_n \cong \mathbf{G}_m^n$  ( $n \geq 0$ ).  $G$  is a **torus** if  $G \cong D_n$  ( $n \geq 0$ ). The **character group** of  $G$  is

$$X^*(G) := \operatorname{Hom}(G, \mathbf{G}_m) \quad (\text{morphisms of algebraic groups})$$

*It is an abelian group under multiplication ( $(\chi_1\chi_2)(g) = \chi_1(g)\chi_2(g)$ ) and is a subgroup of  $k[G]^\times$ .*

Recall the following result:

**Proposition 44** (Dedekind).  $X^*(G)$  is a linearly independent subset of  $k[G]$ .

*Proof.* Suppose that  $\sum_{i=1}^n \lambda_i \chi_i = 0$  in  $k[G]$ ,  $\lambda_i \in k$ . Without loss of generality,  $n \geq 2$  is minimal among all possible nontrivial linear combinations (so that  $\lambda_i \neq 0 \forall i$ ). Then

$$\begin{aligned} \forall g, h, \quad &\begin{cases} 0 = \sum \lambda_i \chi_i(g)\chi_i(h) \\ 0 = \sum \lambda_i \chi_i(g)\chi_n(h) \end{cases} \\ \implies \forall h, \quad &0 = \sum_{i=1}^{n-1} \lambda_i [\chi_i(h) - \chi_n(h)]\chi_i \end{aligned}$$

By the minimality of  $n$ , we must have that the coefficients are all 0; that is,  $\forall i, h \chi_i(h) = \chi_n(h) \implies \chi_i = \chi_n$ . We still arrive at a contradiction. □

**Proposition 45.** *The following are equivalent:*

- (i)  $G$  is diagonalisable.
- (ii)  $X^*(G)$  is a basis of  $k[G]$  and  $X^*(G)$  is finitely-generated.

(iii)  $G$  is commutative and  $G = G_s$ .

(iv) Any  $G$ -representation is a direct sum of 1-dimensional representations

*Proof.*

(i)  $\Rightarrow$  (ii): Fix an embedding  $G \hookrightarrow D_n$ .  $k[D_n] = k[T_1^{\pm 1}, \dots, T_n^{\pm 1}]$  - as seen from restricting  $T_{ij}, \det(T_{ij})^{-1} \in k[\mathrm{GL}_n]$  - has a basis of monomials  $T_1^{a_1} \cdots T_n^{a_n}$ ,  $a_i \in \mathbf{Z}$ , each of which is in  $X^*(G)$ :

$$\mathrm{diag}(x_1, \dots, x_n) \mapsto x_1^{a_1} \cdots x_n^{a_n}$$

Hence,  $X^*(D_n) \cong \mathbf{Z}^n$  (by Proposition 44). The closed immersion  $G \rightarrow D_n$  gives a surjection  $k[D_n] \rightarrow k[G]$ , inducing a map  $X^*(D_n) \rightarrow X^*(G)$ ,  $\chi \mapsto \chi|_G$ .  $\mathrm{im}(X^*(D_n) \rightarrow X^*(G))$  spans  $k[G]$  and is contained in  $X^*(G)$ , which is linearly independent. Hence,  $X^*(G)$  is a basis of  $k[G]$  and we have the surjection

$$\mathbf{Z}^n \cong X^*(D_n) \twoheadrightarrow X^*(G)$$

implying the finite-generation.

(ii)  $\Rightarrow$  (iii): Say  $\chi_1, \dots, \chi_n$  by generators of  $X^*(G)$ . Define the morphism  $\phi : G \rightarrow \mathrm{GL}_n$  by  $g \mapsto \mathrm{diag}(\chi_1(g), \dots, \chi_n(g))$ .

$$\begin{aligned} g \in \ker \phi &\implies \chi_i(g) = 1 \quad \forall i \\ &\implies \chi(g) = 1 \quad \forall \chi \in X^*(G) \\ &\implies f(g) = 0 \quad \forall f \in M_e = \{g = \sum_{\chi} \lambda_{\chi} \chi \in k[X] \mid 0 = g(e) = \sum_{\chi} \lambda_{\chi}\} \\ &\implies M_e \subset M_g \\ &\implies M_e = M_g \\ &\implies g = e \end{aligned}$$

So  $\phi$  is injective, which implies that  $G$  is commutative and  $G = G_s$ .

(iii)  $\Rightarrow$  (iv): Let  $\phi : G \rightarrow \mathrm{GL}_n$  be a representation.  $\mathrm{im} \phi$  is a commuting set of diagonalisable elements, which means we can simultaneously diagonalise them.

(iv)  $\Rightarrow$  (i): Pick  $\phi : G \hookrightarrow \mathrm{GL}_n$  (Theorem 28). By (iii), without loss of generality, suppose that  $\mathrm{im} \phi \subset D_n$ . Hence,  $\phi : G \hookrightarrow D_n$ .  $\square$

**Corollary 46.** *Subgroups and images under morphisms of diagonalisable groups are diagonalisable.*

*Proof.* (iii).  $\square$

*Observations:*

- $\mathrm{char} k = p \implies X^*(G)$  has no  $p$ -torsion.
- $k[G] \cong k[X^*(G)]$  as algebras ( $k[X^*(G)]$  being a group algebra).
- For  $\chi \in X^*(G)$ ,

$$\Delta(\chi) = \chi \otimes \chi, \quad i(\chi) = \chi^{-1}, \quad \epsilon(\chi) = 1$$

Indeed,

$$\begin{aligned}\Delta(\chi)(g_1, g_2) &= \chi(g_1 g_2) = \chi(g_1)\chi(g_2) = (\chi \otimes \chi)(g_1, g_2) \\ i(\chi)(g) &= \chi(g^{-1}) = \chi(g)^{-1} = \chi^{-1}(g) \\ \epsilon(\chi) &= \chi(e) = 1\end{aligned}$$

**Theorem 47.** *Let  $p = \text{char } k$ .*

$\left( \text{diagonalisable algebraic groups} \right) \xrightarrow{X^*} \left( \text{finitely-generated abelian groups (with no } p\text{-torsion if } p > 0) \right)$

$$\begin{array}{ccc} G & \longrightarrow & X^*(G) \\ \downarrow & & \uparrow \\ H & \longrightarrow & X^*(H) \end{array}$$

is a (contravariant) equivalence of categories.

*Proof.* It is well-defined by the above. We will define an inverse functor  $F$ . Given  $X \cong \mathbf{Z}^\oplus \bigoplus_{i=1}^s \mathbf{Z}/n_i \mathbf{Z}$  from the category on the right, we have that its group algebra  $k[X]$  is finitely-generated and reduced:

$$k[X] \cong k[\mathbf{Z}]^{\otimes r} \otimes \bigotimes_{i=1}^s k[\mathbf{Z}/n_i \mathbf{Z}] \cong k[T^{\pm 1}]^{\otimes r} \otimes \bigotimes_{i=1}^s k[T]/(T^{n_i} - 1)$$

Moreover,  $k[X]$  is a Hopf algebra, which is easily checked, defining

$$\Delta : e_x \mapsto e_x \otimes e_x, \quad i : e_x \mapsto e_{x^{-1}} = e_x^{-1}, \quad \epsilon : e_x \mapsto 1$$

where  $X$  has been written multiplicatively and  $k[X] = \bigoplus_{x \in X} k e_x$ . Define  $F$  by  $F(X) = \text{m-Spec}(k[X])$ . Above, we saw that  $F X^*(G) \cong G$  as algebraic groups.

$$\begin{aligned} X^*(F(X)) &= \text{Hom}(F(X), \mathbf{G}_m) \\ &= \text{Hom}_{\text{Hopf-alg}}(k[T, T^{-1}], k[X]) \\ &= \{ \lambda \in k[X]^\times \text{ (corresponding to the images of } T) \mid \Delta(\lambda) = \lambda \otimes \lambda \} \end{aligned}$$

For an element above, write  $\lambda = \sum_{x \in X} \lambda_x e_x$  (almost all of the  $\lambda_x \in k$  of course being zero). Then

$$\Delta(\lambda) = \sum_x \lambda_x (e_x \otimes e_x) \quad \text{and} \quad \lambda \otimes \lambda = \sum_{x, x'} \lambda_x \lambda_{x'} (e_x \otimes e_{x'})$$

Hence,

$$\lambda_x \lambda_{x'} = \begin{cases} \lambda_x, & x = x' \\ 0, & x \neq x' \end{cases}$$

So,  $\lambda_x \neq 0$  for an *unique*  $x \in X$ , and

$$\lambda_x^2 = \lambda \implies \lambda_x = 1 \implies \lambda = e_x \in X$$

Thus we have  $X^*(F(X)) \cong X$  as abelian groups. The two functors are inverse on maps as well, as is easily checked.  $\square$

**Corollary 48.**

- (i) *The diagonalisable groups are the groups  $\mathbf{G}_m^r \times H$ , where  $H$  is a finite group of order prime to  $p$ .*
- (ii) *For a diagonalisable group  $G$ ,*

$$G \text{ is a torus} \iff G \text{ is connected} \iff X^*(G) \text{ is free abelian}$$

*Proof.* Define  $\mu_n := \ker(\mathbf{G}_m \xrightarrow{n} \mathbf{G}_m)$ , which is diagonalisable. If  $(n, p) = 1$ , then  $k[\mu_n] = k[T]/(T^n - 1)$  ( $T^n - 1$  is separable) and  $X^*(\mu_n) \cong \mathbf{Z}/n\mathbf{Z}$ . Since  $X^*(\mathbf{G}_m) \cong \mathbf{Z}$  and  $X^*(G \times H) \cong X^*(G) \oplus X^*(H)$ , the result follows from Theorem 47.  $\square$

**Corollary 49.**  $\text{Aut}(D_n) \cong \text{GL}_n(\mathbf{Z})$

*Fact/Exercise.* If  $G$  is diagonalisable, then

$$G \times X^*(G) \rightarrow \mathbf{G}_m, (g, \chi) \mapsto \chi(g)$$

is a “perfect bilinear pairing”, i.e., it induces isomorphisms  $X^*(G) \xrightarrow{\sim} \text{Hom}(G, \mathbf{G}_m)$  and  $G \xrightarrow{\sim} \text{Hom}_{\mathbf{Z}}(X^*(G), \mathbf{G}_m)$  (as abelian groups). Moreover, it induces inverse bijections

$$\begin{aligned} \{ \text{closed subgroups of } G \} &\longleftrightarrow \{ \text{subgroups } Y \text{ of } X^*(G) \text{ such that } X^*(G)/Y \text{ has no } p\text{-torsion} \} \\ H &\longmapsto H^\perp \\ Y^\perp &\longleftarrow Y \end{aligned}$$

$\square$

*Fact.* Say

$$1 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 1$$

is **exact** if the sequence is set-theoretically exact and the induced sequence of lie algebras

$$0 \rightarrow \text{Lie } G_1 \rightarrow \text{Lie } G_2 \rightarrow \text{Lie } G_3 \rightarrow 0$$

is exact. (See Definition 90.) Suppose the  $G_i$  are diagonalisable, so that  $\text{Lie } G_i \cong \text{Hom}_{\mathbf{Z}}(X^*(G_i), k)$ . Then the sequence of the  $G_i$  is exact if and only if

$$0 \rightarrow X^*(G_3) \rightarrow X^*(G_2) \rightarrow X^*(G_1) \rightarrow 0$$

**Remark 50.**

$$1 \rightarrow \mu_p \rightarrow \mathbf{G}_m \xrightarrow{p} \mathbf{G}_m \rightarrow 1$$

is set-theoretically exact, but

$$0 \rightarrow X^*(\mathbf{G}_m) \xrightarrow{p} X^*(\mathbf{G}_m) \rightarrow X^*(\mu_p) \rightarrow 0$$

is not if  $\text{char } k = p$  (in which case  $X^*(\mu_p) = 0$ ).

**Definition.** The group of **cocharacters** of  $G$  are

$$X_*(G) := \text{Hom}(\mathbf{G}_m, G)$$

If  $G$  is abelian, then  $X_*(G)$  is an abelian group.

**Proposition 51.** *If  $T$  is a torus, then  $X_*(T), X^*(T)$  are free abelian and*

$$X^*(T) \times X_*(T) \rightarrow \text{Hom}(\mathbf{G}_m, \mathbf{G}_m) \cong \mathbf{Z}, \quad (\chi, \lambda) \mapsto \chi \circ \lambda$$

*is a perfect pairing.*

*Proof.*

$$X_*(T) = \text{Hom}(\mathbf{G}_m, T) \cong \text{Hom}(X^*(T), \mathbf{Z}).$$

The isomorphism follows from Theorem 47. Since  $X^*(T)$  is finitely-generated free abelian by Corollary 48, we have that  $X_*(T) \cong \text{Hom}(X^*(T), \mathbf{Z})$  is free abelian as well. Moreover, since

$$\text{Hom}(X, \mathbf{Z}) \times X \rightarrow \mathbf{Z}, \quad (\alpha, x) \mapsto \alpha(x)$$

is a perfect pairing for any finitely-generated free abelian  $X$ , it follows from the isomorphism above that the pairing in question is also perfect.  $\square$

**Proposition 52** (Rigidity of diagonalisable groups). *Let  $G, H$  be diagonalisable groups and  $V$  a connected affine variety. If  $\phi : G \times V \rightarrow H$  is a morphism of varieties such that  $\phi_v : G \rightarrow H, g \mapsto \phi(g, v)$  is a morphism of algebraic groups for all  $v \in V$ , then  $\phi_v$  is independent of  $v$ .*

Under  $\phi^* : k[H] \rightarrow k[G] \otimes k[V]$ , for  $\chi \in X^*(H)$ , write

$$\phi^*(\chi) = \sum_{\chi' \in X^*(G)} \chi' \otimes f_{\chi\chi'}$$

Then

$$\begin{aligned} \phi_v^*(\chi) = \sum_{\chi'} f_{\chi\chi'}(v)\chi \in X^*(G) &\implies \forall \chi', v \quad f_{\chi\chi'}(v) \in \{0, 1\} \\ &\implies \forall \chi' \quad f_{\chi\chi'}^2 = f_{\chi\chi'} \\ &\implies \forall \chi' \quad V = V(f_{\chi\chi'}) \sqcup V(1 - f_{\chi\chi'}) \\ &\implies \forall \chi' \quad f_{\chi\chi'} \text{ is constant, since } V \text{ is connected} \\ &\implies \forall \phi_v \text{ is independent of } v \end{aligned}$$

$\square$

**Corollary 53.** *Suppose that  $H \subset G$  is a closed diagonalisable subgroup. Then  $N_G(H)^0 = \mathcal{Z}_G(H)^0$  and  $N_G(H)/\mathcal{Z}_G(H)$  is finite. ( $N_G(H), \mathcal{Z}_G(H)$  are easily seen to be closed subgroups.)*

*Proof.* Applying the above proposition to the morphism

$$H \times N_G(H)^0 \rightarrow H, \quad (h, n) \mapsto nhn^{-1}$$

we get that  $nhn^{-1} = h$  for all  $h, n$ . Hence

$$N_G(H)^0 \subset \mathcal{Z}_G(H) \subset N_G(H)$$

and the corollary immediately follows.  $\square$

## 2.3 Elementary unipotent groups.

Define  $\mathcal{A}(G) := \text{Hom}(G, \mathbf{G}_a)$ , which is an abelian group under addition of maps; actually, it is an  $R$ -module, where  $R = \text{End}(\mathbf{G}_a)$ . Note that  $\mathcal{A}(\mathbf{G}_a) \cong R^n$ .  $R = \text{End}(\mathbf{G}_a)$  can be identified with

$$\{f \in k[\mathbf{G}_a] = k[x] \mid f(x+y) = f(x) + f(y) \text{ in } k[x, y]\} = \begin{cases} \{\lambda x \mid \lambda \in k\}, & \text{char } k = p = 0 \\ \{\sum \lambda_i x^{p^i} \mid \lambda_i \in k\}, & \text{char } k = p > 0 \end{cases}$$

Accordingly,

$$R \cong \begin{cases} k, & p = 0 \\ \text{noncommutative polynomial ring over } k, & p > 0 \end{cases}$$

**Proposition 54.**  *$G$  is an algebraic group. The following are equivalent:*

- (i)  $G$  is isomorphic to a closed subgroup of  $\mathbf{G}_a^n$  ( $n \geq 0$ ).
- (ii)  $\mathcal{A}(G)$  is a finitely-generated  $R$ -module and generates  $k[G]$  as a  $k$ -algebra.
- (iii)  $G$  is commutative and  $G = G_u$  (and  $G^p = 1$  if  $p > 0$ ).

**Definition 55.** *If one of the above conditions holds, then  $G$  is **elementary unipotent**. Note that (iii) rules out  $\mathbf{Z}/p^n\mathbf{Z}$  as elementary unipotent when  $n > 1$ .*

**Theorem 56.**

$$(\text{ elementary unipotent groups } ) \xrightarrow{\mathcal{A}} (\text{ finitely-generated } R\text{-modules } )$$

*is an equivalence of categories.*

*Proof.* For the inverse functor, see Springer 14.3.6. □

**Corollary 57.**

- (i) *The elementary unipotent groups are  $\mathbf{G}_a^n$  if  $p = 0$ , and  $\mathbf{G}_a^n \times (\mathbf{Z}/p\mathbf{Z})^s$  if  $p > 0$*
- (ii) *For an elementary unipotent group  $G$ ,*

$$G \text{ is isomorphic to a } \mathbf{G}_a^n \iff G \text{ is connected } \iff \mathcal{A}(G) \text{ is free}$$

**Theorem 58.** *Suppose  $G$  is a connected algebraic group of dimension 1, then  $G \cong \mathbf{G}_a$  or  $\mathbf{G}_m$ .*

*Proof.*

**Claim :**  $G$  is commutative.

Fix  $\gamma \in G$  and consider  $\phi : G \rightarrow G$  given by  $g \mapsto g\gamma g^{-1}$ . Then  $\overline{\phi(G)}$  is irreducible and closed, which implies that  $\overline{\phi(G)} = \{\gamma\}$  or  $\overline{\phi(G)} = G$ . Now, either  $\overline{\phi(G)} = \{\gamma\}$  for all  $\gamma \in G$ , in which case  $G$  is commutative and the claim is true, or  $\overline{\phi(G)} = G$  for at least one  $\gamma$ . Suppose the second case holds with a particular  $\gamma$  and fix an embedding  $G \hookrightarrow \text{GL}_n$ . Consider the morphism  $\psi : G \rightarrow \mathbf{A}^{n+1}$  which takes  $g$  to the coefficients of the characteristic polynomial of  $g$ ,  $\det(T \cdot \text{id} - g)$ .  $\psi$  is constant

on the conjugacy class  $\phi(G)$ , implying that  $\psi$  is constant. Hence, every  $g \in G$ ,  $e$  included, has the same characteristic polynomial:  $(T - 1)^n$ . Thus

$$G = G_u \implies G \text{ is nilpotent} \implies G \supsetneq [G, G] \implies [G, G] = 1 \implies G \text{ is commutative}$$

Now, by Proposition 37,

$$G \cong G_s \times G_u \implies G = G_s \text{ or } G = G_u$$

as dimension is additive. In the former case,  $G \cong \mathbf{G}_m$  by Corollary 46. In the latter, if we can prove that  $G$  is elementary unipotent, then  $G \cong \mathbf{G}_a$  by Corollary 57; we must show that  $G^p = 1$  when  $p > 0$  by Proposition 54. Suppose that  $G^p \neq 1$ , so that  $G^p = G$ . Then  $G = G^p = G^{p^2} = \dots$ . But  $(g - 1)^n = 0$  in  $\text{GL}_n$  and so for  $p^r \geq n$ ,

$$0 = (g - 1)^{p^r} = g^{p^r} - 1 \implies g^{p^r} = 1 \implies \{e\} = G^{p^r} = G$$

which is a contradiction. □

### 3. Lie algebras.

If  $X$  is a variety and  $x \in X$ , then the **local ring** at  $x$  is

$$\mathcal{O}_{X,x} := \varinjlim_{\substack{U \text{ open} \\ U \ni x}} \mathcal{O}_X(U) = \text{germs of functions at } x = \frac{\{(f, U) \mid f \in \mathcal{O}_X(U)\}}{\sim}$$

where  $(f, U) \sim (f', U')$  if there is an open neighbourhood  $V \subset U \cap U'$  of  $x$  for which  $f|_V = f'|_V$ . There is a well-defined ring morphism  $\text{ev}_x : \mathcal{O}_{X,x} \rightarrow k$  given by evaluating at  $x$ :  $[(f, U)] \mapsto f(x)$ .  $\mathcal{O}_{X,x}$  is a local ring (hence the name) with unique maximal ideal

$$\mathfrak{m}_x =: \ker \text{ev}_x = \{[(f, U)] \mid f(x) = 0\}$$

for if  $f \notin \mathfrak{m}_x$ , then  $f^{-1}$  is defined near  $x$ , implying that  $f \in \mathcal{O}_{X,x}^\times$ .

*Fact.* If  $X$  is affine and  $x$  corresponds to the maximal ideal  $\mathfrak{m} \subset k[X]$  (via Nullstellensatz), then  $\mathcal{O}_{X,x} \cong k[X]_{\mathfrak{m}}$ . By choosing an affine chart in  $X$  at  $x$ , we see in general that  $\mathcal{O}_{X,x}$  is noetherian.

#### 3.1 Tangent Spaces.

Analogous to the case of manifolds, the **tangent space** to a variety  $X$  at a point  $x$  is

$$T_x X := \text{Der}_k(\mathcal{O}_{X,x}, k) = \{\delta : \mathcal{O}_{X,x} \rightarrow k \mid \delta \text{ is } k\text{-linear, } \delta(fg) = f(x)\delta(g) + g(x)\delta(f)\}$$

(so  $k$  is viewed as a  $\mathcal{O}_{X,x}$ -module via  $\text{ev}_x$ .)  $T_x X$  is a  $k$ -vector space.

**Lemma 59.** *Let  $A$  be a  $k$ -algebra,  $\epsilon : A \rightarrow k$  a  $k$ -algebra morphism, and  $\mathfrak{m} = \ker \epsilon$ . Then*

$$\text{Der}_k(A, k) \xrightarrow{\sim} (\mathfrak{m}/\mathfrak{m}^2)^*, \quad \delta \mapsto \delta|_{\mathfrak{m}}$$

*Proof.* An inverse map is given by sending  $\lambda$  to a derivation defined by  $x \mapsto \begin{cases} 0, & x = 1 \\ \lambda(x), & x \in \mathfrak{m} \end{cases}$ .

Checking this is an exercise. □

Hence,  $T_x X \cong (\mathfrak{m}_x/\mathfrak{m}_x^2)^*$  is finite-dimensional.

*Examples.*

- If  $X = \mathbf{A}^n$ , then  $T_x X$  has basis

$$\left. \frac{\partial}{\partial x_1} \right|_x, \dots, \left. \frac{\partial}{\partial x_n} \right|_x$$

- For a finite-dimensional  $k$ -vector space  $V$ ,  $T_x(V) \cong V$ .

**Definition 60.**  $X$  is **smooth** at  $x$  if  $\dim T_x X = \dim X$ . Moreover,  $X$  is **smooth** if it is smooth at every point. From the above example, we see that  $\mathbf{A}^n$  is smooth.

If  $\phi : X \rightarrow Y$  we get  $\phi^* : \mathcal{O}_{Y, \phi(x)} \rightarrow \mathcal{O}_{X, x}$  and hence

$$d\phi : T_x X \rightarrow T_{\phi(x)} Y, \quad \delta \mapsto \delta \circ \phi^*$$

**Remark 61.** If  $U \subset X$  is an open neighbourhood of  $x$ , then  $d(U \hookrightarrow X) : T_x U \xrightarrow{\sim} T_x X$ . More generally, if  $X \subset Y$  is a locally closed subvariety, then  $T_x X$  embeds into  $T_x Y$ .

**Theorem 62.**

$$\dim T_x X \geq \dim X$$

with equality holding for all  $x$  in some open dense subset.

Note that if  $X$  is affine and  $x$  corresponds to  $\mathfrak{m} \subset k[X]$ , then the natural map  $k[X] \rightarrow k[X]_{\mathfrak{m}} = \mathcal{O}_{X, x}$  induces an isomorphism

$$T_x X \xrightarrow{\sim} \text{Der}_k(k[X], k), \quad (k \text{ being viewed as a } k[X]\text{-modules via } \text{ev}_x)$$

which is isomorphic to  $(\mathfrak{m}/\mathfrak{m}^2)^*$  by Lemma 59. So, we can work without localising.  $\square$

**Remark 63.** If  $G$  is an algebraic group, then  $G$  is smooth by Theorem 62 since

$$d(\ell_g : x \mapsto gx) : T_x G \xrightarrow{\sim} T_{g\gamma} G$$

The same holds for **homogeneous**  $G$ -spaces (i.e.,  $G$ -spaces for which the  $G$ -action is transitive).

## 3.2 Lie algebras.

**Definition 64.** A **Lie algebra** is a  $k$ -vector space  $L$  together with a bilinear map  $[\cdot, \cdot] : L \times L \rightarrow L$  such that

- (i)  $[x, x] = 0 \quad \forall x \in L \quad ( \implies [x, y] = -[y, x] )$
- (ii)  $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0 \quad \forall x, y, z \in L$

*Examples.*

- If  $A$  is an associative  $k$ -algebra (maybe non-unital), then  $[a, b] := ab - ba$  gives  $A$  the structure of a Lie algebra.
- Take  $A = \text{End}(V)$  and as above define  $[\alpha, \beta] = \alpha \circ \beta - \beta \circ \alpha$ .
- For  $L$  an arbitrary  $k$ -vector space, define  $[\cdot, \cdot] = 0$ . When  $[\cdot, \cdot] = 0$  a Lie algebra is said to be **abelian**.

We will construct a functor

$$( \text{ algebraic groups } ) \xrightarrow{\text{Lie}} ( \text{ Lie algebras } )$$

As a vector space,  $\text{Lie } G = T_e G$ .  $\dim \text{Lie } G = \dim G$  by above remarks.

The following is another way to think about  $T_e G$ . Recall that we can identify  $G$  with the functor

$$R \mapsto \text{Hom}_{\text{alg}}(k[G], R) := G(R)$$

(where  $k[G]$  is a reduced finite-dimensional commutative Hopf  $k$ -algebra). The Hopf (i.e., cogroup) structure on  $R$  induces a group structure on  $G(R)$ , even when  $R$  is not reduced..

**Lemma 65.**

$$\text{Lie } G \cong \ker \left( G(k[\epsilon]/(\epsilon^2)) \rightarrow G(k) \right)$$

as abelian groups.

*Proof.* Write the algebra morphism  $\theta : k[G] \rightarrow k[\epsilon]/(\epsilon^2)$  as given by  $f \mapsto \text{ev}_e(f) + \delta(f) \cdot \epsilon$  for some  $\delta : k[G] \rightarrow k$ .  $\delta$  is a derivation.  $\square$

*Examples.*

- For  $G = \text{GL}_n$ ,  $G(R) = \text{GL}_n(R)$ , and we have

$$\text{Lie } G = \ker \left( \text{GL}_n(k[\epsilon]/(\epsilon^2)) \rightarrow \text{GL}_n(k) \right) = \{I + A\epsilon \mid A \in M_n(k)\} \xrightarrow{\sim} M_n(k)$$

Explicitly, the isomorphism  $\text{Lie } \text{GL}_n \rightarrow M_n(k)$  is given by  $\delta \mapsto (\partial(T_{ij}))$ .

- Intrinsically, for a finite-dimensional  $k$ -vector space  $V$ : Since  $\text{GL}(V)$  is an open subset of  $\text{End}(V)$ , we have

$$\text{Lie } \text{GL}(V) \xrightarrow{\sim} T_I(\text{End } V) \xrightarrow{\sim} \text{End } V$$

**Definition 66.** A left-invariant vector field on  $G$  is an element  $D \in \text{Der}_k(k[G], k[G])$  such that the

$$\begin{array}{ccc} k[G] & \xrightarrow{D} & k[G] \\ \Delta \downarrow & & \downarrow \Delta \\ k[G] \otimes k[G] & \xrightarrow{\text{id} \otimes D} & k[G] \otimes k[G] \end{array}$$

commutes.

For a fixed  $D$ , for  $g \in G$ , define  $\delta_g := \text{ev}_g \circ D \in T_g G$ .

Evaluating  $\Delta \circ D$  at  $(g_1, g_2)$  gives  $\delta_{g_1 g_2}$

Evaluating  $(\text{id} \otimes D) \circ \Delta$  at  $(g_1, g_2)$  gives  $\delta_{g_2} \circ \ell_{g_1}^* = d\ell_{g_1}(\delta_{g_2})$

Hence  $D \in$  being left-invariant is equivalent to  $\delta_{g_1 g_2} = d\ell_{g_1}(\delta_{g_2})$  for all  $g_1, g_2 \in G$ . Define

$$\mathcal{D}_G := \text{vector space of left-invariant vector fields on } G$$

**Theorem 67.**

$$\mathcal{D}_G \rightarrow \text{Lie } G, \quad D \mapsto \delta_e = \text{ev}_e \circ D$$

is a linear isomorphism.

*Proof.* We shall prove that  $\delta \mapsto (\text{id} \otimes \delta) \circ \Delta$  is an inverse morphism. Fix  $\delta \in \text{Lie } G$ , set  $D = (\text{id}, \delta) \circ \Delta : k[G] \rightarrow k[G]$ , and check that  $(\text{id}, \delta)$  is a  $k$ -derivation  $k[G] \otimes k[G] \rightarrow k[G]$ , where  $k[G]$  is viewed as a  $k[G] \otimes k[G]$ -module via  $\text{id} \otimes \text{ev}_e$ . First, we shall check that  $D \in \mathcal{D}_G$ :

$$\begin{aligned}
D(fh) &= (\text{id} \otimes \delta)(\Delta(fh)) \\
&= (\text{id} \otimes \delta)(\Delta(f) \cdot \Delta(h)) \\
&= (\text{id} \otimes \text{ev}_e)(\Delta f) \cdot (\text{id} \otimes \delta)(\Delta h) + (\text{id} \otimes \text{ev}_e)(\Delta h) \cdot (\text{id} \otimes \delta)(\Delta f) \\
&= f \cdot D(h) + h \cdot D(f)
\end{aligned}$$

Next, we show that  $D$  is left-invariant:

$$\begin{aligned}
(\text{id} \otimes D) \circ \Delta &= (\text{id} \otimes ((\text{id} \otimes \delta) \circ \Delta)) \circ \Delta \\
&= (\text{id} \otimes (\text{id} \otimes \delta)) \circ (\text{id} \circ \Delta) \circ \Delta \\
&= (\text{id} \otimes (\text{id} \otimes \delta)) \circ (\Delta \circ \text{id}) \circ \Delta \quad (\text{“ co-associativity”}) \\
&= \Delta \circ (\text{id} \otimes \delta) \circ \Delta \quad (\text{easily checked}) \\
&= \Delta \circ D
\end{aligned}$$

Lastly, we show that the maps are inverse:

$$\begin{aligned}
\delta \mapsto (\text{id} \otimes \delta) \circ \Delta &\mapsto \text{ev}_e \circ (\text{id} \otimes \delta) \circ \Delta = \delta \circ (\text{ev}_e \otimes \text{id}) \circ \Delta = \delta \\
D \mapsto \text{ev}_e \circ D &\mapsto (\text{id} \otimes \text{ev}_e) \circ (\text{id} \otimes D) \circ \Delta = (\text{id} \otimes \text{ev}_e) \circ \Delta \circ D = D
\end{aligned}$$

□

Since  $\text{Hom}_k(k[G], k[G])$  is an associative algebra, there is a natural candidate for a Lie bracket on  $\mathcal{D}_G \subset \text{Hom}_k(k[G], k[G])$ :  $[D_1, D_2] = D_1 \circ D_2 - D_2 \circ D_1$ . We must check that  $[\mathcal{D}_G, \mathcal{D}_G] \subset \mathcal{D}_G$ . Let  $D_1, D_2 \in \mathcal{D}_G$ . Since

$$\begin{aligned}
[D_1, D_2](fh) &= D_1(D_2(fh)) - D_2(D_1(fh)) \\
&= D_1(f \cdot D_2(h) + h \cdot D_2(f)) - D_2(f \cdot D_1(h) + h \cdot D_1(f)) \\
&= D_1(f \cdot D_2(h)) + D_1(h \cdot D_2(f)) - D_2(f \cdot D_1(h)) - D_2(h \cdot D_1(f)) \\
&= \left( f D_1(D_2(h)) + D_2(h) D_1(f) \right) + \left( h D_1(D_2(f)) + D_2(f) D_1(h) \right) \\
&\quad - \left( f D_2(D_1(h)) + D_1(h) D_2(f) \right) - \left( h D_2(D_1(f)) + D_1(f) D_2(h) \right) \\
&= f \left( D_1(D_2(h)) - f D_2(D_1(h)) \right) + h \left( D_1(D_2(f)) - h D_2(D_1(f)) \right) \\
&= f \cdot [D_1, D_2](h) + h \cdot [D_1, D_2](f)
\end{aligned}$$

we have that  $[D_1, D_2]$  is a derivation. Moreover,

$$\begin{aligned}
(\text{id} \otimes [D_1, D_2]) \circ \Delta &= (\text{id} \otimes (D_1 \circ D_2)) \circ \Delta - (\text{id} \otimes (D_2 \circ D_1)) \circ \Delta \\
&= (\text{id} \otimes D_1) \circ (\text{id} \otimes D_2) \circ \Delta - (\text{id} \otimes D_2) \circ (\text{id} \otimes D_1) \circ \Delta \\
&= (\text{id} \otimes D_1) \circ \Delta \circ D_2 - (\text{id} \otimes D_2) \circ \Delta \circ D_1 \\
&= \Delta \circ D_1 \circ D_2 - \Delta \circ D_2 \circ D_1 \\
&= \Delta \circ [D_1, D_2]
\end{aligned}$$

and so  $[D_1, D_2]$  is left-invariant. Accordingly,  $[\mathcal{D}_G, \mathcal{D}_G] \subset \mathcal{D}_G$ , and thus by the above theorem  $\text{Lie } G$  becomes a Lie algebra.

**Remark 68.** If  $p > 0$ , then  $\mathcal{D}_G$  is also stable under  $D \mapsto D^p$  (composition with itself  $p$ -times).

**Proposition 69.** If  $\delta_1, \delta_2 \in \text{Lie } G$ , then  $[\delta_1, \delta_2] : k[G] \rightarrow k$  is given by

$$[\delta_1, \delta_2] = ((\delta_1, \delta_2) - (\delta_2, \delta_1)) \circ \Delta$$

*Proof.* Let  $D_i = (\text{id} \otimes \delta_i) \circ \Delta$  for  $i = 1, 2$ . Then

$$\begin{aligned} [\delta_1, \delta_2] &= \text{ev}_e \circ [D_1, D_2] \\ &= \text{ev}_e \circ D_1 \circ D_2 - \text{ev}_e \circ D_2 \circ D_1 \\ &= \delta_1 \circ (\text{id} \otimes \delta_2) \circ \Delta - \delta_2 \circ (\text{id} \otimes \delta_1) \circ \Delta \\ &= (\delta_1 \otimes \delta_2) \circ \Delta - (\delta_2 \otimes \delta_1) \circ \Delta \\ &= ((\delta_1 \otimes \delta_2) - (\delta_2 \otimes \delta_1)) \circ D \end{aligned}$$

□

**Corollary 70.** If  $\phi : G \rightarrow H$  is a morphism of algebraic groups, then  $d\phi : \text{Lie } G \rightarrow \text{Lie } H$  is a morphism of Lie algebras (i.e., brackets are preserved).

*Proof.*

$$\begin{aligned} d\phi([\delta_1, \delta_2]) &= [\delta_1, \delta_2] \circ \phi^* \\ &= (\delta_1 \otimes \delta_2 - \delta_2 \otimes \delta_1) \circ \Delta \circ \phi^*, \quad (\text{by the above Prop.}) \\ &= ((\delta_1 \otimes \phi^*) \otimes (\delta_2 \otimes \phi^*) - (\delta_2 \otimes \phi^*) \otimes (\delta_1 \otimes \phi^*)) \circ \Delta \\ &= (\delta_1 \circ \phi^*, \delta_2 \circ \phi^*) \circ \Delta - (\delta_2 \circ \phi^*, \delta_1 \circ \phi^*) \circ \Delta \\ &= (d\phi(\delta_1), d\phi(\delta_2)) \circ \Delta - (d\phi(\delta_2), d\phi(\delta_1)) \circ \Delta \\ &= [d\phi(\delta_1), d\phi(\delta_2)] \end{aligned}$$

□

**Corollary 71.** If  $G$  is commutative, then so too is  $\text{Lie } G$  (i.e.,  $[\cdot, \cdot] = 0$ ).

*Example.* We have that  $\phi : \text{Lie } \text{GL}_n \cong M_n(k)$  is given by  $\phi : \delta \mapsto (\delta(T_{ij}))$ . Since

$$\begin{aligned} [\delta_1, \delta_2](T_{ij}) &= (\delta_1, \delta_2)(\Delta T_{ij}) - (\delta_2, \delta_1)(\Delta T_{ij}) \\ &= \sum_{l=1}^n \delta_1(T_{il})\delta_2(T_{lj}) - \sum_{l=1}^n \delta_2(T_{il})\delta_1(T_{lj}) \\ &= (\phi(\delta_1)\phi(\delta_2))_{ij} - (\phi(\delta_2)\phi(\delta_1))_{ij} \end{aligned}$$

Hence,

$$\phi([\delta_1, \delta_2]) = \phi(\delta_1)\phi(\delta_2) - \phi(\delta_2)\phi(\delta_1)$$

and so in identifying  $\text{Lie } \text{GL}_n$  with  $M_n(k)$ , we can also identify the Lie bracket with the usual one on  $M_n(k)$ :  $[A, B] = AB - BA$ . Similarly, the Lie bracket on  $\text{Lie } \text{GL}(V) \cong \text{End}(V)$  can be identified with the commutator. □

**Remark 72.** If  $\phi : G \rightarrow H$  is a closed immersion, then  $\phi^*$  is surjective, and so  $d\phi : \text{Lie } G \rightarrow \text{Lie } H$  is injective. Hence, if  $G \hookrightarrow \text{GL}_n$ , then the above example determines  $[\cdot, \cdot]$  on  $\text{Lie } G$ .

*Examples.*

- $\text{Lie } \text{SL}_n =$  trace 0 matrices in  $M_n(k)$
- $\text{Lie } B_n =$  upper-triangular matrices in  $M_n(k)$
- $\text{Lie } U_n =$  strictly upper-triangular matrices in  $M_n(k)$
- $\text{Lie } D_n =$  triangular matrices in  $M_n(k)$

**Exercise.** If  $G$  is diagonal, show that  $\text{Lie } G \cong \text{Hom}_{\mathbf{Z}}(X^*(G), k)$ .

### 3.3 Adjoint representation.

$G$  acts on itself by conjugation: for  $x \in G$ ,

$$c_x : G \rightarrow G, \quad g \mapsto xgx^{-1}$$

is a morphism.  $\text{Ad}(x) := dc_x : \text{Lie } G \rightarrow \text{Lie } G$  is a Lie algebra endomorphism such that

$$\text{Ad}(e) = \text{id}, \quad \text{Ad}(xy) = \text{Ad}(x) \circ \text{Ad}(y)$$

Hence, we have a morphism of groups

$$\text{Ad} : G \rightarrow \text{GL}(\text{Lie } G)$$

**Proposition 73.** *Ad is an algebraic representation of  $G$ .*

*Proof.* We must show that

$$\theta : G \times \text{Lie } G \rightarrow \text{Lie } G, \quad (x, \delta) \mapsto \text{Ad}(x)(\delta) = dc_x(\delta) = \delta \circ c_x^*$$

is a morphism of varieties. It is enough to show that  $\lambda \circ \theta$  is a morphism for all  $\lambda \in (\text{Lie } G)^*$ . Given such a  $\lambda$ , since  $(\text{Lie } G)^* \cong \mathfrak{m}/\mathfrak{m}^2$  we must have  $\lambda(\delta) = \delta(f)$  for some  $f \in \mathfrak{m}$ . Accordingly, for any  $f \in \mathfrak{m}$  we must show that

$$(x, \delta) \mapsto \delta(c_x^* f)$$

is a morphism. Recall from the proof of Proposition 27 that  $c_x^* f = \sum_i h_i(x) f_i$  for some  $f_i, h_i \in k[G]$ , which implies that

$$(x, \delta) \mapsto \delta(c_x^* f) = \sum_i h_i(x) \delta(f_i)$$

is a morphism as  $x \mapsto h_i(x)$  and  $\delta \mapsto \delta(f_i)$  are morphisms. □.

**Exercises.**

- Show that  $\text{ad} := d(\text{Ad}) : \text{Lie } G \rightarrow \text{End}(\text{Lie } G)$  is

$$\delta_1 \mapsto (\delta_2 \mapsto [\delta_1, \delta_2])$$

This is hard, but is easiest to manage in reducing to the case of  $\text{GL}_n$  using an embedding  $G \hookrightarrow \text{GL}_n$ .

- Show that  $d(\det : \text{GL}_n \rightarrow \text{GL}_1) : M_n(k) \rightarrow k$  is the trace map.

### 3.4 Some derivatives.

If  $X_1, X_2$  are varieties with points  $x_1 \in X_1$  and  $x_2 \in X_2$ , then the morphisms

$$\begin{array}{ccccc}
 & & X_1 & & \\
 & \nearrow \pi_1 & & \searrow i_{x_1}: x \mapsto (x_1, x) & \\
 X_1 \times X_2 & & & & X_1 \times X_2 \\
 & \searrow \pi_2 & & \nearrow i_{x_2}: x \mapsto (x, x_2) & \\
 & & X_2 & & 
 \end{array}$$

induce inverse isomorphisms  $T_{x_1}X_1 \oplus T_{x_2}X_2 \xrightarrow{\sim} T_{(x_1, x_2)}(X_1 \times X_2)$ . In particular, for algebraic groups  $G_1, G_2$  we have inverse isomorphisms

$$\mathrm{Lie} G_1 \oplus \mathrm{Lie} G_2 \xrightarrow{\sim} \mathrm{Lie} (G_1 \times G_2)$$

**Proposition 74.**

- (i)  $d(\mu : G \times G \rightarrow G) = (\mathrm{Lie} G \oplus \mathrm{Lie} G \xrightarrow{(X, Y) \mapsto X+Y} \mathrm{Lie} G)$
- (ii)  $d(i : G \rightarrow G) = (\mathrm{Lie} G \xrightarrow{X \mapsto -X} \mathrm{Lie} G)$

*Proof.*

(i). It is enough to show that  $d\mu$  is the identity on each factor. Since  $\mathrm{id}_G$  can be factored as

$$G \xrightarrow{i_e} G \times G \xrightarrow{\mu} G$$

where  $i_e : x \mapsto (e, x)$  or  $x \mapsto (x, e)$ , we are done.

(ii). Since  $x \mapsto e$  can be factored  $G \xrightarrow{(\mathrm{id}, i)} G \times G \xrightarrow{\mu} G$ . From (i) we have that  $0 : \mathrm{Lie} G \rightarrow \mathrm{Lie} G$  can be factored as

$$\mathrm{Lie} G \xrightarrow{(\mathrm{id}, di)} \mathrm{Lie} G \otimes \mathrm{Lie} G \xrightarrow{\pm} \mathrm{Lie} G$$

□

**Remark 75.** The open immersion  $G^0 \hookrightarrow G$  induces an isomorphism  $\mathrm{Lie} G^0 \xrightarrow{\sim} \mathrm{Lie} G$ .

**Proposition 76** (Derivative of a linear map). *If  $V, W$  be vector spaces and  $f : V \rightarrow W$  a linear map (hence a morphism), then, for all  $v \in V$ , we have the commutative diagram*

$$\begin{array}{ccc}
 T_v V & \xrightarrow{T_v(f)} & T_{f(v)} W \\
 \wr \downarrow & & \downarrow \wr \\
 V & \xrightarrow{f} & W
 \end{array}$$

*Proof.* Exercise. □

**Proposition 77.** *Suppose that  $\sigma : G \rightarrow \mathrm{GL}(V)$  is a representation and  $v \in V$ . Define  $o_v : G \rightarrow V$  by  $g \mapsto \sigma(g)v$ . Then*

$$do_v(X) = d\sigma(X)(v)$$

in  $T_vV \cong V$ .

*Proof.* Factor  $o_v$  as

$$\begin{array}{ccc} G & \xrightarrow{\phi} & \mathrm{GL}(V) \times V & \xrightarrow{\psi} & V \\ g & \mapsto & (\sigma(g), v) & & \\ & & (A, w) & \mapsto & Aw \end{array}$$

$d\phi = (d\sigma, 0) : \mathrm{Lie} G \rightarrow \mathrm{End} V \oplus V$ . By 76, under the identification  $V \cong T_vV$ , we have that the derivative at  $(e, v)$  of the first component of  $\psi$ , which sends  $A \rightarrow Av$ , is the same map. The result follows.  $\square$

**Proposition 78.** *Suppose that  $\rho_i : G \rightarrow \mathrm{GL}(V_i)$  are representations for  $i = 1, 2$ . Then the derivative of  $\rho_1 \otimes \rho_2 : G \rightarrow \mathrm{GL}(V_1 \otimes V_2)$  is*

$$d(\rho_1 \otimes \rho_2)X = d\rho_1(X) \otimes \mathrm{id} + \mathrm{id} \otimes d\rho_2(X)$$

(i.e.,  $X(v_1 \otimes v_2) = (Xv_1) \otimes v_2 + v_1 \otimes (Xv_2)$ .) Similarly for  $V_1 \otimes \cdots \otimes V_n$ ,  $\mathrm{Sym}^n V$ ,  $\Lambda^n V$ .

*Proof.* We have the commutative diagram

$$\begin{array}{ccccc} \rho_1 \otimes \rho_2 : G & \longrightarrow & \mathrm{GL}(V_1) \times \mathrm{GL}(V_2) & \longrightarrow & \mathrm{GL}(V_1 \otimes V_2) \\ & & \downarrow \text{open} & & \downarrow \text{open} \\ & & \mathrm{End}(V_1) \times \mathrm{End}(V_2) & \xrightarrow{\phi} & \mathrm{End}(V_1 \otimes V_2) \end{array}$$

where  $\phi : (A, B) \mapsto A \otimes B$ . (Note that  $\phi$  being a morphism implies that  $\rho_1 \otimes \rho_2$ .) Computing  $d\phi$  component-wise at  $(1, 1)$ , we get that  $d\phi|_{\mathrm{End}(V_1)}$  is the derivative of the linear map  $\mathrm{End}(V_1) \rightarrow \mathrm{End}(V_1 \otimes V_2)$  given by  $A \mapsto A \otimes 1$ , which is the same map; likewise for  $d\phi|_{\mathrm{End}(V_2)}$ . Hence,

$$d\phi(A, B) = A \otimes 1 + 1 \otimes B$$

and we are done.  $\square$

**Proposition 79** (Adjoint representation for  $\mathrm{GL}(V)$ ). *For  $g \in \mathrm{GL}(V)$ ,  $A \in \mathrm{Lie} \mathrm{GL}(V) \cong \mathrm{End}(V)$ ,*

$$\mathrm{Ad}(g)A = gAg^{-1}$$

*Proof.* This follows from Proposition 76 with  $V = \mathrm{GL}(V) \hookrightarrow \mathrm{End}(V) = W$  and  $f = c_g : A \mapsto gAg^{-1}$ .  $\square$

**Exercise.** Deduce that, for  $\mathrm{GL}(V)$ ,  $\mathrm{ad}(A)(B) = AB - BA$ .

### 3.5 Separable morphisms.

Let  $\phi : X \rightarrow Y$  be a *dominant* morphism of varieties (i.e.,  $\overline{\phi(X)} = Y$ ). From the induced maps  $\mathcal{O}_Y(V) \rightarrow \mathcal{O}_X(\phi^{-1}(V))$  - note that  $\phi^{-1}(V) \neq \emptyset$ , as  $\phi$  is dominant - given by  $f \mapsto f \circ \phi$ , we get a morphism of fields  $\phi^* : k(Y) \rightarrow k(X)$ . That is,  $k(X)$  is a finitely-generated field extension of  $k(Y)$ .

**Remark 80.** *This field extension has transcendence degree  $\dim X - \dim Y$ , and hence is algebraic if and only if  $\dim X = \dim Y$ .*

**Definition 81.** *A dominant  $\phi$  is **separable** if  $\phi^* : k(Y) \rightarrow k(X)$  is a separable field extension.*

**Recall.**

- An algebraic field extension  $E/F$  being separable means that every  $\alpha \in E$  has a minimal polynomial without repeated roots.
- A finitely-generated field extension  $E/F$  is separable if it is of the form

$$\begin{array}{c} E \\ \left| \begin{array}{l} \text{finite separable} \end{array} \right. \\ F(x_1, \dots, x_n) \\ \left| \begin{array}{l} x_1, \dots, x_n \text{ algebraically independent} \end{array} \right. \\ F \end{array}$$

*Facts.*

- If  $E'/E$  and  $E/F$  are separable then  $E'/F$  is separable.
- If  $\text{char } k = 0$ , all extensions are separable; in characteristic 0 being dominant is equivalent to being separable. (As an example, if  $\text{char } k = p > 0$ , then  $F(t^{1/p})/F(t)$  is never separable.)
- The composition of separable morphisms is separable.

*Example.* If  $p > 0$ , then  $\mathbf{G}_m \xrightarrow{p} \mathbf{G}_m$  is not separable.

**Theorem 82.** *Let  $\phi : X \rightarrow Y$  be a morphism between irreducible varieties. The following are equivalent:*

- (i)  $\phi$  is separable.
- (ii) There is a dense open set  $U \subset X$  such that  $d\phi_x : T_x X \rightarrow T_{\phi(x)} Y$  is surjective for all  $x \in U$ .
- (iii) There is an  $x \in X$  such that  $X$  is smooth at  $x$ ,  $Y$  is smooth at  $\phi(x)$ , and  $d\phi_x$  is surjective.

**Corollary 83.** *If  $X, Y$  are irreducible, smooth varieties, then  $\phi : X \rightarrow Y$*

$$\text{is separable} \iff d\phi_x \text{ is surjective for all } x \iff d\phi_x \text{ is surjective for one } x$$

**Remark 84.** *The corollary applies in particular if  $X, Y$  are algebraic groups or homogeneous spaces.*

### 3.6 Fibres of morphisms.

**Theorem 85.** Let  $\phi : X \rightarrow Y$  be a dominant morphism between irreducible varieties and let  $r := \dim X - \dim Y \geq 0$ .

- (i) For all  $y \in \phi(X)$ ,  $\dim \phi^{-1}(y) \geq r$
- (ii) There is a nonempty open subset  $V \subset Y$  such that for all irreducible closed  $Z \subset Y$  and for all irreducible components  $Z' \subset \phi^{-1}(Z)$  with  $Z' \cap \phi^{-1}(V) \neq \emptyset$ ,  $\dim Z' = \dim Z + r$  (which implies that  $\dim \phi^{-1}(y) = r$  for all  $y \in V$ .) If  $r = 0$ ,  $|\phi^{-1}(y)| = [k(X), k(Y)]_s$  for all  $y \in V$ .

**Theorem 86.** If  $\phi : X \rightarrow Y$  is a dominant morphism between irreducible varieties, then there is a nonempty open  $V \subset Y$  such that  $\phi^{-1}(V) \xrightarrow{\phi} V$  is universally open, i.e., for all varieties  $Z$

$$\phi^{-1}(V) \times Z \xrightarrow{\phi \times \text{id}_Z} V \times Z$$

is an open map.

**Corollary 87.** If  $\phi : X \rightarrow Y$  is a  $G$ -equivariant morphism of homogeneous spaces,

- (i) For all varieties  $Z$ ,  $\phi \times \text{id}_Z : X \times Z \rightarrow Y \times Z$  is an open map.
- (ii) For all closed, irreducible  $Z \subset Y$  and for all irreducible components  $Z' \subset \phi^{-1}(Z)$ ,  $\dim Z' = \dim Z + r$ . (In particular, all fibres are equidimensional of dimension  $r$ .)
- (iii)  $\phi$  is an isomorphism if and only if  $\phi$  is bijective and  $d\phi_x$  is an isomorphism for one (or, equivalently, all)  $x$ .

**Corollary 88.** For all  $G$ -spaces,  $\dim \text{Stab}_G(x) + \dim(Gx) = \dim G$

*Proof.* Apply the above to  $G \rightarrow Gx$ . □

**Corollary 89.** Let  $\phi : G \rightarrow H$  be a surjective morphism of algebraic groups.

- (i)  $\phi$  is open
- (ii)  $\dim G = \dim H + \dim \ker \phi$
- (iii)

$$\phi \text{ is an isomorphism} \iff \phi \text{ and } d\phi \text{ are bijective} \iff \phi \text{ is bijective and separable}$$

*Proof.* They are homogeneous  $G$ -spaces by left-translation,  $H$  via  $\phi$ . □

**Definition 90.** A sequence of algebraic groups

$$1 \rightarrow K \xrightarrow{\phi} G \xrightarrow{\psi} H \rightarrow 1$$

is **exact** if

- (i) it is set-theoretically exact and
- (ii)

$$0 \rightarrow \text{Lie } K \xrightarrow{d\phi} \text{Lie } G \xrightarrow{d\psi} \text{Lie } H \rightarrow 0$$

is an exact sequence of lie algebras (i.e., of vector spaces).

**Exercise.** Show that condition (ii) above can be replaced above by (ii')  $\phi$  being a closed immersion and  $\psi$  being separable. In characteristic 0, show that (ii') is automatic.

**Theorem 91** (Weak form of Zariski's Main Theorem). *If  $\phi : X \rightarrow Y$  is a morphism between irreducible varieties such that  $Y$  is smooth, and  $\phi$  is birational (i.e.,  $k(Y) = k(X)$ ) and bijective, then  $\phi$  is an isomorphism.*

### 3.7 Semisimple automorphisms.

**Definition 92.** *An automorphism  $\sigma : G \rightarrow G$  is **semisimple** if there is a  $G \hookrightarrow \mathrm{GL}_n$  and a semisimple element  $s \in \mathrm{GL}_n$  such that  $\sigma(g) = sgs^{-1}$  for all  $g \in G$ .*

*Example.* If  $s \in G_s$ , then the inner automorphism  $g \mapsto sgs^{-1}$  is semisimple.

**Definitions 93.** *Given a semisimple automorphism of  $G$ , define*

$$G_\sigma := \{g \in G \mid \sigma(g) = g\}, \text{ which is a closed subgroup}$$

$$\mathfrak{g}_\sigma := \{X \in \mathfrak{g} := \mathrm{Lie} G \mid d\sigma(X) = X\}$$

*Let  $\tau : G \rightarrow G$ ,  $g \mapsto \sigma(g)g^{-1}$ . Then  $G_\sigma = \tau^{-1}(e)$  and  $d\tau = d\sigma - \mathrm{id}$  by Proposition 74, which implies that  $\ker d\tau = \mathfrak{g}_\sigma$ . Since  $G_\sigma \hookrightarrow G \xrightarrow{\tau} G$  is constant, we have*

$$d\tau(\mathrm{Lie} G_\sigma) = 0 \implies \mathrm{Lie} G_\sigma \subset \mathfrak{g}_\sigma$$

**Lemma 94.**

$$\mathrm{Lie} G_\sigma = \mathfrak{g}_\sigma \iff G \xrightarrow{\tau} \tau(G) \text{ is separable} \iff d\tau : \mathrm{Lie} G \rightarrow T_e(\tau(G)) \text{ is surjective}$$

*Proof.*  $\tau$  is a  $G$ -map of homogeneous spaces, acting by  $x * g = \sigma(x)gx^{-1}$  on the codomain.  $\tau(G)$  is smooth and is, by Proposition 24, locally closed. Hence, by Theorem 82

$$\begin{aligned} \tau \text{ is separable} &\iff d\tau \text{ is surjective} \\ &\iff \dim \mathfrak{g}_\sigma = \dim \ker d\tau = \dim G - \dim \tau(G) = \dim G_\sigma = \dim \mathrm{Lie} G_\sigma \\ &\iff \mathfrak{g}_\sigma = \mathrm{Lie} G_\sigma \end{aligned}$$

□

**Proposition 95.**  $\tau(G)$  is closed and  $\mathrm{Lie} G_\sigma = \mathfrak{g}_\sigma$ .

*Proof.* Without loss of generality  $G \subset \mathrm{GL}_n$  is a closed subgroup and  $\sigma(g) = sgs^{-1}$  for some semisimple  $s \in \mathrm{GL}_n$ . Without loss of generality,  $s$  is diagonal with

$$s = a_1 I_{m_1} \times \cdots \times a_n I_{m_n}$$

with the  $a_i$  distinct and  $n = m_1 + \cdots + m_n$ . Then, extending  $\tau, \sigma$  to  $\mathrm{GL}_n$ , we have

$$(\mathrm{GL}_n)_\sigma = \mathrm{GL}_{m_1} \times \cdots \times \mathrm{GL}_{m_n} \quad \text{and} \quad (\mathfrak{gl}_n)_\sigma = M_{m_1} \times \cdots \times M_{m_n}$$

So,  $\text{Lie}(\text{GL}_n)_\sigma = (\mathfrak{gl}_n)_\sigma$ . Hence

$$\begin{array}{ccc} \mathfrak{gl}_n & \xrightarrow{d\tau} & T_e(\tau(\text{GL}_n)) \\ \uparrow & & \uparrow \\ \mathfrak{g} & \xrightarrow{d\tau} & T_e(\tau(G)) \end{array}$$

So, if  $X \in T_e(\tau(G))$ , there is  $Y \in \mathfrak{gl}_n$  such that  $X = d\tau(Y) = (d\sigma - 1)Y$ . But, since  $d\sigma : A \mapsto sAs^{-1}$  acts semisimply on  $\mathfrak{gl}_n$  and preserves  $\mathfrak{g}$ , we can write  $\mathfrak{gl}_n = \mathfrak{g} \oplus V$ , with  $V$  a  $d\sigma$ -stable complement. Without loss of generality,  $Y \in \mathfrak{g}$ , so  $d\tau$  is surjective and  $\text{Lie } G_\sigma = \mathfrak{g}_\sigma$ .

Consider  $S := \{x \in \text{GL}_n \mid \text{(i), (ii), (iii)}\}$  where

- (i)  $xGx^{-1} = G$ , which implies that  $\text{Ad}(x)$  preserves  $\mathfrak{g}$
- (ii)  $m(x) = 0$ , where  $m(T) = \prod_i (T - a_i)$  is the minimal polynomial of  $s$  on  $k^n$
- (iii)  $\text{Ad}(x)$  has the same characteristic polynomial on  $\mathfrak{g}$  as  $\text{Ad}(s)$

Note that  $s \in S$ ,  $S$  is closed (check), and if  $x \in S$  then (ii) implies that  $x$  is semisimple.  $G$  acts on  $S$  by conjugation. Define  $G_x, \mathfrak{g}_x$  as  $G_\sigma, \mathfrak{g}_\sigma$  were defined. Then

$$\mathfrak{g}_x = \{X \in \mathfrak{g} \mid \text{Ad}(x)X = X\}$$

and

$$\dim \mathfrak{g}_x = \text{multiplicity of eigenvalue 1 in } \text{Ad}(x) \text{ on } \mathfrak{g} \stackrel{\text{(iii)}}{=} \dim \mathfrak{g}_\sigma$$

and

$$\dim G_x = \dim G_\sigma$$

by what we proved above. The stabilisers of the  $G$ -action on  $S$  (conjugation) all  $G_x, x \in S$ , and have the same dimension. This implies that the orbits of  $G$  on  $S$  all have the same dimension, which further gives that all orbits are closed (Proposition. 24) in  $S$  and hence in  $G$ . We have

$$\text{orbit of } s = \{gsg^{-1} \mid g \in G\} = \{g\sigma(g^{-1})s \mid g \in G\}$$

and that the map from the orbit to  $\tau(G)$  given by  $z \mapsto sz^{-1}$  is an isomorphism.  $\square$

**Corollary 96.** *If  $s \in G_s$ , then  $\text{cl}_G(s)$ , the conjugacy class of  $s$ , is closed and*

$$G \rightarrow \text{cl}_G(s), \quad g \mapsto gsg^{-1}$$

*is separable.*

**Remark 97.** *The conjugacy class of  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  in  $B_2$  is not closed!*

**Proposition 98.** *If a torus  $D$  is a closed subgroup of a connected  $G$ , then  $\text{Lie } \mathcal{Z}_G(D) = \mathfrak{z}_{\mathfrak{g}}(D)$ , where*

$$\begin{aligned}\mathcal{Z}_G(D) &= \{g \in G \mid dgd^{-1} = g \ \forall d \in D\} \text{ is the centraliser of } D \text{ in } G, \text{ and} \\ \mathfrak{z}_{\mathfrak{g}}(D) &= \{X \in \mathfrak{g} \mid \text{Ad}(d)(X) = X \ \forall d \in D\}\end{aligned}$$

**Note:**  $\mathcal{Z}_G(D) = \bigcap_{d \in D} G_d$  and  $\mathfrak{z}_{\mathfrak{g}}(D) = \bigcap_{d \in D} \mathfrak{g}_d$  ( $G_d, \mathfrak{g}_d$  as above) since, for  $d \in G_s$  and  $\text{Lie } G_d = \mathfrak{g}_d$  by above.

*Proof.* Use induction on  $\dim G$ . When  $G = 1$  this is trivial.

Case 1: If  $\mathfrak{z}_{\mathfrak{g}}(D) = \mathfrak{g}$ , then  $\mathfrak{g}_d = \mathfrak{g}$  for all  $d \in D$  so  $G_d = G$  for all  $d \in D$ , implying that  $\mathcal{Z}_G(D) = G$ .

Case 2: Otherwise, there exists  $d \in D$  such that  $\mathfrak{g}_d \subsetneq \mathfrak{g}$ . Hence,  $G_d \subsetneq G$ . Also have  $D \subset G_d^0$ , as  $D$  is connected. Note that  $\mathcal{Z}_{G_d^0}(D) = \mathcal{Z}_G(D) \cap G_d^0$  has finite index in  $\mathcal{Z}_G(D) \cap G_d = \mathcal{Z}_G(D)$  and so their Lie algebras coincide. By induction,

$$\text{Lie } \mathcal{Z}_G(D) = \text{Lie } \mathcal{Z}_{G_d^0}(D) = \mathfrak{z}_{\text{Lie } G_d^0}(D) = \mathfrak{z}_{\mathfrak{g}_d}(D) = \mathfrak{z}_{\mathfrak{g}}(D) \cap \mathfrak{g}_d = \mathfrak{z}_{\mathfrak{g}}(D)$$

□

**Proposition 99.** *If  $G$  is connected, nilpotent, then  $G_s \subset \mathcal{Z}_G$  (which implies that  $G_s$  is a subgroup).*

*Proof.* Pick  $s \in G_s$  and set  $\sigma : g \mapsto sgs^{-1}$  and  $\tau : g \mapsto \sigma(g)g^{-1} = [s, g]$ . Since  $G$  is nilpotent, there is an  $n > 0$  such that  $\tau^n(g) = [s, [s, \dots, [s, g] \cdots]] = e$  for all  $g \in G$  and so

$$\begin{aligned}\tau^n = \text{id} &\implies d\tau^n = 0 \\ &\implies d\tau = d\sigma - 1 \text{ is nilpotent, but is also semisimple by above, since } d\sigma \text{ is semisimple} \\ &\implies d\tau = 0 \\ &\implies \tau(G) = \{e\} \text{ as } G \xrightarrow{\tau} \tau(G) \text{ is separable} \\ &\implies sgs^{-1} = g \text{ for all } g \in G\end{aligned}$$

□

# 4. Quotients.

## 4.1 Existence and uniqueness as a variety.

Given a closed subgroup  $H \subset G$ , we want to give the coset space  $G/H$  the structure of a variety such that  $\pi : G \rightarrow G/H, g \mapsto gH$  is a morphism satisfying the a natural universal property.

**Proposition 100.** *There is a  $G$ -representation  $V$  and a subspace  $W \subset V$  such that*

$$H = \{g \in G \mid gW \subset W\} \quad \text{and} \quad \mathfrak{h} = \text{Lie } H = \{X \in \mathfrak{g} \mid XW \subset W\}$$

(We only need the characterisation of  $\mathfrak{h}$  when  $\text{char } k > 0$ .)

*Proof.* Let  $I = I_G(H)$ , so that  $0 \rightarrow I \rightarrow k[G] \rightarrow k[H] \rightarrow 0$ . Since  $k[G]$  is noetherian,  $I$  is finitely-generated; say,  $I = (f_1, \dots, f_n)$ . Let  $V \supset \sum k f_i$  be a finite-dimensional  $G$ -stable subspace of  $k[G]$  (with  $G$  acting by right translation). This gives a  $G$ -representation  $\rho : G \rightarrow \text{GL}(V)$ . Let  $W = V \cap I$ . If  $g \in H$ , then  $\rho(g)I \subset I \implies \rho(g)W \subset W$ . Conversely,

$$\begin{aligned} \rho(g)W \subset W &\implies \rho(g)(f_i) \in I \quad \forall i \\ &\implies \rho(g)I \subset I, \quad \text{as } \rho(g) \text{ is a ring morphism } k[G] \rightarrow k[G] \\ &\implies g \in H \quad (\text{easy exercise. Note that } \rho(g)I = I_G(Hg^{\pm 1})) \end{aligned}$$

Moreover, if  $X \in \mathfrak{h}$ , then  $d\phi(X)W \subset W$  from the above from the above. For the converse  $d\phi(X)W \subset W \implies X \in \mathfrak{h}$ , we first need a lemma.

**Lemma 101.**  $d\phi(X)f = D_X(f) \quad \forall X \in \mathfrak{g}, f \in V$

*Proof.* We know (Proposition 77) that  $d\phi(X)f = d\mathfrak{o}_f(X)$ , identifying  $V$  with  $T_f V$ , where

$$\mathfrak{o}_f : G \rightarrow V, \quad g \mapsto \rho(g)f$$

That is, for all  $f^\vee \in V^*$

$$\langle d\phi(X)f, f^\vee \rangle = \langle d\mathfrak{o}_f(X), f^\vee \rangle$$

Extend any  $f^\vee$  to  $k[G]^*$  arbitrarily. We need to show that

$$\langle d\mathfrak{o}_f(X), f^\vee \rangle = \langle D_X(f), f^\vee \rangle$$

or, equivalently,

$$X(\mathfrak{o}_f^*(f^\vee)) = \langle d\mathfrak{o}_f(X), f^\vee \rangle = \langle D_X(f), f^\vee \rangle = (1, X)\Delta f, f^\vee = (f^\vee, X)\Delta f.$$

We have

$$\mathfrak{o}_f^*(f^\vee) = f^\vee \circ \mathfrak{o}_f : g \mapsto \langle \rho(g)f, f^\vee \rangle = \langle f(\cdot g), f^\vee \rangle = \langle (\text{id}, \text{ev}_g)\Delta f, f^\vee \rangle = (f^\vee, \text{ev}_g)\Delta f$$

and so

$$\mathfrak{o}_f^*(f^\vee) = (f^\vee, \text{id})\Delta f \implies X(\mathfrak{o}_f^*(f^\vee)) = (f^\vee, X)\Delta f$$

□

Now,

$$\begin{aligned} d\phi(X)W \subset W &\implies D_X(f_i) \in I \quad \forall i \\ &\implies D_X(I) \subset I \quad (\text{as } D_X \text{ is a derivation}) \\ &\implies X(I) = 0 \quad \text{easy exercise} \end{aligned}$$

which implies that  $X$  factors through  $k[H]$ :

$$\begin{array}{ccc} k[G] & \twoheadrightarrow & k[H] \\ & \searrow X & \downarrow \bar{X} \\ & & k \end{array}$$

It is easy to see that  $\bar{X}$  is a derivation, which means that  $X \in \mathfrak{h}$ .

**Corollary 102.** *We can even demand  $\dim W = 1$  in Proposition 100 above.*

*Proof.* Let  $d = \dim W$ ,  $V' = \Lambda^d V$ , and  $W' = \Lambda^d W$ , which has dimension 1 and is contained in  $V'$ . We have actions

$$\begin{aligned} g(v_1 \wedge \cdots \wedge v_d) &= gv_1 \wedge \cdots \wedge gv_d \\ X(v_1 \wedge \cdots \wedge v_d) &= (Xv_1 \wedge \cdots \wedge v_d) + (v_1 \wedge Xv_2 \wedge \cdots \wedge v_d) + \cdots + (v_1 \wedge \cdots \wedge Xv_d) \end{aligned}$$

We need to show that

$$\begin{aligned} gW' \subset W' &\iff gW \subset W \\ XW' \subset W' &\iff XW \subset W \end{aligned}$$

which is just a lemma in linear algebra (see Springer). □

**Corollary 103.** *There is a quasiprojective homogeneous space  $X$  for  $G$  and  $x \in X$  such that*

- (i)  $\text{Stab}_G(x) = H$
- (ii) *If  $\mathfrak{o}_x : G \rightarrow X$ ,  $g \mapsto gx$ , then*

$$0 \rightarrow \text{Lie } H \rightarrow \text{Lie } G \xrightarrow{d\mathfrak{o}_x} T_x X \rightarrow 0$$

*is exact.*

Note that (ii) follows from (i) if  $\text{char } k = 0$  (use Corollaries 83 and 87.)

*Proof.* Take a line  $W \subset V$  as in the corollary above. Let  $x = [W] \in \mathbf{P}V$  and let  $X = Gx \subset \mathbf{P}V$ .  $G$  is a subvariety and is a quasiprojective homogeneous space. Then (i) is clear.  $\square$

*Exercise.* The natural map  $\phi : V - \{0\} \rightarrow \mathbf{P}V$  induces an isomorphism

$$V/x \cong T_v V/x \cong T_x(\mathbf{P}V)$$

for all  $x \in \mathbf{P}V$  and  $v \in \phi^{-1}(x)$ . (Hint:

$$k^\times \xrightarrow{\lambda \mapsto \lambda v} V - \{0\} \xrightarrow{\phi} \mathbf{P}V$$

is constant. Use an affine chart in  $\mathbf{P}V$  to prove that  $d\phi$  is surjective.)

**Claim.**  $\ker(d\mathfrak{o}_x) = \mathfrak{h}$  (then (ii) follows by dimension considerations.)  
Fix  $v \in \phi^{-1}(x)$ .

$$\begin{aligned} \phi \circ \mathfrak{o}_x : G &\xrightarrow{g \mapsto (\rho(g), v)} \text{GL}(V) \times (V - \{0\}) \xrightarrow{(\rho(g), v) \mapsto \rho(g)v} V - \{0\} \xrightarrow{\phi: \rho(g)v \mapsto [\rho(g)v]} \mathbf{P}V \\ d\phi \circ d\mathfrak{o}_x : \mathfrak{g} &\xrightarrow{X \mapsto (d\rho(X), 0)} \text{End}(V) \oplus V \xrightarrow{(d\rho(X), 0) \mapsto d\rho(X)v} V \xrightarrow{d\phi: d\rho(X)v \mapsto [d\rho(X)v]} V/x. \end{aligned}$$

We have

$$[d\phi(X)v] = 0 \iff XW \subset W \iff X \in \mathfrak{h}$$

$\square$

**Definition 104.** If  $H \subset G$  is a closed subgroup (not necessarily normal). A **quotient** of  $G$  by  $H$  is a variety  $G/H$  together with a morphism  $\pi : G \rightarrow G/H$  such that

- (i)  $\pi$  is constant on  $H$ -cosets, i.e.,  $\pi(g) = \pi(gh)$  for all  $g \in G, h \in H$ , and
- (ii) if  $G \rightarrow X$  is a morphism that is constant on  $H$ -cosets, then there exists a unique morphism  $G/H \rightarrow X$  such that

$$\begin{array}{ccc} G & \xrightarrow{\pi} & G/H \\ \downarrow & \swarrow & \\ X & & \end{array}$$

commutes. Hence, if a quotient exists, it is unique up to unique isomorphism.

**Theorem 105.** A quotient of  $G$  by  $H$  exists; it is quasiprojective. Moreover,

- (i)  $\pi : G \rightarrow G/H$  is surjective whose fibers are the  $H$ -cosets.
- (ii)  $G/H$  is a homogeneous  $G$ -space under

$$G \times G/H \rightarrow G/H, \quad (g, \pi(\gamma)) \mapsto \pi(g\gamma)$$

*Proof.* Let  $G/H = \{\text{cosets } gH\}$  as a set with natural surjection  $\pi : G \rightarrow G/H$  and give it the quotient topology (so that  $G/H$  is the quotient in the category of topological spaces).  $\pi$  is open. For  $U \subset G/H$  let  $\mathcal{O}_{G/H}(U) := \{f : U \rightarrow k \mid f \circ \pi \in \mathcal{O}_G(\pi^{-1}(U))\}$ . Easy check:  $\mathcal{O}_{G/H}$  is a sheaf of  $k$ -valued functions on  $G/H$  and so  $(G/H, \mathcal{O}_{G/H})$  is a ringed space.

If  $\phi : G \rightarrow X$  is a morphism constant on  $H$ -cosets, then we get

$$\begin{array}{ccc} G & \xrightarrow{\pi} & G/H \\ \phi \downarrow & \swarrow \exists! & \\ X & & \end{array}$$

in the category of *ringed spaces* .

By the second corollary 103 to Proposition 100 there is a quasiprojective homogeneous space  $X$  of  $G$  and  $x \in X$  such that

- (i)  $\text{Stab}_G(x) = H$
- (ii) If  $\mathfrak{o}_x : G \rightarrow X, g \mapsto gx$ , then

$$0 \rightarrow \text{Lie } H \rightarrow \text{Lie } G \xrightarrow{d\mathfrak{o}_x} T_x X \rightarrow 0$$

is exact.

Since  $\mathfrak{o}_x$  is constant on  $H$ -cosets, we get a map  $\psi : G/H \rightarrow X$  of ringed spaces (from the above universal property).  $\psi$  is necessarily given by  $gH \mapsto gx$  and is bijective. If we show that  $\psi$  is an isomorphism of ringed spaces and that  $(G/H, \mathcal{O}_{G/H})$  is a variety, then the theorem follows.

$\psi$  is a homeomorphism:

We need only show that  $\psi$  is open. If  $U \subset G/H$  is open then

$$\psi(U) = \psi(\pi(\pi^{-1}(U))) = \phi(\pi^{-1}(U))$$

is open, as  $\phi$  is.

$\psi$  gives an isomorphism of sheaves:

We must show that for  $V \subset X$  open

$$\mathcal{O}_X(V) \rightarrow \mathcal{O}_{G/H}(\psi^{-1}(V))$$

is an isomorphism of rings. Clearly it is injective. To get surjectivity we need that for all  $f : V \rightarrow k$

$$f \circ \phi : \phi^{-1}(V) \rightarrow k \text{ regular} \implies f \text{ regular}$$

Since

$$\begin{array}{ccc} G & \xrightarrow{\pi} & G/H \\ \phi \downarrow & \swarrow \psi & \\ X & & \end{array}$$

and  $\psi$  is a homeomorphism, we need only focus on  $(X, \phi)$ . A lemma:

**Lemma 106.** *Let  $X, Y$  be irreducible varieties and  $f : X \rightarrow Y$  a map of sets. If  $f$  is a morphism, then the graph  $\Gamma_f \subset X \times Y$  is closed. The converse is true if  $X$  is smooth if  $\Gamma_f$  is irreducible, and  $\Gamma_f \rightarrow X$  is separable.*

*Proof.*

( $\Rightarrow$ ): If  $f$  is a morphism, then  $\Gamma_f = \theta^{-1}(\Delta_Y)$  is closed, where

$$\theta : X \times Y \rightarrow Y \times Y, \quad (x, y) \mapsto (f(x), y).$$

( $\Leftarrow$ ): We have

$$\begin{array}{ccc} \Gamma_f & \hookrightarrow & X \times Y \\ \eta \downarrow & & \swarrow \\ & & X \end{array}$$

with  $\Gamma_f \hookrightarrow X \times Y$  the closed immersion.

$$\eta \text{ bijective} \xrightarrow{85} \dim \Gamma_f = \dim X \text{ and } 1 = [k(\Gamma_f) : k(X)]_s = [k(\Gamma_f) : k(X)]$$

as  $\eta$  is separable. Hence  $\eta$  is birational and bijective with  $X$  smooth, meaning that  $\eta$  is an isomorphism by Theorem 91 and

$$f : X \xrightarrow{\eta^{-1}} \Gamma_f \rightarrow Y$$

is a morphism. □

Now, for simplicity, assume that  $G$  is connected, which implies that  $X, V, \phi^{-1}(V)$  are irreducible. (For the general case, see Springer.) Suppose that  $f \circ \phi$  is regular. It follows from the lemma that  $\Gamma_{f \circ \phi} \subset \phi^{-1}(V) \times \mathbf{A}^1$  is closed, surjecting onto  $\Gamma_f$  via  $\phi \times \text{id}$ . By Corollary 87,  $\phi : G \rightarrow X$  is “universally open” and so

$$V \times \mathbf{A}^1 - \Gamma_{f \circ \phi} = (\phi \times \text{id})(\phi^{-1}(V) \times \mathbf{A}^1 - \Gamma_{f \circ \phi})$$

is open:  $\Gamma_f$  is closed. (The point is that  $\Gamma_{f \circ \phi}$  is a union of fibers of  $\phi \times \text{id}$ .)

Also,  $\Gamma_{f \circ \phi} \cong \phi^{-1}(V)$  is irreducible, implying that  $\Gamma_f$  is irreducible, and

$$\begin{array}{ccc} \Gamma_{f \circ \phi} & \xrightarrow[\text{pr}_1]{\sim} & \phi^{-1}(V) \\ \downarrow & & \downarrow \\ \Gamma_f & \xrightarrow[\text{pr}_1]{} & V \end{array}$$

and

$$d\phi \text{ surjective} \implies d(\text{pr}_1) \text{ surjective} \implies \Gamma_f \rightarrow V \text{ separable and } V \text{ smooth.}$$

By Lemma 106,  $f$  is a morphism. □

**Corollary 107.** (i)  $\dim(G/H) = \dim G - \dim H$

(ii)

$$0 \rightarrow \text{Lie } H \rightarrow \text{Lie } G \xrightarrow{d\pi} T_e(G/H) \rightarrow 0$$

is exact.

*Proof.*

(i):  $G/H$  is a homogeneous with stabilisers equal to  $H$ .

(ii): Implied by Corollary 103. □

**Exercise.** Recall that a sequence  $1 \rightarrow K \xrightarrow{\phi} G \xrightarrow{\psi} H \rightarrow 1$  of algebraic groups is *exact* if (i) it is set-theoretically and (ii)  $0 \rightarrow \text{Lie } K \xrightarrow{d\phi} \text{Lie } G \xrightarrow{d\psi} \text{Lie } H \rightarrow 0$  is exact.

(a) Show that  $\phi$  is a closed immersion if and only if  $\phi$  is injective and  $d\phi$  injective.

(b) Show that  $\psi$  is separable if and only if  $\psi$  is surjective and  $d\psi$  surjective.

(c) Deduce that the sequence is exact if and only if (i) as above and (ii')  $\phi$  is a closed immersion and  $\psi$  is separable.

**Lemma 108.** *Let  $H_1 \subset G_1, H_2 \subset G_2$  be closed subgroups. The natural map*

$$(G_1 \times G_2)/(H_1 \times H_2) \rightarrow G_1/H_1 \times G_2/H_2$$

*is an isomorphism.*

*Proof.* This is a bijective map of homogeneous  $G_1 \times G_2$  spaces, which is bijective on tangent spaces by the above. The rest follows from Corollary 89. □

## 4.2 Quotient algebraic groups.

**Proposition 109.** *Suppose that  $N \trianglelefteq G$  is a closed normal subgroup. Then  $G/N$  is an algebraic group that is affine (and  $\pi : G \rightarrow G/N$  is a morphism of algebraic groups).*

*Proof.* Inversion  $G/N \rightarrow G/N$  is a morphism, along with multiplication  $G/N \times G/N \rightarrow G/N$  by Lemma 108, which gives that  $G/N$  is an algebraic group.

By Corollary 102, there exists a  $G$ -representation  $\rho : G \rightarrow \text{GL}(V)$  and a line  $L \subset V$  such that  $N = \text{Stab}_G(L)$  and  $\text{Lie } N = \text{Stab}_{\mathfrak{g}}(L)$ . For  $\chi \in X(N)$ , let  $V_\chi$  be the  $\chi$ -eigenspace of  $V$ . (Note that  $L \subset V_\chi$  for some  $\chi$ .) Let  $V' = \sum_{\chi \in X(H)} V_\chi = \bigoplus_{\chi} V_\chi$ . As  $N \trianglelefteq G$ ,  $G$  permutes the  $V_\chi$ . Define

$$W = \{f \in \text{End}(V) \mid f(V_\chi) \subset V_\chi \ \forall \chi\} \subset \text{End}(V).$$

Let  $\sigma : G \rightarrow \text{GL}(W)$  by

$$\sigma(g)f := \rho(g)f\rho(g)^{-1}$$

which is an algebraic representation.

**Claim.**  $\sigma$  induces a closed immersion  $G/N \hookrightarrow \text{GL}(W)$ .

It is enough to show that  $\ker \sigma = N$  and  $\ker(d\sigma) = \text{Lie } N$ .

$$\begin{aligned} g \in \ker \sigma &\iff \rho(g)f = f\rho(g) \\ &\iff \rho(g) \text{ acts as a scalar on each } V_\chi \\ &\implies \rho(g)L = L \text{ as } L \subset V_\chi \text{ for some } \chi \\ &\implies g \in N \end{aligned}$$

The converse is trivial:  $\ker \sigma = N$ .

By Proposition 77,  $\phi_f : G \rightarrow W$ ,  $g \mapsto \sigma(g)f$  has derivative

$$d\phi : \mathfrak{g} \rightarrow W, \quad X \mapsto d\sigma(X)f.$$

Check that  $d\sigma(X)f = d\phi(X)f - fd\phi(X)$ . We have

$$\begin{aligned} d\sigma(X) = 0 &\iff d\phi(X)f = fd\phi(X) \quad \text{for all } f \in W \\ &\iff && d\phi(X) \text{ acts as a scalar on each } V_\chi \\ &\implies X \in \text{Lie } N \text{ (as above)} \end{aligned}$$

□

**Corollary 110.** *Suppose  $\phi : G \rightarrow H$  is a morphism of algebraic groups with  $\phi(N) = 1$ ,  $N \trianglelefteq G$ . Then*

$$\begin{array}{ccc} G & \longrightarrow & G/N \\ & \searrow \phi & \downarrow \\ & & H \end{array}$$

*In particular, we get that  $G/\ker \phi \rightarrow \text{im } \phi$  is bijective and is an isomorphism when in characteristic 0.*

(Note that in characteristic  $p$ ,  $\mathbf{G}_m \xrightarrow{p} \mathbf{G}_m$  is bijective and not an isomorphism.)

**Remark 111.**

$$1 \rightarrow N \rightarrow G \rightarrow G/N \rightarrow 1$$

*is exact by Corollary 107.*

**Exercise.** If  $N \subset H \subset G$  are closed subgroups with  $N \trianglelefteq G$ , then the natural map  $H/N \rightarrow G/N$  is a closed immersion (so we can think of  $H/N$  as a closed subgroup of  $G/N$ ) and we have a canonical isomorphism  $(G/N)/(H/N) \xrightarrow{\sim} G/H$ . of homogeneous  $G$ -spaces.

**Exercise.** Assume that  $\text{char } k = 0$ . Suppose  $N, H \subset G$  are closed subgroups such that  $H$  normalises  $N$ . Show that  $HN$  is a closed subgroup of  $G$  and that we have a canonical isomorphism  $HN/N \cong H/(H \cap N)$  of algebraic groups. Find a counterexample when  $\text{char } k > 0$ .

**Exercise.** Suppose  $H$  is a closed subgroup of an algebraic group  $G$ . Show that if both  $H$  and  $G/H$  are connected, then  $G$  is connected. (Use, for example, Exercise 5.5.9 (1) in Springer.)

**Exercise.** Suppose  $\phi : G \rightarrow H$  is a morphism of algebraic groups. If  $H_1 \subset H_2 \subset H$  are closed subgroups, show that we have a canonical isomorphism  $\phi^{-1}(H_2)/\phi^{-1}(H_1) \xrightarrow{\sim} H_2/H_1$ . (Hint: show  $\text{Lie } \phi^{-1}(H_i) = (d\phi^{-1})^{-1}(\text{Lie } H_i)$ .)

*Example.* The group  $\mathbf{PSL}_2$ :

Let  $Z = \left\{ \begin{pmatrix} x & \\ & x \end{pmatrix} \mid x \in \mathbf{G}_m \right\}$ .  $\text{GL}_2/Z$  is affine and the composition

$$\text{SL}_2 \hookrightarrow \text{GL}_2 \twoheadrightarrow \text{GL}_2/Z$$

is surjective, inducing the inclusion of Hopf algebras

$$k[\mathrm{GL}_2]^Z = k[\mathrm{GL}_2/Z] \hookrightarrow k[\mathrm{SL}_2].$$

Check that the image is generated by the elements  $\frac{T_i T_j}{\det^2}$ . (See Springer Exercise 2.1.5(3).)

# 5. Parabolic and Borel subgroups.

## 5.1 Complete varieties.

**Recall:** A variety  $X$  is **complete** if for all varieties  $Z$ ,  $X \times Z \xrightarrow{\text{pr}_2} Z$  is a closed map. In the category of locally compact Hausdorff topological spaces, the analogous property is equivalent to compactness.

**Proposition 112.** *Let  $X$  be complete.*

- (i)  $Y \subset X$  closed  $\implies Y$  complete.
- (ii)  $Y$  complete  $\implies X \times Y$  complete
- (iii)  $\phi : X \rightarrow Y$  morphism  $\implies \phi(X) \subset Y$  is closed and complete, which implies that if  $X \subset Z$  is a subvariety, then  $X$  is closed in  $Z$
- (iv)  $X$  irreducible  $\implies \mathcal{O}_X(X) = k$
- (v)  $X$  affine  $\implies X$  finite

*Proof.* An exercise (or one can look in Springer). □

**Theorem 113.**  $X$  projective  $\implies X$  complete

**Note:** The converse is not true.

**Lemma 114.** *Let  $X, Y$  be homogeneous  $G$ -spaces with  $\phi : X \rightarrow Y$  a bijective  $G$ -map. Then  $X$  is complete  $\iff Y$  is complete.*

*Proof.* For all varieties  $Z$ , then projection  $X \times Z \rightarrow Z$  can be factored as

$$X \times Z \xrightarrow{\phi \times \text{id}} Y \times Z \xrightarrow{\text{pr}_2} Z$$

$\phi \times \text{id}$  is bijective and open (by Corollary 87) and is thus a homeomorphism:  $Y$  being complete implies that in  $X$ . Applying the same reasoning to  $\phi^{-1} : Y \rightarrow X$  gives the converse. □

**Definition 115.** *A closed subgroup  $P \subset G$  is **parabolic** if  $G/P$  is complete.*

**Remark 116.** For a closed subgroup  $P \subset G$ ,  $G/P$  is quasi-projective by Theorem 105 and so

$$G/P \text{ projective} \iff G/P \text{ complete} \iff P \text{ parabolic.}$$

The implication of  $G/P$  being complete implying that  $G/P$  being projective follows from Proposition 112 (iii) applying to the embedding of  $G/P$  into some projective space.

**Proposition 117.** If  $Q \subset P$  and  $P \subset G$  are parabolic, then  $Q \subset G$  is parabolic.

*Proof.* For all varieties  $Z$  we need to show that  $G/Q \times Z \xrightarrow{\text{pr}_2} Z$  is closed. Fix a closed subset  $C \subset G/Q \times Z$ . Letting  $\pi : G \rightarrow G/P$  denote the natural projection, set  $D = (\pi \times \text{id}_Z)^{-1}(C) \subset G \times Z$ , which is closed. For all  $q \in Q$ , note that  $(g, z) \in D \implies (gq, z) \in D$ . It is enough to show that  $\text{pr}_2(D) \subset Z$  is closed.

Let

$$\theta : P \times G \times Z \rightarrow G \times Z, \quad (p, g, z) \mapsto (gp, z)$$

Then  $\theta^{-1}(D)$  is closed for all  $q \in Q$

$$(*) \quad (p, g, z) \in \theta^{-1}(D) \implies (pq, g, z) \in \theta^{-1}(D)$$

Let  $\alpha : P \times G \times Z \rightarrow P/Q \times G \times Z$  be the natural map.

$$\begin{array}{ccc} P \times G \times Z & \xrightarrow{\alpha} & P/Q \times G \times Z \\ & \searrow \text{pr}_{23} & \downarrow \text{pr}_{23} \\ & & G \times Z \end{array}$$

By Corollary 87  $\alpha$  is open.  $(*)$  implies that  $\alpha(\theta^{-1}(D))$  is closed.  $P/Q$  being complete implies that

$$\text{pr}_{23}(\theta^{-1}(D)) = \{(gp^{-1}, z) \mid (g, z) \in D, p \in P\}$$

is closed. Now,

$$\begin{array}{ccc} G \times Z & \xrightarrow{\beta} & Z \\ & \searrow \text{pr}_2 & \downarrow \text{pr}_2 \\ & & G \times Z \end{array}$$

Similarly  $\beta$  is open, and so  $\beta(\text{pr}_{23}(\theta^{-1}(D)))$  is closed.  $G/P$  being complete implies

$$\text{pr}_2(\beta(\text{pr}_{23}(\theta^{-1}(D)))) = \text{pr}_2(\text{pr}_{23}(\theta^{-1}(D))) = \text{pr}_2(D) = \text{pr}_2(C)$$

is closed. □

## 5.2 Borel subgroups.

**Theorem 118** (Borel's fixed point theorem). *Let  $G$  be a connected, solvable algebraic group and  $X$  a (nonempty) complete  $G$ -space. Then  $X$  has a fixed point.*

*Proof.* We show this by inducting on the dimension of  $G$ . When  $\dim G = 0 \implies G = \{e\}$  the theorem trivially holds. Now, let  $\dim G > 0$  and suppose that the theorem holds for dimensions less than  $\dim G$ . Let  $N = [G, G] \trianglelefteq G$ , which is a connected normal subgroup by Proposition 19 and is a proper subgroup as  $G$  is solvable. Since  $N$  is connected and solvable, by induction

$$X^N = \{x \in X \mid nx = x \ \forall n \in N\} \neq \emptyset$$

Since  $X^N \subset X$  is closed (both topologically and under the action of  $G$ , as  $N$  is normal), by Proposition 112,  $X^N$  is complete; so, without loss of generality suppose that  $N$  acts trivially on  $X$ . Pick a closed orbit  $Gx \subset X$ , which exists by Proposition 24 and is complete. Since  $G/\text{Stab}_G(x) \rightarrow Gx$  is a bijective map of homogeneous  $G$ -spaces,  $G/\text{Stab}_G(x)$  is complete by Proposition 114.

$$\begin{aligned} N \subset \text{Stab}_G(x) &\implies \text{Stab}_G(x) \text{ is normal} \\ &\implies G/\text{Stab}_G(x) \text{ is affine and complete (and connected)} \\ &\implies G/\text{Stab}_G(x) \text{ is a point, by Proposition 112} \\ &\implies x \in X^G \end{aligned}$$

□

**Proposition 119** (Lie-Kolchin). *Suppose that  $G$  is connected and solvable. If  $\phi : G \rightarrow \text{GL}_n$ , then there exists  $\gamma \in \text{GL}_n$  such that  $\gamma(\text{im } \phi)\gamma^{-1} \subset B_n$ .*

*Proof.* Induct on  $n$ . When  $n = 1$ , then theorem trivially holds. Let  $n > 1$  and suppose that it holds for all  $m < n$ . Write  $\text{GL}_n = \text{GL}(V)$  for an  $n$ -dimensional vector space  $V$ .  $G$  acts on  $\mathbf{P}V$  via  $\phi$ . By Borel's fixed point theorem, there exists  $v_1 \in V$  such that  $G$  stabilises the line  $V_1 := kv_1 \subset V$ , implying that  $G$  acts on  $V/V_1$ . By induction there exists a flag

$$0 = V_1/V_1 \subsetneq V_2/V_1 \subsetneq \cdots \subsetneq V/V_1$$

stabilised by  $G$ ; hence  $G$  stabilises the flag

$$0 \subsetneq V_1 \subsetneq \cdots \subsetneq V_n = V$$

□

**Definition 120.** *A Borel subgroup of  $G$  is a closed subgroup  $B$  of  $G$  that is maximal among connected solvable subgroups.*

**Remarks 121.**

- Any  $G$  has a Borel subgroup since if  $B_1 \subsetneq B_2$  is irreducible  $\implies \dim B_1 < \dim B_2$ .
- $B_n \subset \text{GL}_n$  is a Borel by Lie-Kolchin.

**Theorem 122.**

- (i) A closed subgroup  $P \subset G$  is parabolic  $\iff P$  contains a Borel subgroup.
- (ii) Any two Borel subgroups are conjugate.

*In particular, a Borel subgroup is precisely a minimal - or, equivalently, a connected, solvable - parabolic.*

*Proof.* For simplicity, assume that  $G$  is connected.

(i) ( $\Rightarrow$ ): Suppose that  $B$  is a Borel and  $P$  is parabolic.  $B$  acts on  $G/P$ . By the Borel fixed point theorem, there is a coset  $gP$  such that  $Bg \subset gP \implies g^{-1}Bg \subset P$ .  $g^{-1}Bg$  is Borel.

(i) ( $\Leftarrow$ ): Let  $B$  be a Borel. We first show that  $B$  is parabolic, inducting on  $\dim G$ . Pick a closed immersion  $G \hookrightarrow \mathrm{GL}(V)$ .  $G$  acts on  $\mathbf{P}V$ . Let  $Gx$  be a closed - hence complete - orbit. Since  $G/\mathrm{Stab}_G(x) \rightarrow Gx$  is a bijective map of homogeneous spaces,  $P := \mathrm{Stab}_G(x)$  is parabolic. By above,  $B \subset gPg^{-1}$ , for some  $g \in G$ . Without loss of generality,  $B \subset P$ . If  $P \neq G$ , then  $B$  is Borel in  $P$ . Since  $P \subset G$  is parabolic and  $B \subset P$  is parabolic by induction, it follows that  $B \subset G$  is parabolic, by Proposition 117. Suppose  $P = G$ .  $G$  stabilises some line  $V_1 \subset V$ , which gives a morphism  $G \rightarrow \mathrm{GL}(V/V_1)$ . By induction on  $\dim V$ , we either obtain a proper parabolic subgroup, in which case we are done by the above, or  $G$  stabilises some flag  $0 \subset V_1 \subset \cdots \subset V_n = V$ , giving that

$$G \hookrightarrow B_n \implies G \text{ is solvable} \implies G = B \text{ is parabolic}$$

Now, suppose that  $P$  is a closed subgroup containing a Borel  $B$ . Then  $G/B \rightarrow G/P$ . Since  $G/B$  is complete, by Proposition 112 we get that  $G/P$  is complete  $\implies P$  is parabolic.

(ii). Let  $B_1, B_2$  be Borel subgroups, which are parabolic by (i). By (i), there is  $g \in G$  such that  $gB_1g^{-1} \subset B_2 \implies \dim B_1 \leq \dim B_2$ . Similarly,

$$\dim B_2 \leq \dim B_1 \implies \dim B_1 = \dim B_2 \implies gB_1g^{-1} = B_2$$

□

**Corollary 123.** *Let  $\phi : G \rightarrow G'$  be a surjective morphism of algebraic groups.*

(i) *If  $B \subset G$  is Borel, then  $\phi(B) \subset G'$  is Borel.*

(ii) *If  $P \subset G$  is parabolic, then  $\phi(P) \subset G'$  is parabolic.*

*Proof.* It is enough to prove (i). Since  $B \rightarrow \phi(B)$ ,  $\phi(B)$  is connected and solvable. Since  $G/B$  is complete and  $G/B \rightarrow G'/\phi(B)$  it follows that  $G'/\phi(B)$  is complete and  $\phi(B)$  is parabolic. Now,  $\phi(B)$  is connected, solvable, and contains a Borel:  $\phi(B)$  is Borel by the maximality in the definition of a Borel subgroup. □

**Corollary 124.** *If  $G$  be connected and  $B \subset G$  Borel, then  $\mathcal{Z}_G^0 \subset \mathcal{Z}_B \subset \mathcal{Z}_G$ .*

*Proof.*

$$\begin{aligned} \mathcal{Z}_G^0 \text{ connected, solvable} &\implies \mathcal{Z}_G^0 \subset gBg^{-1}, \text{ for some } g \in G \\ &\implies \mathcal{Z}_G^0 = g^{-1}\mathcal{Z}_G^0g \subset B \\ &\implies \mathcal{Z}_G^0 \subset \mathcal{Z}_B \end{aligned}$$

Now, fix  $b \in \mathcal{Z}_B$  and define the morphism  $\phi : G/B \rightarrow G$  of varieties by  $gB \mapsto gbg^{-1}$ .  $\phi(G/B)$  is complete and closed - hence affine - and irreducible:

$$\phi(G/B) = \{b\} \implies \forall g \in G gbg^{-1} = b \implies b \in \mathcal{Z}_G \implies \mathcal{Z}_B \subset \mathcal{Z}_G$$

□

**Proposition 125.** *Let  $G$  be a connected group and  $B \subset G$  a Borel. If  $B$  is nilpotent, then  $G$  is solvable; that is,  $B$  nilpotent  $\implies B = G$ .*

$B$  being nilpotent means that

$$B \supsetneq CB \supsetneq \cdots \supsetneq C^n B = 1$$

for some  $n$  (where  $C^i B = [B, C^{i-1} B]$  is connected and closed). Let  $N = C^{n-1} B$ , so that

$$1 = [B, N] \implies N \subset \mathcal{Z}_B \subset \mathcal{Z}_G \text{ (above corollary)} \implies N \trianglelefteq G$$

Hence we have the morphism  $B/N \hookrightarrow G/N$  of algebraic groups, which is a closed immersion by the exercise after Theorems 85, 86. Also,  $B/N$  is a Borel of  $G/N$ , by the corollary above, and  $B/N$  is nilpotent.

Inducting on  $\dim G$ , we get that  $G/N$  is solvable, which implies that  $G$  is solvable.  $\square$

### 5.3 Structure of solvable groups.

**Proposition 126.** *Let  $G$  be connected and nilpotent. Then  $G_s, G_u$  are (connected) closed normal subgroups and  $G_s \times G_u \xrightarrow{\text{mult.}} G$  is an isomorphism of algebraic groups. Moreover,  $G_s$  is a central torus.*

*Proof.* Without loss of generality,  $G \subset \text{GL}(V)$  is a closed subgroup. By Proposition 99  $G_s \subset \mathcal{Z}_G$ . The eigenspaces of elements  $G_s$  coincide; let  $V = \bigoplus_{\lambda: G_s \rightarrow k^\times} V_\lambda$  be a simultaneous eigenspace decomposition. Since  $G_s$  is central,  $G$  preserves each  $V_\lambda$ . By Lie-Kolchin (Proposition 119), we can choose a basis for each  $V_\lambda$  such that the  $G$ -action is upper-triangular. Therefore,  $G \subset B_n$ , and  $G_s = G \cap D_n$ ,  $G_u = G \cap U_n$  are closed subgroups,  $G_u$  being normal. We can now show that  $G_s \times G_u \xrightarrow{\sim} G$  as in the proof of Proposition 37. Moreover,  $G_s$  is a torus, being connected and commutative.  $\square$

**Proposition 127.** *Let  $G$  be connected and solvable.*

- (i)  $[G, G]$  is a connected, normal closed subgroup and is unipotent.
- (ii)  $G_u$  is a connected, normal closed subgroup and  $G/G_u$  is a torus.

*Proof.*

(i).

$$\begin{aligned} \text{Lie-Kolchin} &\implies G \hookrightarrow B_n \\ &\implies [G, G] \hookrightarrow [B_n, B_n] \subset U_n \\ &\implies [G, G] \text{ unipotent} \end{aligned}$$

We already know that it is connected, closed, and normal.

(ii).  $G_u = G \cap U_n$  is a closed subgroup.  $G_u \supset [G, G]$  implies that  $G_u \trianglelefteq G$  and that  $G/G_u$  is commutative. For  $[g] \in G/G_u$ ,  $[g] = [g_s] = [g]_s$ : all elements of  $G/G_u$  are semisimple. Since  $G/G_u$

is furthermore connected, it follows that  $G/G_u$  is a torus. It now remains to show that  $G_u$  is connected.

$$1 \rightarrow G_u/[G, G] \rightarrow G/[G, G] \rightarrow G/G_u \rightarrow 1$$

is exact (by the exercise on exact sequences). By Proposition 37,

$$G/[G, G] \cong (G/[G, G])_s \times (G/[G, G])_u$$

Hence  $(G/[G, G])_u = G_u/[G, G]$ , which is connected by the above. Since  $[G, G]$  is also connected, it follows from Springer 5.5.9.(1) (exercise) that  $G_u$  is connected.  $\square$

**Lemma 128.** *Let  $G$  be connected and solvable with  $G_u \neq 1$ . Then there exists a closed subgroup  $N \subset \mathcal{Z}_{G_u}$  such that  $N \cong \mathbf{G}_a$  and  $N \trianglelefteq G$ .*

*Proof.* Since  $G_u$  is unipotent, it is nilpotent. Let  $n$  be such that

$$G_u \supseteq \mathcal{C}G_u \supseteq \cdots \supseteq \mathcal{C}^n G_u = 1$$

The  $\mathcal{C}^i G_u$  are connected closed subgroups and are normal as  $G_u$  is normal. Let  $N = \mathcal{C}^{n-1} G_u$ . Then

$$1 = [G_u, N] \implies N \subset \mathcal{Z}_G(G_u)$$

If  $\text{char } k = p > 0$ , let  $N \hookrightarrow U_m$ , for some  $m$ , and let  $r$  be the minimal such that  $p^r \geq n$  so that  $N^{p^r} = 1$ . then

$$N \supseteq N^p \supseteq \cdots \supseteq N^{p^r} = 1$$

The  $N^{p^i}$  are connected, closed, and normal. Without loss of generality, suppose that  $r = 1$  taking  $N^{p^{r-1}}$  otherwise. Then  $N$  is a connected elementary unipotent group and hence is isomorphic to  $\mathbf{G}_a^r$  for some  $r$ , by Corollary 57.

$G$  act on  $N$  by conjugation, with  $G_u$  acting trivially. This induces an action  $G/G_u \times N \rightarrow N$  (use Lemma 108).  $G/G_u$  acts on  $k[N]$  in a locally algebraic manner, preserving  $\text{Hom}(N, \mathbf{G}_a) = \mathcal{A}(N)$ . Since  $G/G_u$  is a torus, there is a nonzero  $f \in \text{Hom}(N, \mathbf{G}_a)$  that is a simultaneous eigenvector. So,  $(\ker f)^0 \subset N$  has dimension  $r - 1$  and is still normal in  $G$ . Induct on  $r$ .  $\square$

**Definitions 129.** *A maximal torus of  $G$  is a closed subgroup that is a torus and is a maximal such subgroup with respect to inclusion; they exist by dimension considerations. A temporary definition: a torus  $T$  of a connected solvable group is **Maximal** (versus maximal) if  $\dim T = \dim(G/G_u)$ . (Recall that  $G/G_u$  is a torus). It is easy to see that **Maximal**  $\implies$  *maximal*. We shall soon see that the converse is true as well, after a corollary to the following theorem (so that we can then dispense with the capital M):*

**Theorem 130.** *Let  $G$  be connected and solvable.*

- (i) *Any semisimple element lies in a Maximal torus. (In particular, Maximal tori exist.)*
- (ii)  *$\mathcal{Z}_G(s)$  is connected for all semisimple  $s$ .*
- (iii) *Any two Maximal tori are conjugate in  $G$ .*
- (iv) *If  $T$  is a Maximal torus, then  $G \cong G_u \rtimes T$  (i.e.,  $G_u \trianglelefteq G$  and  $G_u \times T \xrightarrow{\text{mult.}} G$  is an isomorphism of varieties).*

*Proof.*

(iv): Let  $T$  be Maximal and consider  $\phi : T \rightarrow G/G_u$ . Since  $\ker \phi = T \cap G_u = 1$  (Jordan decomposition), we have that

$$\dim \phi(T) = \dim T - \dim \ker \phi = \dim T = \dim G/G_u \implies \phi(T) = G/G_u :$$

$\phi$  is surjective and so  $G = TG_u$ . Thus multiplication  $T \times G_u \rightarrow G$  is a bijective map of homogeneous  $T \times G_u$ -spaces. To see that it is an isomorphism, (if  $p > 0$ ) we need an isomorphism - just an injection by dimension considerations - on Lie algebras, which is equivalent to  $\text{Lie } T \cap \text{Lie } G_u = 0$ , as is to be shown.

Now, pick a closed immersion  $G \hookrightarrow \text{GL}(V)$ . Picking a basis for  $V$  such that  $G_u \subset U_n$  gives that

$$\text{Lie } G_u \subset \text{Lie } U_n = \begin{pmatrix} 0 & * & * \\ & \ddots & * \\ & & 0 \end{pmatrix}$$

consists of nilpotent elements. Picking a basis for  $V$  such that  $T \subset D_n$  gives that

$$\text{Lie } T \subset \text{Lie } D_n = \text{diag}(*, \dots, *)$$

consist of semisimple elements. Thus,  $\text{Lie } T \cap \text{Lie } G_u = 0$ .

(i)-(iii):

If  $G_u = 1$ , then  $G$  is a torus and there is nothing to show. Suppose that  $\dim G_u > 0$ .

Case 1.  $\dim G_u = 1$ :

$G_u$  is connected, unipotent and so  $G_u \cong \mathbf{G}_a$  by Theorem 58. Let  $\phi : \mathbf{G}_a \rightarrow G_u$  be an isomorphism.  $G$  acts on  $G_u$  by conjugation with  $G_u$  acting trivially. We have

$$\text{Aut } G_u \cong \text{Aut } \mathbf{G}_a \cong \mathbf{G}_m \text{ (exercise).}$$

Hence

$$g\phi(x)g^{-1} = \phi(\alpha(g)x)$$

for all  $g \in G, x \in \mathbf{G}_a$ , for some character  $\alpha : G/G_u \rightarrow \mathbf{G}_m$ .

$\alpha = 1$ :  $G_u \subset Z_G$ .

$$\begin{aligned} [G, G] \subset G_u \text{ (Proposition 127)} &\implies [G, [G, G]] = 1, \text{ so } G \text{ is nilpotent} \\ &\implies G \cong G_u \times G_s \text{ (Proposition 126)} \end{aligned}$$

and so  $G$  is commutative and  $G_s$  is the unique maximal torus. (i)-(iii) are immediate.

$\alpha \neq 1$ : Given  $s \in G_s$ , let  $Z = Z_G(s)$ .

$$\begin{aligned} G/G_u \text{ commutative} &\implies \text{cl}_G(s) \text{ maps to } [s] \in G/G_u \\ &\implies \text{cl}_G(s) \subset sG_u \\ &\implies \dim \text{cl}_G(s) \leq 1 \\ &\implies \dim Z = \dim G - \dim \text{cl}_G(s) \geq \dim G - 1 \end{aligned}$$

$\alpha(s) \neq 1$ : For all  $x \neq 0$

$$s\phi(x)s^{-1} = \phi(\alpha(s)x) \neq \phi(x)$$

which implies that  $Z \cap G_u = 1$ , further giving  $\dim Z = \dim G - 1$  and

$$\begin{aligned} Z_u = 1 &\implies Z^0 \text{ is a torus - which is Maximal - by Proposition 127 (it is connected, solvable and } Z_u^0 = 1) \\ &\implies G = Z^0 G_u, \quad \text{by (iv)} \end{aligned}$$

If  $z \in Z$ , then  $z = z_0 u$  for some  $z_0 \in Z^0$  and  $u \in G_u$ . But

$$u = z_0^{-1} z \in Z \cap G_u = 1 \implies z = z_0 \in Z^0.$$

Therefore,  $Z = Z^0$ , giving (iii), and  $s \in Z$ , giving (i).

$\alpha(s) = 1$ : For all  $x \neq 0$

$$s\phi(x)s^{-1} = \phi(\alpha(s)x) = \phi(x)$$

and so  $G_u \subset Z$ . By the Jordan decomposition, since  $s$  commutes with  $G_u$ ,  $sG_u \cap G_s = \{s\}$ , which means that

$$\text{cl}_G(s) = \{s\} \implies s \in \mathcal{Z}_G \implies Z = G.$$

(ii) follows.

Note that since  $\alpha \neq 1$  there is  $g = g_s g_u$  such that  $\alpha(g_s) = \alpha(g) \neq 1$  and so  $\mathcal{Z}(g_s)$  is a Maximal torus by the previous case. Hence, since  $\mathcal{Z}_G(s) = G$ , we have  $s \in \mathcal{Z}_G(g_s)$ : (i) follows.

Now it remains to prove (iii) in the general case in which  $\alpha \neq 1$ . Let  $s$  be such that  $T, T'$  be Maximal tori. With the identification  $T \xrightarrow{\sim} G/G_u$  (see (iv)), let  $s \in T$  be such that  $\alpha(s) \neq 1$ . Then  $\mathcal{Z}_G(s)$  is Maximal (by the above) and

$$T \subset \mathcal{Z}_G(s) \implies T = \mathcal{Z}_G(s) \quad \text{by dimension considerations.}$$

Likewise, with the identification  $T' \xrightarrow{\sim} G/G_u$ , pick  $s' \in T'$  with  $[s] = [s']$  in  $G/G_u$  so that  $T' = \mathcal{Z}_G(s')$ .  $s' = su$  for some  $u \in G_u$ . The conjugacy class of  $s$  (resp.  $s'$ ) - which has dimension 1 by the above - is contained in  $sG_u = s'G_u$ , which is irreducible of dimension 1:

$$\text{cl}_G(s) = sG_u = s'G_u = \text{cl}_G(s')$$

since the conjugacy classes are closed (Corollary 96). Therefore,  $s'$  is conjugate to  $s$  and thus  $T, T'$  are conjugate.

Case 2.  $\dim G_u > 1$ : Induct on the dimension of  $G$ .

Lemma 128 implies that there exists a closed, normal subgroup  $N \subset \mathcal{Z}_{G_u}$  isomorphic to  $\mathbf{G}_a$ . Set  $\overline{G} = G/N$  and  $\overline{G}_u = G_u/N$ , so  $\overline{G}/\overline{G}_u \cong G/G_u$ . Let  $\pi : G \rightarrow \overline{G}$  be the natural surjection.

(i): If  $s \in G_s$ , define  $\bar{s} = \pi(s) \in \overline{G}_s := \pi(G_s)$ . By induction, there is a Maximal torus  $\overline{T}$  in  $\overline{G}$  containing  $\bar{s}$ . Let  $H = \pi^{-1}(\overline{T})$ , which is connected since  $N$  and  $H/N \cong \overline{T}$  (exercise) is connected. Also,  $H_u = N$  (as  $H/N \cong \overline{T}$ ) has dimension 1. Case 1 implies that there is a torus  $T \ni s$  in  $H$  (Maximal in  $H$ ) of dimension  $\dim H/H_u = \dim \overline{T} = \dim G/G_u$ ; hence,  $T$  is Maximal

in  $G$ , containing  $s$ .

(iii): Let  $T, T'$  be Maximal tori. Then  $\pi(T) = \pi(T')$  are Maximal tori in  $\overline{G}$  and by induction are conjugate: there is  $g \in G$  such that

$$\pi(T) = \pi(gT'g^{-1}) \implies T, gT'g^{-1} \in \pi^{-1}(\pi(T)) =: H.$$

As above  $H_u$  is 1-dimensional and so  $T, gT'g^{-1}$  - being Maximal tori in  $H$  - are conjugate in  $H$  and hence in  $G$ .

(ii): Again, for  $s \in G_s$ , set  $\bar{s} = \pi(s)$ .  $\mathcal{Z}_{\overline{G}}(\bar{s})$  is connected by induction.  $H := \pi^{-1}(\mathcal{Z}_{\overline{G}}(\bar{s}))$  is connected since  $N$  and  $H/N \cong \mathcal{Z}_{\overline{G}}(\bar{s})$  (exercise) are connected. Since  $\pi(\mathcal{Z}_G(s)) \subset \mathcal{Z}_{\overline{G}}(\bar{s})$ , we have  $\mathcal{Z}_G(s) = \mathcal{Z}_H(s)$ . If  $H \neq G$ ,  $\mathcal{Z}_H(s)$  is connected by induction and we are done. If  $H = G$ , then  $\mathcal{Z}_{\overline{G}}(\bar{s}) = \overline{G}$ . Hence,

$$\text{cl}_G(\bar{s}) = \{\bar{s}\} \implies \text{cl}_G(s) \subset \pi^{-1}(\bar{s}) = sN$$

and so the conjugacy class of  $s$  has dimension 0 or 1. In the former case,  $\mathcal{Z}_G(s) = G$  is connected and we are done. In the second, conjugating by  $s$  gives rise to an  $\alpha : G/G_u \rightarrow \text{Aut}(N) \cong \mathbf{G}_m$  and we can proceed as in Case 1...  $\square$

*Example.*  $D_n$  is a maximal torus of  $B_n$  and  $B_n \cong U_n \rtimes D_n$ .

**Remark 131.** (i), (iii) above carry over to all connected  $G$ , as we shall see soon. However, (ii) can fail in general. (For example, take  $G = \text{PSL}_2$  in characteristic  $\neq 2$  and  $s = [\text{diag}(1, -1)]$ .)

**Lemma 132.** If  $\phi : H \rightarrow G$  is injective, then  $\dim H \leq \dim G$ .

*Proof.* Since  $\dim \ker \phi = 0$ ,  $\dim H = \dim \phi(H) \leq \dim G$ .  $\square$

**Proposition 133.** Let  $G$  be connected and solvable with  $H \subset G$  a closed diagonalisable subgroup.

- (i)  $H$  is contained in a Maximal torus.
- (ii)  $\mathcal{Z}_G(H)$  is connected.
- (iii)  $\mathcal{Z}_G(H) = N_G(H)$

*Proof.* We shall induct on  $\dim G$ .

If  $H \subset \mathcal{Z}_G$ : Let  $T$  be a Maximal torus. For  $h \in H$ , for some  $g \in G$ ,

$$h \in gTg^{-1} \implies h = g^{-1}hg \in T \implies H \subset T$$

Also,  $\mathcal{Z}_G(H) = N_G(H) = G$ .

If  $H \not\subset \mathcal{Z}_G$ : let  $s \in H - \mathcal{Z}_G$ . Then  $H \subset Z := \mathcal{Z}_G(s) \neq G$  and so  $Z$  is connected by induction. Also by induction,  $s \in T$  for some Maximal torus  $T$ ; hence  $T \subset Z$ . We have injective morphisms

$$T \rightarrow Z/Z_u \rightarrow G/G_u \implies \dim T \leq \dim(Z/Z_u) \leq \dim(G/G_u)$$

But  $T$  is maximal, and so all of the dimensions must coincide:  $T$  is a Maximal torus of  $Z$ . By induction  $H \subset gTg^{-1}$  for some  $g \in Z$ , implying (i). Also,  $\mathcal{Z}_G(H) = \mathcal{Z}_Z(H)$  is connected by induction, giving (ii). For (iii), if  $n \in N_G(H), h \in H$ , then

$$[n, h] \in H \cap [G, G] \subset H \cap G_u = 1 \implies n \in \mathcal{Z}_G(H) \implies N_G(H) \subset \mathcal{Z}_G(H)$$

□

**Corollary 134.** *Let  $G$  be connected and solvable, and let  $T \subset G$  be a torus. Then*

$$T \text{ is maximal} \iff T \text{ is Maximal}$$

*Proof.* If  $T$  is Maximal and  $T \subset T'$  for some torus  $T'$ , then  $T \rightarrow T' \rightarrow G/G_u$  are injective morphisms, giving

$$\dim(G/G_u) = \dim T \leq \dim T' \leq \dim(G/G_u)$$

Hence,  $T = T'$  and  $T$  is maximal. If  $T$  is not Maximal, then  $T \subset T'$  for some Maximal  $T'$  by the above proposition, so  $T$  is not maximal. □

## 5.4 Cartan subgroups.

**Remark 135.** *From now on,  $G$  denotes a connected algebraic group.*

**Theorem 136.** *Any two maximal tori in  $G$  are conjugate.*

Let  $T, T'$  be maximal. Since both are connected and solvable they are each contained in Borels:  $T \subset B, T' \subset B'$ . There is a  $g \in G$  such that  $gBg^{-1} = B'$ .  $gTg^{-1}$  and  $T'$  are two maximal tori in  $B$  and so, by Proposition 130, for some  $b \in B, bgTg^{-1}b^{-1} = T'$ . □

**Corollary 137.** *A maximal torus in a Borel subgroup of  $G$  is a maximal torus in  $G$ .*

**Definition 138.** *A Cartan subgroup of  $G$  is  $\mathcal{Z}_G(T)^0$ , for a maximal torus  $T$ . All Cartan subgroups are conjugate. (We will see in Proposition 144 that  $\mathcal{Z}_G(T)$  is connected.)*

*Examples.*

- $G = \text{GL}_n, T = D_n, \mathcal{Z}_G(T) = T = D_n$
- $G = U_n, T = 1, \mathcal{Z}_G(T) = G = U_n$

**Proposition 139.** *Let  $T \subset G$  be a maximal torus.  $C := \mathcal{Z}_G(T)^0$  is nilpotent and  $T$  is its (unique) maximal torus.*

*Proof.*  $T \subset C$  and so  $T$  is a maximal torus of  $C$ . Moreover,  $T \subset \mathcal{Z}_G(C)$  and all maximal tori in  $C$  are conjugate, and so  $T$  is the unique maximal torus of  $C$ . Since any semisimple element lies in a maximal torus,

$$C_s = T \implies C/T \text{ unipotent} \implies C/T \text{ nilpotent} \implies C^n C \subset T \text{ for some } n \geq 0$$

. But  $T$  is central and so  $C^{n+1}C = [C, C^n C] \subset [C, T] = 1$ ; hence  $C$  is nilpotent. □

**Lemma 140.** *Let  $S \subset G$  be a torus. There exists  $s \in S$  such that  $\mathcal{Z}_G(S) = \mathcal{Z}_G(s)$ .*

*Proof.* Let  $G \hookrightarrow \mathrm{GL}_n$  be a closed immersion. Since  $S$  is a collection of commuting, diagonalisable elements, without loss of generality,  $S \hookrightarrow D_n$ . It is enough to show that  $\mathcal{Z}_{\mathrm{GL}_n}(S) = \mathcal{Z}_{\mathrm{GL}_n}(s)$ , for some  $s \in S$ . Let  $\chi_i \in X^*(D_n)$  be given by  $\mathrm{diag}(x_1, \dots, x_n) \mapsto x_i$ . It is easy to show that

$$\mathcal{Z}_G(S) = \{(x_{ij}) \in \mathrm{GL}_n \mid \forall i, j \quad x_{ij} = 0 \text{ if } \chi_i|_S \neq \chi_j|_S\}$$

The set

$$\bigcap_{\substack{i, j \\ \chi_i|_S \neq \chi_j|_S}} \{s \in S \mid \chi_i(s) \neq \chi_j(s)\}$$

is nonempty and open, and thus is dense; any  $s$  from the set will do.  $\square$

**Lemma 141.** *For a closed, connected subgroup  $H \subset G$ , let  $X = \bigcup_{x \in G} xHx^{-1} \subset G$ .*

- (i)  $X$  contains a nonempty open subset of  $\overline{X}$ .
- (ii)  $H$  parabolic  $\implies X$  closed
- (iii) If  $(N_G(H) : H) < \infty$  and there is  $y \in G$  lying in only finitely many conjugates of  $H$ , then  $\overline{X} = G$ .

*Proof.*

(i):

$$Y := \{(x, y) \mid x^{-1}yx \in H\} = \{(x, y) \mid y \in xHx^{-1}\} \subset G \times G$$

is a closed subset. Note that

$$\mathrm{pr}_2(Y) = \{y \in G \mid y \in xHx^{-1} \text{ for some } x\} = X$$

By Chevalley,  $X$  contains a nonempty open subset of  $\overline{X}$ .

(ii): Let  $P$  be parabolic.

$$\begin{array}{ccc} G \times G & \xrightarrow{\pi \times \mathrm{id}} & G/H \times G \\ & \searrow \mathrm{pr}_2 & \downarrow \mathrm{pr}'_2 \\ & & G \end{array}$$

Note that  $\pi \times \mathrm{id}$  is open (Corollary 87) and that

$$(x, y) \in Y \iff \forall h \in H \quad (xh, y) \in Y.$$

By the usual argument,  $(\pi \times \mathrm{id})(Y)$  is closed. Since  $G/P$  is complete,

$$\mathrm{pr}'_2((\pi \times \mathrm{id})(Y)) = \mathrm{pr}_2(Y) = X$$

is closed.

(iii): We have an isomorphism

$$Y \xrightarrow{\sim} G \times H, \quad (x, y) \mapsto (x, x^{-1}yx)$$

and so  $Y$  is irreducible (as  $H, G$  are connected). Consider the diagram

$$G \xleftarrow{\text{pr}_1} Y \xrightarrow{\text{pr}_2} G.$$

$$\begin{aligned} \text{pr}_1^{-1}(x) = \{(x, xhx^{-1}) \mid h \in H\} \cong H &\implies \text{all fibers of } \text{pr}_1 \text{ have dimension } \dim H \\ &\implies \dim Y = \dim G + \dim H \quad (\text{Theorem 85}). \end{aligned}$$

Moreover,

$$\text{pr}_2^{-1}(y) = \{(x, y) \mid y \in xHx^{-1}\} \cong \{x \mid y \in xHx^{-1}\}$$

Pick  $y \in G$  lying in finitely many conjugates of  $H$ :  $x_1Hx_1^{-1}, \dots, x_nHx_n^{-1}$ . Then

$$\text{pr}_2^{-1}(y) = \bigcup_{i=1}^n x_i N_G(H)$$

which is a finite union of  $H$  cosets by hypothesis ( $(N_G(H) : H) < \infty$ ). This implies that

$$\begin{aligned} \dim \text{pr}_2^{-1}(y) = \dim H &\implies \text{pr}_2 : Y \rightarrow \overline{\text{pr}_2(Y)} \text{ is a dominant map with minimal fibre dimension } \leq \dim H \\ &\implies \dim Y - \dim \overline{\text{pr}_2(Y)} \leq \dim H \\ &\implies \dim \overline{\text{pr}_2(Y)} \geq \dim Y - \dim H = \dim G \\ &\implies \overline{\text{pr}_2(Y)} = G \end{aligned}$$

□

**Theorem 142.**

- (i) *Every  $g \in G$  is contained in a Borel subgroup.*
- (ii) *Every  $s \in G_s$  is contained in a maximal torus.*

*Proof.*

(i): Pick a maximal torus  $T \subset G$ . Let  $C = \mathcal{Z}_G(T)^0$  be the associated Cartan subgroup. Because  $C$  is connected and nilpotent (Proposition 139), there is a Borel  $B \supset C$ .

$$\begin{aligned} T = C_s \quad (\text{Proposition 139}) &\implies N_G(C) = N_G(T) \quad (\text{“}\supset\text{” is obvious}) \\ &\implies (N_G(C) : C) = (N_G(T) : \mathcal{Z}_G(T)^0) < \infty \quad (\text{Corollary 53}) \end{aligned}$$

By Lemma 140 there is  $t \in T$  such that  $\mathcal{Z}_G(t)^0 = \mathcal{Z}_G(T)^0 = C$ .  $t$  is contained in a unique conjugate, i.e.,

$$t \in xCx^{-1} \implies xCx^{-1} = C$$

by the following.

$$\begin{aligned} t \in xCx^{-1} &\implies x^{-1}tx \in C, \quad \text{which is a semisimple element} \\ &\implies x^{-1}tx \in C_s = T \subset \mathcal{Z}_G(C) \\ &\implies C \subset \mathcal{Z}_G(x^{-1}tx)^0 = x^{-1}\mathcal{Z}_G(t)^0x = x^{-1}Cx \\ &\implies C = x^{-1}Cx \quad (\text{compare dimensions}) \end{aligned}$$

Hence, we can apply Lemma 141 (iii) with  $H = C$  to get

$$G = \overline{\bigcup_x xCx^{-1}} \subset \overline{\bigcup_x xBx^{-1}} = \bigcup_x xBx^{-1}$$

with the last equality following from Lemma 141 (ii) (this time with  $H = B$ ). Hence,  $G = \bigcup_x xBx^{-1}$ , giving (i) of the theorem.

(ii):

$$\begin{aligned} s \in G_s &\implies s \in B, \quad \text{for some Borel } B \text{ by (i)} \\ &\implies s \in T, \quad \text{for some maximal torus } T \text{ of } B \text{ by Theorem 130 (i).} \end{aligned}$$

(A maximal torus in  $B$  is a maximal torus in  $G$  by Theorem 136.)

**Corollary 143.** *If  $B \subset G$  is a Borel then  $\mathcal{Z}_B = \mathcal{Z}_G$ .*

*Proof.* The inclusion  $\mathcal{Z}_B \subset \mathcal{Z}_G$  follows Corollary 124. For the reverse inclusion, if  $z \in \mathcal{Z}_G$ , we have  $z \in gBg^{-1}$  for some  $g$  by the above Theorem, and so  $z = g^{-1}zg \in B$ .  $\square$

**Proposition 144.** *Let  $S \subset G$  be a torus.*

- (i)  $\mathcal{Z}_G(S)$  is connected.
- (ii) If  $B \subset G$  is a Borel containing  $S$ , then  $\mathcal{Z}_G(S) \cap B$  is a Borel in  $\mathcal{Z}_G(S)$ , and all Borels of  $\mathcal{Z}_G(S)$  arise this way.

*Proof.*

(i): Let  $g \in \mathcal{Z}_G(S)$  and  $B$  a Borel containing  $g$ . Define

$$X = \{xB \mid g \in xBx^{-1}\} \subset G/B$$

which is nonempty by Theorem 142. Consider the diagram

$$G/B \xleftarrow{\pi} G \xrightarrow{\alpha} G$$

in which  $\pi$  is the natural surjection and  $\alpha : x \mapsto x^{-1}gx$ . We have  $X = \pi(\alpha^{-1}(B))$ . Since  $\pi^{-1}(B)$  is a union of fibres of  $\pi$  and is closed, and  $\pi$  is open, we have that  $X$  is closed.  $X$  is thus complete, being a closed subset of the complete  $G/B$ .

$S$  acts on  $X \subset G/B$ , as for all  $s \in S$

$$xBx^{-1} \ni g \implies sxBx^{-1}s^{-1} \ni g \quad (\text{since } g = s^{-1}gs).$$

By the Borel Fixed Point Theorem (118),  $S$  has some fixed point  $xB \in X$ , so

$$sxB = xB \implies Sx \subset xB \implies S \subset xBx^{-1}.$$

Hence, since  $g$  also lies in  $xBx^{-1}$ , we have

$$g \in \mathcal{Z}_{xBx^{-1}}(S) \subset \mathcal{Z}_G(S)^0$$

where  $\mathcal{Z}_{xBx^{-1}}(S)$  is connected by Proposition 133. Thus,  $\mathcal{Z}_G(S) \subset \mathcal{Z}_G(S)^0$ : equality.

(ii): Let  $B$  be a Borel containing  $S$  and set  $Z = \mathcal{Z}_G(S)$ .  $Z \cap B = \mathcal{Z}_B(S)$  is connected by Proposition 133 and is also solvable. Therefore,  $Z \cap B$  is a Borel of  $Z$  if and only if it is parabolic, i.e., if  $Z/Z \cap B$  is complete. By the bijective map

$$Z/(Z \cap B) \rightarrow ZB/B$$

of homogeneous  $Z$ -spaces, we see that suffices to show that

$ZB/B \subset G/B$  is closed  $\iff Y := ZB \subset G$  is closed (by the definition of the quotient topology)

$Z$  being irreducible implies that

$$Y = \text{im}(Z \times B \xrightarrow{\text{mult}} G) \text{ is irreducible} \implies \bar{Y} \text{ irreducible.}$$

Let  $\pi : B \rightarrow B/B_u$  be the natural surjection and define

$$\phi : \bar{Y} \times S \rightarrow B/B_u, \quad (y, s) \mapsto \pi(y^{-1}sy).$$

(To make sure that this definition makes sense, i.e., that  $y^{-1}sy \in B$ , first check it when  $y \in Y = ZB$ .) For fixed  $y$ ,

$$\phi_y : S \rightarrow B/B_u, \quad s \mapsto \phi(y, s) = \pi(y^{-1}sy)$$

is a homomorphism. Therefore, by rigidity (Theorem 52), for all  $y \in Y$ ,  $\phi_e = \phi_y$ : for all  $s \in S$

$$\pi(y^{-1}sy) = \pi(s).$$

If  $T \supset S$  is a maximal torus, by the conjugacy of maximal tori in  $B$ , we have

$$uy^{-1}Syu^{-1} = T$$

for some  $u \in B_u$ . But then, by the above,

$$\pi(uy^{-1}uyu^{-1}) = \pi(y^{-1}sy) = \pi(s) \quad \text{for all } s \in S$$

while  $\pi|_T : T \rightarrow B/B_u$  is injective (an isomorphism even) (Jordan decomposition). Therefore,

$$uy^{-1}Syu^{-1} = S \implies yu^{-1} \in \mathcal{Z}_G(S) = Z \implies y \in ZB = Y$$

and thus  $Y$  is closed:  $Z \cap B \subset Z$  is Borel. Moreover, any other Borel of  $Z$  is

$$z(Z \cap B)z^{-1} = Z \cap (zBz^{-1}),$$

$zBz^{-1}$  containing  $S$ . □

### Corollary 145.

(i) *The Cartan subgroups are the  $\mathcal{Z}_G(T)$ , for maximal tori  $T$ .*

(ii) *If a Borel  $B$  contains a maximal torus  $T$ , then it contains  $\mathcal{Z}_G(T)$ .*

*Proof.*

(i) follows immediately from the above. For (ii), we have that  $\mathcal{Z}_G(T)$  is a Borel of  $\mathcal{Z}_G(T)$ . But  $\mathcal{Z}_G(T)$  is nilpotent (Proposition 139) and so  $\mathcal{Z}_G(T) \cap B = \mathcal{Z}_G(T)$ . □

## 5.5 Conjugacy of parabolic and Borel subgroups.

**Theorem 146.**

- (i) If  $B \subset G$  is Borel, then  $N_G(B) = B$ .
- (ii) If  $P \subset G$  is parabolic, then  $N_G(P) = P$  and  $P$  is connected.

*Proof.*

(i): Induct on the dimension of  $G$ . If  $G$  is solvable, then  $B = G$  and we are done; suppose otherwise. Let  $H = N_G(B)$  and  $x \in H$ . We want to show that  $x \in B$ . Pick a maximal torus  $T \subset B$ . Then  $xTx^{-1} \subset B$  is another maximal torus, and so  $T, xTx^{-1}$  are  $B$ -conjugate. Without loss of generality - changing  $x$  modulo  $B$  if necessary - suppose that  $T = xTx^{-1}$ . Consider

$$\phi : T \rightarrow T, \quad t \mapsto [x, t] = (xTx^{-1})t^{-1}.$$

Check that  $\phi$  is a homomorphism. (Use that  $T$  is commutative.)

Case 1.  $\text{im } \phi \neq T$ :

Let  $S = (\ker \phi)^0$ , which is a torus and is nontrivial since  $\text{im } \phi \neq T$ .  $x$  lies in  $Z = \mathcal{Z}_G(S)$  and normalises  $Z \cap B$  (which is a Borel of  $Z$  by Proposition 144). If  $Z \neq G$ , then  $x \in Z \cap B \subset B$  by induction. Otherwise, if  $Z = G$ , then  $S \subset \mathcal{Z}_G$  and  $B/S \subset G/S$  is a Borel by Corollary 123; hence,

$$[x] \text{ normalises } B/S \implies [x] \in B/S \text{ by induction} \implies x \in B.$$

Case 2.  $\text{im } \phi = T$ :

If  $\text{im } \phi = T$ , then

$$T \subset [x, T] \subset [H, H].$$

By Corollary 102, there is a  $G$ -representation  $V$  and a line  $kv \subset V$  such that  $H = \text{Stab}_G(kv)$ . Say  $hv = \chi(h)v$  for some character  $\chi : H \rightarrow \mathbf{G}_m$ .  $\chi(T) = \{e\}$  since  $T \subset [H, H]$  and  $\chi(B_u) = \{e\}$  by Jordan decomposition. Thus, as  $B = TB_u$  (Theorem 130),  $B$  fixes  $v$ . By the universal property of quotients, we have a morphism

$$G/B \rightarrow V, \quad gB \mapsto gv.$$

However, the image of the morphism must be a point, as  $V$  is affine, while  $G/B$  is complete and connected; hence,  $G$  fixes  $v$  and  $H = G$ , i.e.,  $B \trianglelefteq G$ . Therefore,  $G/B$  is affine, complete, and connected, and we must have  $G = B$ . (In particular,  $x \in B$ .)

(ii): By Theorem 122,  $P \supset B$  for some Borel  $B$  of  $G$ . Suppose  $n \in N_G(P)$ . Then  $nBn^{-1}, B$  are both contained in - and are Borels of -  $P^0$ . Therefore, there must be  $g \in P^0$  such that

$$nBn^{-1} = gBg^{-1} \implies g^{-1}n \in N_G(B) = B \text{ by (i)} \implies n \in gB \subset P^0.$$

Hence,

$$P \subset N_G(P) \subset P^0 \subset P.$$

□

**Proposition 147.** Fix a Borel  $B$ . Any parabolic subgroup is conjugate to a unique parabolic containing  $B$ .

**Remark 148.** For a fixed  $B$ , the parabolics containing  $B$  are called **standard parabolic subgroups**.

*Example.* If  $G = \mathrm{GL}_n$  and  $B = B_n$ , then the standard parabolic subgroups are the subgroups, for integers  $n_i \geq 1$  with  $n = \sum_i^m n_i$ , consisting of matrices

$$\begin{pmatrix} A_{n_1} & * & * & * \\ & A_{n_2} & * & * \\ & & \ddots & * \\ & & & A_{n_m} \end{pmatrix}$$

where  $A_{n_i} \in \mathrm{GL}_{n_i}$ .

*Proof of proposition.*

Let  $P$  be a parabolic.  $P$  contains some Borel  $gBg^{-1}$ , so  $B \subset g^{-1}Pg$ . This takes care of existence. For uniqueness, let  $P, Q \supset B$  be two conjugate parabolics; say,  $P = gQg^{-1}$ .

$$\begin{aligned} gBg^{-1}, B \subset Q \text{ Borels} &\implies g^{-1}Bg = qBq^{-1} \text{ for some } q \in Q \\ &\implies gq \in N_G(B) = B \\ &\implies g \in Bq^{-1} \subset Q \\ &\implies P = Q \end{aligned}$$

□

**Proposition 149.** If  $T$  is a maximal torus and  $B$  is a Borel containing  $T$ , then we have a bijection

$$\begin{aligned} N_G(T)/\mathcal{Z}_G(T) &\xrightarrow{\sim} \{\text{Borels containing } T\} \\ [n] &\mapsto nBn^{-1} \end{aligned}$$

**Exercise.** If  $G = \mathrm{GL}_n$ ,  $B = B_n$ , and  $T = D_n$ , we have that  $\mathcal{Z}_G(T) = T$ ,  $N_G(T) =$  permutation matrices, and that  $N_G(T)/\mathcal{Z}_G(T) \cong S_n$ . When  $n = 2$ , the two Borels containing  $T$  are  $\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}$

and  $\begin{pmatrix} * & 0 \\ * & * \end{pmatrix}$ .

*Proof of proposition.*

If  $B' \supset T$  is a Borel, then

$$\begin{aligned} B' = gBg^{-1} \text{ for some } g &\implies g^{-1}Tg, T \subset B \text{ are maximal tori} \\ &\implies g^{-1}Tg = bTb^{-1} \text{ for some } b \in B \\ &\implies n := gb \in N_G(T) \\ &\implies B' = gBg^{-1} = nBn^{-1}. \end{aligned}$$

Also,

$$nBn^{-1} = B \iff n \in N_G(B) \cap N_G(T) = B \cap N_G(T) = N_B(T) \stackrel{133}{=} \mathcal{Z}_B(T) \stackrel{145}{=} \mathcal{Z}_G(T).$$

□

**Remark 150.** Given a Borel  $B \subset G$ , we have a bijection

$$\begin{aligned} G/B &\xrightarrow{\sim} \{\text{Borels of } G\} \\ gB &\mapsto gBg^{-1} \end{aligned}$$

The projective variety  $G/B$  is called the **flag variety** of  $G$  (independent of  $B$  up to isomorphism).

*Example.* When  $G = \text{GL}_n$ ,  $B = B_n$

$$\begin{aligned} G/B &\xrightarrow{\sim} \{\text{flags } 0 \subsetneq V_1 \subsetneq V_2 \subsetneq \cdots \subsetneq V_n = k^n\} \\ gB &\mapsto g \left( 0 \subsetneq \begin{pmatrix} * \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \subsetneq \begin{pmatrix} * \\ * \\ 0 \\ \vdots \\ 0 \end{pmatrix} \subsetneq \cdots \subsetneq \begin{pmatrix} * \\ * \\ * \\ \vdots \\ * \end{pmatrix} = k^n \right) \end{aligned}$$

# 6. Reductive groups.

## 6.1 Semisimple and reductive groups.

**Definitions 151.** The radical  $RG$  of  $G$  is the unique maximal connected, closed, solvable, normal subgroup of  $G$ . Concretely,

$$RG = \left( \bigcap_{B \text{ Borel}} B \right)^0$$

(Recall that any two Borels are conjugate.) The **unipotent radical** of  $G$  is the unique maximal connected, closed, unipotent, normal subgroup of  $G$ :

$$R_u G = (RG)_u = \left( \bigcap_{B \text{ Borel}} B_u \right)^0$$

$G$  is **semisimple** if  $RG = 1$  and is **reductive** if  $R_u G = 1$ .

**Remarks 152.**

- $G$  semisimple  $\implies G$  reductive
- $G/RG$  is semisimple and  $G/R_u G$  is reductive. (Exercise!)
- If  $G$  is connected and solvable, then  $G = RG$  and  $G/R_u G = G/G_u$  is a torus. Hence a connected, solvable  $G$  is reductive  $\iff G$  is a torus.

*Example.*

- $\mathrm{GL}_n$  is reductive. Indeed,

$$R(\mathrm{GL}_n) \subset \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \cap \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} = D_n \implies R_u(\mathrm{GL}_n) = 1$$

Similarly,  $\mathrm{SL}_n$  is reductive.

- $\mathrm{GL}_n$  is not semisimple, as  $\{\mathrm{diag}(x, x, \dots, x) \mid x \in k^\times\} \trianglelefteq \mathrm{GL}_n$ .  $\mathrm{SL}_n$  is semisimple by Proposition 153 (iii) below.

**Proposition 153.**  $G$  is connected, reductive.

- $RG = Z_G^0$ , a central torus.
- $RG \cap \mathcal{D}G$  is finite.
- $\mathcal{D}G$  is semisimple.

**Remark 154.** In fact,  $RG \cdot \mathcal{D}G = G$ , so  $G = \mathcal{D}G$  when  $G$  is semisimple. Hence, by (ii) above,  $RG \times \mathcal{D}G \xrightarrow{\text{mult.}} G$  is surjective with finite kernel.

*Proof.*

(i).  $1 = R_u G = (RG)_u \implies RG$  is a torus, by Proposition 127. Hence, by rigidity (Corollary 53)  $N_G(RG)^0 = \mathcal{Z}_G(RG)^0$ . Moreover, since  $RG \trianglelefteq G$

$$G = N_G(RG)^0 = \mathcal{Z}_G(RG)^0 \implies G = \mathcal{Z}_G(RG) \implies RG \subset \mathcal{Z}_G^0$$

The reverse inclusion is clear.

(ii).  $S := RG$  is a torus. Embed  $G \hookrightarrow \mathrm{GL}(V)$ .  $V$  decomposes as  $V = \bigoplus_{\chi \in X(S)} V_\chi$ .

$$S \text{ is central} \implies G \text{ stabilises each } V_\chi \implies G \hookrightarrow \prod_x \mathrm{GL}(V_\chi)$$

It follows that  $\mathcal{D}G \hookrightarrow \prod_x \mathrm{SL}(V_\chi)$  and  $RG$  acts by scalars on each  $V_\chi$ . Since the scalars in  $\mathrm{SL}_n$  are given by the  $n$ -th roots of unity, the result follows.

(iii).

$$\begin{aligned} \mathcal{D}G \trianglelefteq G &\implies R(\mathcal{D}G) \subset RG \\ &\implies R(\mathcal{D}G) \subset RG \cap \mathcal{D}G, \text{ which is finite} \\ &\implies R(\mathcal{D}G) = 1 \end{aligned}$$

□

**Definition 155.** For a maximal torus  $T \subset G$ ,

$$I(T) := \left( \bigcap_{\substack{B \text{ Borel} \\ B \supset T}} B \right)^0$$

which is a connected, closed, solvable subgroup with maximal torus  $T$ :  $I(T) = I(T)_u \rtimes T$  (see Theorem 130).

*Claim:*

$$I(T)_u = \left( \bigcap_{B \supset T} B_u \right)^0$$

*Proof.*

“ $\subset$ ” : For all Borels  $B \supset T$

$$I(T) \subset B \implies I(T)_u \subset B_u \implies I(T)_u \subset \bigcap_{B \supset T} B_u \implies I(T)_u \subset \left( \bigcap_{B \supset T} B_u \right)^0$$

as  $I(T)_u$  is connected.

“ $\supset$ ” :  $\left( \bigcap_{B \supset T} B_u \right)^0 \subset I(T)$  and consists of unipotent elements. □

**Remark 156.**

$$I(T) \supset \left( \bigcap_B B \right)^0 = RG \implies I(T)_u \supset R_u G$$

In fact, the converse is true and equality holds.

**Theorem 157** (Chevalley).  $I(T)_u = R_u G$ . Hence,

$$G \text{ reductive} \iff I(T)_u = 1 \iff I(T) = T$$

**Corollary 158.** Let  $G$  be connected, reductive.

(i)  $S \subset G$  subtorus  $\implies \mathcal{Z}_G(S)$  connected, reductive.

(ii)  $T$  maximal torus  $\implies \mathcal{Z}_G(T) = T$ .

(iii)  $\mathcal{Z}_G$  is the intersection of all maximal tori. (In particular,  $\mathcal{Z}_G \subset T$  for all maximal tori  $T$ .)

*Proof of corollary.*

(i):  $\mathcal{Z}_G(S)$  is connected by Proposition 144. Let  $T \supset S$  be a maximal torus, so that  $T \subset \mathcal{Z}_G(S) =: Z$ . Again by Proposition 144

$$\begin{aligned} \{ \text{Borels of } Z \text{ containing } T \} &= \{ Z \cap B \mid B \supset T \text{ Borel of } G \} \\ \implies I_Z(T) &= \left( \bigcap_{B \supset T} (Z \cap B) \right)^0 \subset I(T) \stackrel{157}{=} T \\ \implies I_Z(T) &= T \\ \implies Z &\text{ is reductive, by the theorem} \end{aligned}$$

(ii):  $\mathcal{Z}_G(T)$  is reductive by (i) and solvable (as it is a Cartan subgroup, which is nilpotent by Proposition 139). Hence,  $\mathcal{Z}_G(T)$  is a torus:  $T = \mathcal{Z}_G(T)$ , by maximality, since  $T \subset \mathcal{Z}_G(T)$ .

(iii):  $T$  maximal  $\implies T = \mathcal{Z}_G(T) \supset \mathcal{Z}_G$ . For the converse, let  $H = \bigcap_{T \text{ max.}} T$ , which is a closed, normal subgroup of  $G$  (normal because all maximal tori are conjugate). Since  $H$  is commutative and  $H = H_s$ ,  $H$  is diagonalisable, and by Corollary 53

$$G = N_G(H)^0 = \mathcal{Z}_G(H)^0 \implies G = \mathcal{Z}_G(H) \implies H \subset \mathcal{Z}_G$$

□

We will now build up several results in order to prove Theorem 157, following D. Luna's proof from 1999<sup>1</sup>.

**Proposition 159.** Suppose  $V$  is a  $\mathbf{G}_m$ -representation.  $\mathbf{G}_m$  acts on  $\mathbf{P}V$ . If  $v \in V - \{0\}$ , write  $[v]$  for its image in  $\mathbf{P}V$ . Then either,  $\mathbf{G}_m \cdot [v] = [v]$ , i.e.,  $v$  is a  $\mathbf{G}_m$ -eigenvector, or  $\overline{\mathbf{G}_m \cdot [v]}$  contains two distinct  $\mathbf{G}_m$ -fixed points.

Precise version of the proposition: Write  $V = \bigoplus_{n \in \mathbf{Z} = X^*(\mathbf{G}_m)} V_n$ , where

$$V_n = \{v \in V \mid t \cdot v = t^n v \quad \forall t \in \mathbf{G}_m\}, \text{ i.e., "v has weight n"}$$

For  $v \in V$ , write  $v = \sum_{n \in \mathbf{Z}} v_n$  with  $v_n \in V_n$ . Then

$$[v_r], [v_s] \in \overline{\mathbf{G}_m \cdot [v]}$$

where  $r = \min\{n \mid v_n \neq 0\}$  and  $s = \max\{n \mid v_n \neq 0\}$ . Clearly,  $[v_r], [v_s]$  are  $\mathbf{G}_m$ -fixed. In fact, if  $\mathbf{G}_m \cdot [v] \neq [v]$ , then

$$\overline{\mathbf{G}_m \cdot [v]} = (\mathbf{G}_m \cdot [v]) \sqcup \{[v_r]\} \sqcup \{[v_s]\}$$

<sup>1</sup>See for example P. Polo's M2 course notes (§21 in Séance 5/12/06) at [www.math.jussieu.fr/~polo/M2](http://www.math.jussieu.fr/~polo/M2)

*Proof.* Pick a basis  $e_0, e_1, \dots, e_n$  of  $V$  such that  $e_i \in V_{m_i}$ . Without loss of generality  $m_0 \leq m_1 \leq \dots \leq m_n$ . Write  $v = \sum_i \lambda_i e_i$ ,  $\lambda_i \in k$ . The orbit map  $f : \mathbf{G}_m \rightarrow \mathbf{P}V$  is given by mapping  $t$  to

$$t \cdot [v] = (t^{m_0} \lambda_0 : t^{m_1} \lambda_1 : \dots : t^{m_n} \lambda_n) = (0 : \dots : 0 : \lambda_u : \dots : t^{m_i-r} \lambda_i : \dots : t^{s-r} \lambda_v : 0 : \dots : 0)$$

where  $u = \min\{i \mid \lambda_i \neq 0\}$  and  $v = \max\{i \mid \lambda_i \neq 0\}$ , so that  $m_u = r$  and  $m_v = s$ .

Define  $\tilde{f} : \mathbf{P}^1 \rightarrow \mathbf{P}V$  by

$$(T_0 : T_1) \mapsto (0 : \dots : 0 : T_1^{s-r} \lambda_u : \dots : T_0^{m_i-r} T_1^{s-m_i} \lambda_i : \dots : T_0^{s-r} \lambda_v : 0 : \dots : 0)$$

Check that this is a morphism and that  $\tilde{f}|_{\mathbf{G}_m} = f$ . (In fact,  $\tilde{f}$  is the unique extension of  $f$ , since  $\mathbf{P}V$  is separated and  $\mathbf{G}_m$  is dense.) We have

$$\tilde{f}(\mathbf{P}^1) = \tilde{f}(\overline{\mathbf{G}_m}) \subset \overline{\tilde{f}(\mathbf{G}_m)} = \overline{\mathbf{G}_m \cdot [v]}$$

and

$$\tilde{f}(0 : 1) = (0 : \dots : \lambda_u : \dots : 0 : \dots : 0) = [v_r] \quad \text{and} \quad \tilde{f}(1 : 0) = \dots = [v_s]$$

(In fact, we actually have  $\tilde{f}(\mathbf{P}^1) = \overline{\mathbf{G}_m \cdot [v]}$ , using the fact that  $\mathbf{P}^1$  is complete).  $\square$

Informally, above, we have

$$\begin{aligned} [v_r] &= \lim_{t \rightarrow 0} t \cdot [v] \in (\mathbf{P}V)^{\mathbf{G}_m} \\ [v_s] &= \lim_{t \rightarrow \infty} t \cdot [v] \in (\mathbf{P}V)^{\mathbf{G}_m} \end{aligned}$$

**Lemma 160.** *Let  $M$  be a free abelian group, and  $M_1, \dots, M_r \subsetneq M$  subgroups such that each  $M/M_i$  is torsion-free. Then*

$$M \neq M_1 \cup \dots \cup M_r$$

*Proof.* Since  $M/M_i$  is torsion-free, it is free abelian, and

$$0 \rightarrow M_i \rightarrow M \rightarrow M/M_i \rightarrow 0$$

splits, giving that  $M_i$  is a (proper) direct summand of  $M$ . Thus,  $M_i \otimes \mathbf{C} \subsetneq M \otimes \mathbf{C}$ ; hence

$$M \otimes \mathbf{C} \neq \bigcup_{i=1}^r M_i \otimes \mathbf{C}$$

as the former is irreducible and the latter are proper closed subsets.  $\square$

**Lemma 161.** *Let  $T$  be a torus and  $V$  an algebraic representation of  $T$ , so that  $T$  acts on  $\mathbf{P}V$ . Then, there is a cocharacter  $\lambda : \mathbf{G}_m \rightarrow T$  such that  $(\mathbf{P}V)^T = (\mathbf{P}V)^{\lambda(\mathbf{G}_m)}$ .*

*Proof.* Let  $\chi_1, \dots, \chi_r \in X^*(T)$  be distinct such that  $V = \bigoplus_{i=1}^r V_{\chi_i}$  and  $V_{\chi_i} \neq 0$  for all  $i$ . Then

$$[v] \in (\mathbf{P}V)^T \iff v \in V_{\chi_i} \quad \text{for some } i$$

So it is enough to show that there is a cocharacter  $\lambda$  such that

$$\forall i \neq j \quad \chi_i \circ \lambda \neq \chi_j \circ \lambda \iff (\chi_i - \chi_j) \circ \lambda \neq 0$$

Recall from Proposition 33 we have that

$$X^*(T) \times X_*(T) \rightarrow X^*(\mathbf{G}_m) \cong \mathbf{Z}, \quad (\chi, \lambda) \mapsto \chi \circ \lambda$$

is a perfect pairing.

Let  $M = X_*(T)$ , which is free abelian, and for all  $i \neq j$

$$M_{ij} := \{\lambda \in X_*(T) \mid \langle \chi_i - \chi_j, \lambda \rangle = 0\} \neq M \quad (\text{as } \chi_i \neq \chi_j)$$

For  $n > 0$ , if  $n\lambda \in M_{ij}$ , then  $\lambda \in M_{ij}$ , and so  $M/M_{ij}$  is torsion-free. By the above lemma,  $M \neq \bigcup_{i \neq j} M_{ij}$ , so there is a  $\lambda \in M$  such that

$$\forall i \neq j \quad 0 \neq \langle \chi_i - \chi_j, \lambda \rangle = (\chi_i - \chi_j) \circ \lambda$$

□

**Theorem 162** (Konstant-Rosenlicht). *Suppose that  $G$  is unipotent and  $X$  is an affine  $G$ -space. Then all orbits are closed.*

*Proof.* Let  $Y \subset X$  be an orbit. Without loss of generality, we replace  $X$  by  $\overline{Y}$  (which is affine). Since  $Y$  is locally closed and dense, it is open. Let  $Z = X - Y$ , which is closed.  $G$  acts (locally-algebraic) on  $k[X]$ , preserving  $I_X(Z) \subset k[X]$ .  $I_X(Z) \neq 0$ , as  $Z \neq X$ . By Theorem 39, since  $G$  is unipotent, it has a nonzero fixed point, say,  $f$  in  $I_X(Z)$ .  $f$  is  $G$ -invariant and hence is constant on  $Y$ . But then

$$Y \text{ is dense} \implies f \text{ is constant } (\neq 0) \implies k[X] = I_X(Z) \implies Z = \emptyset \implies Y = X \text{ is closed}$$

□

Now, we want to prove Theorem 157. Fix a Borel  $B \subset G$  and set  $X = G/B$ , a homogeneous  $G$ -space. Note that

$$X^T = \{gB \mid Tg \subset gB \iff T \subset gBg^{-1}\} \leftrightarrow \{\text{Borel subgroups containing } T\}$$

Furthermore, by Proposition 149,  $X^T$  is in bijection with  $N_G(T)/\mathcal{Z}_G(T)$  and hence is finite. Thus  $N_G(T)/\mathcal{Z}_G(T)$  acts simply transitively on  $X^T$ . For  $p \in X^T$ , define

$$X(p) = \{x \in X \mid p \in \overline{Tx}\}$$

**Proposition 163** (Luna). *For  $p \in X^T$ ,  $X(p)$  is open (in  $X$ ), affine, and  $I(T)$ -stable.*

*Proof.* By Corollary 102 there exists a  $G$ -representation  $V$  and a line  $L \subset V$  such that  $B = \text{Stab}_G(L)$  and  $\text{Lie } B = \text{Stab}_{\mathfrak{g}}(L)$ . This gives a map of  $G$ -spaces

$$i : X = G/B \rightarrow \mathbf{P}V, \quad g \mapsto gL.$$

$i$  and  $di$  are injective (Corollary 103); hence,  $i$  is a closed immersion (Corollary 103). Without loss of generality,  $X \subset \mathbf{P}V$  is a closed  $G$ -stable subvariety - and, replacing  $V$  by the  $G$ -stable  $\langle G \cdot L \rangle$ ,

we may also suppose that  $X$  is not contained in any  $\mathbf{P}V' \subset \mathbf{P}V$  for any subspace  $V' \subset V$ .

By Lemma 161, there is a cocharacter  $\lambda : \mathbf{G}_m \rightarrow T$  such that  $X^T = X^{\mathbf{G}_m}$ , considering  $X$  and  $\mathbf{P}V$  as  $\mathbf{G}_m$ -spaces via  $\lambda$ . For  $p \in X^T$ , write  $p = [v_p]$  for some  $v_p \in V_{m(p)}$ ,  $m(p) \in \mathbf{Z}$  (weight). Pick  $p_0 \in X^T$  such that  $m_0 := m(p_0)$  is minimal. Set  $e_0 = v_{p_0}$  and extend  $e_0$  to a basis  $e_0, e_1, \dots, e_n$  of  $V$  such that  $\lambda(t)e_i = t^{m_i}e_i$ . Without loss of generality,  $m_1 \leq \dots \leq m_n$ . Let  $e_0^*, \dots, e_n^* \in V^*$  denote the dual basis.

Claim 1.  $m_0 < m_1$ :

Suppose that  $m_0 > m_1$ . There is  $[v] \in X$  such that  $e_1^*(v) \neq 0$  (otherwise  $X \subset \mathbf{P}(\ker e_1^*) \subsetneq \mathbf{P}V$ ). Then, by Proposition 159,

$$[v_{m_1}] = \lim_{t \rightarrow 0} \lambda(t)[v] \in (\mathbf{P}V)^{\mathbf{G}_m} \cap X = X^T$$

(with the inclusion following from the fact that  $X$  is complete). This contradicts the minimality of  $m_0$ , so we must have  $m_0 \leq m_1$ .

Suppose that  $m_0 = m_1$ . Define

$$Z = \{z \in k \mid \text{there is some point of the form } (1 : z : \dots) \text{ in } X\}$$

If  $(1 : z : \dots) \in X$ , then by Proposition 159, as  $m_0 = m_1$ ,

$$(1 : z : \dots)' = \lim_{t \rightarrow 0} \lambda(t)(1 : z : \dots) \in X^T.$$

Since  $X^T$  is finite, so too is  $Z$ . Writing  $Z = \{z_1, \dots, z_r\}$ , we have

$$X \subset \mathbf{P}(\ker e_0^*) \cup \bigcup_{i=1}^r \mathbf{P}(\ker(e_1^* - z_i e_0^*)).$$

Since  $X$  is irreducible, it is contained in one of these subspaces, which is a contradiction.

Therefore,  $m_0 < m_1$ .

Claim 2.  $X(\lambda, p_0) := \{x \in X \mid e_0^*(x) \neq 0\}$  is open in  $X$ , affine, and  $T$ -stable. Also,  $X(\lambda, p_0) = X(p_0)$ , and it is  $I(T)$ -stable:

$X(\lambda, p_0) = X \cap (e_0^* \neq 0)$  is open in  $X$  and affine (as  $(e_0^* \neq 0)$  is open and affine in  $\mathbf{P}V$ ). It is  $T$ -stable, as  $e_0^*$  is an eigenvector for  $T$  (as  $e_0$  is an eigenvector for  $T$ ).

If  $x \in X(\lambda, p_0)$ , as  $m_0 < m_i$  for all  $i \neq 0$  (Claim 1),

$$\lim_{t \rightarrow 0} \lambda(t)x = [e_0] = p_0.$$

Hence,  $p_0 \in \overline{\mathbf{G}_m \cdot x} \subset \overline{T x}$ , so  $x \in X(p_0)$ . Let  $x \in X(p_0)$  and suppose that  $e_0^*(x) = 0$ . Then

$$p_0 \in \overline{T x} \subset X - X(\lambda, p_0)$$

with  $X - X(\lambda, p_0)$   $T$ -stable and closed. This is a contradiction and so we must have  $x \in X(\lambda, p_0)$ . Hence,  $X(\lambda, p_0) = X(p_0)$ .

To show that the set is  $I(T)$ -stable, we need to show that from the of  $G$  on  $\mathbf{P}(V^*)$  (which arises from the action on  $V^*$ ), we have

$$e_0^\perp = \{\ell \in V^* \mid \langle \ell, e_0 \rangle = 0\}$$

First, let us adress a third claim.

**Claim 3.** (i) Each  $G$ -orbit in  $\mathbf{P}(V^*)$  intersects the open subset  $\mathbf{P}(V^*) - \mathbf{P}(e_0^\perp)$  and (ii)  $G \cdot [e_0^*]$  is closed in  $\mathbf{P}(V^*)$ : (i): Pick  $v \in V^* - \{0\}$ . If  $G\ell \subset e_0^\perp$ , then for all  $g \in G$

$$0 = \langle g\ell, e_0 \rangle = \langle \ell, g^{-1}e_0 \rangle.$$

But  $Ge_0$  spans  $V$  (otherwise,  $X = Ge_0 \subset \mathbf{P}(V') \subsetneq \mathbf{P}V$ , which is a contradiction) and so

$$\langle \ell, V \rangle = 0 \implies \ell = 0$$

which is another contradiction. Hence,  $G[\ell] \not\subset \mathbf{P}(e_0^\perp)$ .

(ii):  $e_i^*$  has weight  $-m_i$  under the  $\mathbf{G}_m$ -action and

$$-m_n \leq \dots \leq -m_1 < -m_0.$$

Hence by Proposition 159, if  $x \in \mathbf{P}(V^*) - \mathbf{P}(e_0^\perp)$  then  $[e_0^*] \in \overline{\mathbf{G}_m \cdot x}$ . So, for all  $x \in \mathbf{P}(V^*)$ , by (i),

$$[e_0^*] \in \overline{Gx} \implies G[e_0^*] \subset \overline{Gx}.$$

If  $Gx$  is a closed orbit (which exists), we deduce that it is equal to  $G[e_0^*]$ .

Let us return to Claim 2, that  $X(\lambda, p_0)$  is  $I(T)$ -stable. Recall that  $I(T) = \left( \bigcap_{B' \supset T} B' \right)^0$ . Define  $P = \text{Stab}_G([e_0^*])$ . Since  $G/P \rightarrow G[e_0^*]$  is bijective map of  $G$ -spaces and the latter space is complete (Claim 3), it follows that  $P$  is parabolic. Hence, there is a parabolic  $B'$  of  $G$  contained in  $P$ . Moreover, since  $e_0^*$  is a  $T$ -eigenvector,  $T \subset P$ . There is a maximal torus of  $B'$  conjugate to  $T$  in  $P$ , so without loss of generality suppose that  $T \subset B' \subset P$ . It follows that  $I(T) (\subset B')$  stabilises  $[e_0^*]$  and hence also stabilises the set

$$X(\lambda, p_0) = \{x \in X \mid e_0^*(x) \neq 0\},$$

completing claim 2.

Now,  $N_G(T)$  acts transitively on  $X^T$  by above. If  $p \in X^T$ , then  $p = np_0$  for some  $n \in N_G(T)$ ; hence  $X(p) = nX(p_0)$  is open, affine, and stable under  $nI(T)n^{-1} = I(T)$  (equality following from the fact that  $n$  permutes the Borels containing  $T$ ).  $\square$

**Corollary 164.**  $\dim X \leq 1 + \dim(X - X(p_0))$

*Proof.* Either  $X = X(p_0)$  or otherwise. If equality holds, then  $X$  is complete, affine, and connected, and is thus a point. In this case,  $\dim X = 0$  and the inequality is true. Suppose that  $X \neq X(p_0)$  ( $= X(\lambda, p_0)$ ). Pick  $y \in X - X(\lambda, p_0)$ . Then  $e_0^*(y) = 0$ , and  $e_i^*(y) \neq 0$  for some  $i > 0$ . Let

$$U = \{x \in X \mid e_i^*(x) \neq 0\} \subset X,$$

which is nonempty and open. Define the morphism

$$f : U \rightarrow \mathbf{A}^1, \quad x \mapsto \frac{e_0^*(x)}{e_i^*(x)}$$

$f^{-1}(0) \subset X - X(\lambda, p_0)$ . By Corollary 87,

$$\dim(X - X(\lambda, p_0)) \geq \dim U - \dim \overline{f(U)} \geq \dim U - 1 = \dim X - 1$$

□

**Proposition 165** (Luna).  $I(T)_u$  acts trivially on  $X = G/B$ .

*Proof.*  $J := I(T)_u$ . If  $x \in X$ , then  $\overline{T x}$  contains a  $T$ -fixed point by the Borel Fixed Point Theorem; hence

$$X = \bigcup_{x \in X^T} X(p).$$

Fix  $x \in X$ .  $J$  being connected, solvable implies that  $\overline{J x}$  contains a  $J$ -fixed point  $y$ . By the above, we see that  $y \in X(p)$  for some  $p \in X^T$ . If

$$J x \cap (X - X(p)) \neq \emptyset,$$

with  $X - X(p)$  closed and  $J$ -stable by Proposition 163, then

$$y \in \overline{J x} \subset X - X(p)$$

which is a contradiction. Hence,  $J x \subset X(p)$ ,  $X(p)$  being affine by Proposition 163, and  $J$  being unipotent implies that  $J x \subset X(p)$  is closed by Konstant-Rosenlicht (162). But

$$\begin{aligned} y \in X(p) \cap \overline{J x} = J x \quad (J x \text{ is closed}) &\implies J x = J y = y, \quad \text{as } y \text{ is } J\text{-fixed} \\ &\implies x = y \text{ is } J\text{-fixed} \\ &\implies J \text{ acts trivially on } X. \end{aligned}$$

□

*Proof of Theorem 157.*

Let  $J = I(T)_u$  again. We want to show that  $J = R_u G$  and we already know that  $J \supset R_u G$ . For the reverse inclusion, we have that for all  $g \in G$ ,

$$\begin{aligned} J(gB) = gB \quad (\text{Theorem 165}) &\implies Jg \subset gB \\ &\implies J \subset gBg^{-1} \\ &\implies J \subset (gBg^{-1})_u, \quad \text{as } J \text{ is unipotent} \\ &\implies J \subset \left( \bigcap_g (gBg^{-1})_u \right)^0 = R_u G, \quad \text{as } J \text{ is connected} \end{aligned}$$

□

## 6.2 Overview of the rest.

**Plan for the rest of the course:** Given connected, reductive  $G$  (and a maximal torus  $T$ ) we want to show the following:

- $\mathfrak{g} = \text{Lie } T \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$ , under the adjoint action of  $T$ , where  $\Phi \subset X^*(T)$  is finite.
- There is a natural bijection  $\Phi \xrightarrow{\sim} \Phi^\vee$ , where  $\Phi^\vee \subset X_*(T)$  is such that  $(X^*(T), \Phi, X_*(T), \Phi^\vee)$  is a root datum (to be defined shortly).
- For all  $\alpha \in \Phi$ , there is a unique closed subgroup  $U_\alpha \subset G$ , normalised by  $T$ , such that  $\text{Lie } U_\alpha = \mathfrak{g}_\alpha$ .
- $G = \langle T \cup \bigcup_{\alpha \in \Phi} U_\alpha \rangle$ .

From now on  $G$  denotes a connected, reductive algebraic group. Fix a maximal torus  $T$ , so that

$$\mathfrak{g} = \bigoplus_{\lambda \in X^*(T)} \mathfrak{g}_\lambda$$

for the adjoint  $T$ -action. We write  $X^*(T)$  additively, so

$$\mathfrak{g}_0 = \{X \in \mathfrak{g} \mid \text{Ad}(t)X = X \text{ for all } t \in T\} = \mathfrak{z}_{\mathfrak{g}}(T) \stackrel{98}{=} \text{Lie } \mathcal{Z}_G(T) \stackrel{158}{=} \text{Lie } T = \mathfrak{t}$$

Define  $\Phi = \Phi(G, T) := \{\alpha \in X^*(T) - \{0\} \mid \mathfrak{g}_\alpha \neq 0\}$ , which is finite. The  $\alpha \in \Phi$  are the **roots** of  $G$  (with respect to  $T$ ). Hence,

$$\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$$

**Definition 166.** *The Weyl group of  $(G, T)$  is*

$$W = W(G, T) := N_G(T) / \mathcal{Z}_G(T) \stackrel{158}{=} N_G(T) / T$$

*which is finite by Corollary 53.  $W$  acts faithfully on  $T$  by conjugation, and hence acts on  $X^*(T)$  and  $X_*(T)$ :*

$$w \in W \mapsto \begin{cases} (w^{-1})^* : X^*(T) \rightarrow X^*(T) \\ w_* : X_*(T) \rightarrow X_*(T) \end{cases}$$

Explicitly,

$$\begin{aligned} w\mu &= \mu(\dot{w}^{-1}(\cdot)\dot{w}), & \text{for } \mu \in X^*(T) \\ w\lambda &= \dot{w}\lambda(\cdot)\dot{w}^{-1}, & \text{for } \lambda \in X_*(T) \end{aligned}$$

where  $\dot{w} \in N_G(T)$  lifts  $w$ .

**Remarks 167.**

- *The natural perfect pairing  $X^*(T) \times X_*(T) \rightarrow \mathbf{Z}$  is  $W$ -invariant:  $\langle w\mu, w\lambda \rangle = \langle \mu, \lambda \rangle$ .*
- *$W$  preserves  $\Phi \subset X^*(T)$  because  $N_G(T)$  permutes the eigenspaces  $\mathfrak{g}_\alpha$ . (Check that  $\text{Ad}(\dot{w})\mathfrak{g}_\alpha = \mathfrak{g}_{w\alpha}$ .)*

*Example.*  $G = \text{GL}_n$ ,  $T = D_n$ .

$\mathfrak{g} = M_n(k)$  and  $T$  acts by conjugation.

$$\mathfrak{g} = \begin{pmatrix} * & & & \\ & * & & \\ & & \ddots & \\ & & & * \end{pmatrix} \oplus \bigoplus_{\substack{i,j \\ i \neq j}} \begin{pmatrix} * & & & \\ & & & \\ & & & \\ & & & \end{pmatrix}$$

where in the summands on the right  $*$  appears in the  $(i, j)$ -th entry. On the  $(i, j)$ -th summand,  $\text{diag}(x_1, \dots, x_n) \in T$  acts as multiplication by  $x_i x_j^{-1}$ . Letting  $\epsilon_i \in X^*(T)$  denote  $\text{diag}(x_1, \dots, x_n) \mapsto x_i$ , we get that  $\Phi = \{\epsilon_i - \epsilon_j \mid i \neq j\}$ . Also,  $W = N_G(T)/T \cong S_n$  acts by permuting the  $\epsilon_i$ .

**Lemma 168.** *If  $\phi : H \rightarrow H'$  is a surjective morphism of algebraic groups and  $T \subset H$  is a maximal torus, then  $\phi(T) \subset H'$  is a maximal torus.*

*Proof.* Pick a Borel  $B \supset T$ , so that  $B = B_u \rtimes T$  and  $\phi(B) = \phi(B_u)\phi(T)$ .  $\phi(B)$  is a Borel of  $H'$  by Corollary 123.  $\phi(T)$  is a torus, as it is connected, commutative, and consists of semisimple elements.  $\phi(B_u) \subset \phi(B)_u$  is unipotent (Jordan decomposition). Finally,

$$\begin{aligned} \phi(T) \rightarrow \phi(B)/\phi(B)_u \text{ bijective (Jordan decomposition)} &\implies \dim \phi(T) = \dim \phi(B) - \dim \phi(B)_u \\ &\implies \phi(T) \subset \phi(B) \text{ maximal torus} \\ &\implies \phi(T) \subset H' \text{ maximal torus} \end{aligned}$$

□

**Lemma 169.** *If  $S \subset T$  be a subtorus, then*

$$\mathcal{Z}_G(S) \supseteq T \iff S \subset (\ker \alpha)^0 \text{ for some } \alpha \in \Phi$$

*Proof.* We already have  $\mathcal{Z}_G(S) \supset T$ . Note that

$$\text{Lie } \mathcal{Z}_G(S) \stackrel{98}{=} \mathfrak{z}_{\mathfrak{g}}(S) = \{X \in \mathfrak{g} \mid \text{Ad}(s)(X) = X \text{ for all } s \in S\} = \mathfrak{t} \oplus \bigoplus_{\substack{\alpha \in \Phi \\ \alpha|_S=1}} \mathfrak{g}_{\alpha}$$

$$\begin{aligned} \text{“} \supseteq \text{”} &\iff \text{Lie } \mathcal{Z}_G(S) \supseteq \mathfrak{t}, \text{ by dimension considerations} \\ &\iff \mathfrak{t} \oplus \bigoplus_{\substack{\alpha \in \Phi \\ \alpha|_S=1}} \mathfrak{g}_{\alpha} \supseteq \mathfrak{t} \\ &\iff S \subset \ker \alpha, \text{ for some } \alpha \in \Phi \end{aligned}$$

□

For  $\alpha \in \Phi$ , define  $T_{\alpha} := (\ker \alpha)^0$ , which is a torus of dimension  $\dim T - 1$ , as  $\text{im } \alpha = \mathbf{G}_m$ . Define  $G_{\alpha} := \mathcal{Z}_G(T_{\alpha})$ , which is connected, reductive by Corollary 158. Note that

$$T_{\alpha} \subset \mathcal{Z}_{G_{\alpha}}^0 \stackrel{153}{=} R(G_{\alpha})$$

Let  $\pi$  denote the natural surjection  $G_{\alpha} \rightarrow G_{\alpha}/R(G_{\alpha})$ . By Lemma 168,  $\pi(T)$  is a maximal torus of  $G_{\alpha}/R(G_{\alpha})$ .

$$T_{\alpha} \subset R(G_{\alpha}) \implies T/T_{\alpha} \twoheadrightarrow \pi(T) \implies \dim \pi(T) \leq 1$$

If  $\dim \pi(T) = 0$ , then

$$T \subset R(G_{\alpha}) \subset \mathcal{Z}_{G_{\alpha}} \implies G_{\alpha} \subset \mathcal{Z}_G(T) = T$$

which is a contradiction by Lemma 169. Hence,  $\dim \pi(T) = 1$ .

**Definitions 170.**

the **rank** of  $G = \text{rk } G := \dim T$ , where  $T$  is a maximal torus  
the **semisimple rank** of  $G = \text{ss-rk } G := \text{rk}(G/RG)$

Hence,  $\text{ss-rk } G_\alpha = 1$ . Note that since all maximal tori are conjugate, rank is well-defined, and that  $\text{ss-rk } G \leq \text{rk } G$  by Lemma 168.

*Example.*  $G = \text{GL}_n$ ,  $\alpha = \epsilon_i - \epsilon_{i+1}$ . We have

$$T_\alpha = \{\text{diag}(x_1, \dots, x_n) \mid x_i = x_{i+1}\}$$

and

$$G_\alpha = D_{i-1} \times \text{GL}_2 \times D_{n-i-1}.$$

$G_\alpha/RG_\alpha \cong \text{PGL}_2$  and  $\mathcal{D}G_\alpha \cong \text{SL}_2$ .

### 6.3 Reductive groups of rank 1.

**Proposition 171.** *Suppose that  $G$  is not solvable and  $\text{rk } G = 1$ . Pick a maximal torus  $T$  and a Borel  $B$  containing  $T$ . Let  $U = B_u$ .*

- (i)  $\#W = 2$ ,  $\dim G/B = 1$ , and  $G = B \sqcup UnB$ , where  $n \in N_G(T) - T$ .
- (ii)  $\dim G = 3$  and  $G = \mathcal{D}G$  is semisimple.
- (iii)  $\Phi = \{\alpha, -\alpha\}$  for some  $\alpha \neq 0$ , and  $\dim \mathfrak{g}_{\pm\alpha} = 1$ .
- (iv)  $\psi : U \times B \rightarrow UnB$ ,  $(u, b) \mapsto unb$ , is an isomorphism of varieties.
- (v)  $G \cong \text{SL}_2$  or  $\text{PSL}_2$

**Remark 172.** *In either case,  $G/B \cong \mathbf{P}^1$ . For example,*

$$\text{SL}_2 / \begin{pmatrix} * & * \\ & * \end{pmatrix} \xrightarrow{\sim} \mathbf{P}^1, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto (a : c)$$

*Proof of proposition.*

(i):

$$W \hookrightarrow \text{Aut}(X^*(T)) \cong \text{Aut}(\mathbf{Z}) = \{\pm 1\} \implies \#W \leq 2$$

If  $W = 1$ , then  $B$  is the only Borel containing  $T$ , and so by Theorem 157

$$B = I(T) = T \implies B \text{ nilpotent} \xrightarrow{125} G \text{ solvable}$$

which contradicts our hypothesis; hence,  $\#W = 2$ .

Set  $X := G/B$ .  $\dim X > 0$  since  $B \neq G$ . By Proposition 149 we have  $\#X^T = \#W = 2$ . By Corollary 164

$$\dim X \leq 1 + \dim(X - X(p_0))$$

Since  $X - X(p_0)$  is  $T$ -stable and closed (Proposition 163), it can contain at most one  $T$ -fixed point (as  $\#X^T = 2, p_0 \in X(p_0)$ ). By Proposition 159,  $T$  acts trivially and so  $X - X(p_0)$  is finite:

$$\dim X \leq 1.$$

Now,

$$\begin{aligned} \#W = 2 &\implies B, nBn^{-1} \text{ are the two Borels containing } T \\ &\implies X^T = \{x, nx\}, \text{ where } x := B \in G/B \end{aligned}$$

We want to show that  $X = \{x\} \sqcup Unx$ , which will imply that  $G = B \sqcup UnB$ . Note that  $x$  is  $U$ -fixed, so  $\{x\}$  and  $Unx$  are disjoint (as  $x \neq nx$ ). Also,  $Unx$  is  $T$ -stable, as

$$TUnx = UTnx = UnTx = Unx,$$

and  $Unx \neq \{nx\}$ , as otherwise

$$\{nx\} = Unx = Bnx \implies \{x\} = n^{-1}Bnx \implies n^{-1}Bn \subset \text{Stab}_G(x) = B \implies \text{contradiction}$$

Hence,  $\overline{Unx} = X$ , by dimension considerations, so  $Unx \subset X$  is open,  $X - Unx$  is finite (as  $\dim X = 1$ ), and  $X - Unx$  is  $T$ -stable.  $T$  is connected and so

$$U - Unx \subset X^T = \{x, nx\} \implies X - Unx = \{x\}$$

(ii):

$$\begin{aligned} 1 &= \dim Unx \\ &= \dim U - \dim(U \cap nUn^{-1}), \text{ as } Unx \text{ is a } U\text{-orbit} \\ &= \dim U, \text{ as } U \cap nUn^{-1} = \text{Stab}_U(nx) \text{ is finite by Theorem 157} \end{aligned}$$

Hence,

$$\begin{aligned} \dim B &= \dim T + \dim U = 1 + 1 = 2 \\ \dim G &= \dim B + \dim(G/B) = 2 + 1 = 3 \end{aligned}$$

$\mathcal{D}G$  is semisimple by Proposition 153 and is not solvable (as  $G$  is not).  $\text{rk } \mathcal{D}G \leq \text{rk } G = 1$ . If  $\text{rk } \mathcal{D}G = 0$ , then a Borel of  $\mathcal{D}G$  is unipotent, which by Proposition 125 implies that  $\mathcal{D}G$  is solvable: contradiction. (Or,  $T_1 = \{1\}$  is a maximal torus and  $T_1 = \mathcal{Z}_{\mathcal{D}G}(T_1) = \mathcal{D}G$ : contradiction.) Hence,  $\text{rk } \mathcal{D}G = 1$ , so  $\dim \mathcal{D}G = 3$  by the above:  $\mathcal{D}G = G$ .

(iii):  $\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$ . Since  $\dim \mathfrak{g} = 3$  and  $\dim \mathfrak{t} = 1$ , we have  $\#\Phi = 2$ . Moreover,  $\Phi$  is  $W$ -stable and  $[n] \in W$  acts by  $-1$  on  $X^*(T)$ , and so  $\Phi = \{\alpha, \alpha\}$  for some  $\alpha$ :  $\dim \mathfrak{g}_{\pm\alpha} = 1$ . From  $B = U \rtimes T$  we have  $\text{Lie } B = \mathfrak{t} \oplus \text{Lie } U$  and  $\text{Lie } U = \mathfrak{g}_\alpha$  or  $\mathfrak{g}_{-\alpha}$ , as  $\text{Lie } U$  is a  $T$ -stable subspace of  $\mathfrak{g}$  of dimension 1. Without loss of generality,  $\text{Lie } U = \mathfrak{g}_\alpha$ . Likewise,

$$nBn^{-1} = nUn^{-1} \rtimes T \implies \text{Lie}(nBn^{-1}) = \mathfrak{t} \oplus \text{Lie}(nUn^{-1})$$

Since  $\text{Lie}(nUn^{-1}) = \text{Ad}(n)(\text{Lie } U)$  and  $[n] \in W$  acts as  $-1$  on  $X^*(T)$ ,  $\text{Lie}(nUn^{-1}) = \mathfrak{g}_{-\alpha}$ .

(iv). This is a surjective map of homogeneous  $U \times B$  spaces.

$$\begin{aligned} unb = n &\implies u \in U \cap nBn^{-1} = U \cap nUn^{-1}, \text{ which is finite by Theorem 157} \\ &\implies U \cap nUn^{-1} = 1, \end{aligned}$$

(as  $T$ , being connected, acts trivially by conjugation  $\implies U \cap nUn^{-1} \subset \mathcal{Z}_G(T) = T$ )  
 $\implies \psi$  is injective, hence bijective

$$\begin{aligned}
d\phi \text{ bijective} &\iff d\left( U \times B \xrightarrow{(u,b) \mapsto unb^{-1}} UnBn^{-1} \right) \text{ injective} \\
&\iff d(U \times (nBn^{-1}) \xrightarrow{\text{mult.}} UnBn^{-1}) \text{ injective} \\
&\iff 0 = \text{Lie } U \cap \text{Lie}(nBn^{-1}) = \mathfrak{g}_\alpha \cap (\mathfrak{t} \oplus \mathfrak{g}_{-\alpha})
\end{aligned}$$

(v). See Springer 7.2.4. □

## 6.4 Reductive groups of semisimple rank 1.

**Lemma 173.** *If  $\phi : H \rightarrow K$  is a morphism of algebraic groups, then*

$$d\phi(\text{Ad}(h) \cdot X) = \text{Ad}(\phi(h)) \cdot d\phi X$$

*Proof.* Exercise. (Easy!) □

**Proposition 174.** *Suppose that  $\text{ss-rk } G = 1$ . Set  $\overline{G} := G/RG$  and  $\overline{T} :=$  image of  $T$  in  $\overline{G}$  ( $T$  being a maximal torus). Note that  $X^*(\overline{T}) \subset X^*(T)$  as  $T \twoheadrightarrow \overline{T}$ .*

(i) *There is  $\alpha \in X^*(\overline{T})$  such that  $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}$ , and  $\dim \mathfrak{g}_{\pm\alpha} = 1$ .*

(ii)  *$\mathcal{D}G \cong \text{SL}_2$  or  $\text{PSL}_2$*

(iii)  *$\#W = 2$ , so there are precisely two Borels containing  $T$ , and, if  $B$  is one, then*

$$\text{Lie } B = \mathfrak{t} \oplus \mathfrak{g}_{\pm\alpha} \quad \text{and} \quad \text{Lie } B_u = \mathfrak{g}_{\pm\alpha}$$

(iv) *If  $T_1$  denotes the unique maximal torus of  $\mathcal{D}G$  contained in  $T$ , then  $\exists! \alpha^\vee \in X_*(T_1) \subset X_*(T)$  such that  $\langle \alpha, \alpha^\vee \rangle = 2$ . Moreover, letting  $W = \{1, s_\alpha\}$ , we have*

$$\begin{aligned}
s_\alpha \mu &= \mu - \langle \mu, \alpha^\vee \rangle \alpha \quad \text{for all } \mu \in X^*(T) \\
s_\alpha \lambda &= \lambda - \langle \alpha, \lambda \rangle \alpha^\vee \quad \text{for all } \lambda \in X_*(T)
\end{aligned}$$

*Proof.*

(i):  $\overline{G}$  is semisimple of rank 1.

We have

$$0 \rightarrow \text{Lie } RG \rightarrow \text{Lie } G \rightarrow \text{Lie } \overline{G} \rightarrow 0$$

From Lemma 173, restricting actions, we have that the morphisms  $T \rightarrow \overline{T}$  and  $\text{Lie } G \rightarrow \text{Lie } \overline{G}$  are compatible with the action of  $T$  on  $\text{Lie } G$  and  $\overline{T}$  on  $\text{Lie } \overline{G}$ .  $T$  acts trivially on  $\text{Lie } RG$  (as  $RG \subset T$ ). Thus,

$$\Phi = \Phi(\overline{G}, \overline{T}) = \{\alpha, -\alpha\} \subset X^*(\overline{T}) \subset X^*(T)$$

and  $\dim \mathfrak{g}_{\pm\alpha} = 1$ .

(ii):  $\mathcal{D}G$  is semisimple by Proposition 153. If  $T_1 \subset \mathcal{D}G$  is a maximal torus with image  $\overline{T}_1$  in  $\overline{G}$ , then

$$\dim T_1 = \dim \overline{T}_1 + \dim(T_1 \cap RG) \leq 1$$

the inequality being due to the fact that  $T_1 \cap RG \subset \mathcal{D}G \cap RG$  is finite by Proposition 153. If  $\dim T_1 = 0$ , then the Borel of  $\mathcal{D}G$  is unipotent, implying that  $\mathcal{D}G$  is solvable, which gives that  $G$  is solvable, a contradiction. Hence,  $\text{rk } \mathcal{D}G = 1$ . By Proposition 171,  $\mathcal{D}G \cong \text{SL}_2$  or  $\text{PSL}_2$ .

(iii): First a lemma.

**Lemma 175.** *Suppose that  $\pi : G \rightarrow G'$  with  $\ker \pi$  connected and solvable. Then  $\pi(T)$  is a maximal torus of  $G'$  and we have a bijection*

$$\{\text{Borels of } G \text{ containing } T\} \xrightleftharpoons[\pi^{-1}]{\pi} \{\text{Borels of } G' \text{ containing } \pi(T)\}$$

Moreover,  $G'$  is reductive.

*Proof of lemma.* In the proposed bijection,  $\xrightarrow{\pi}$  is well-defined by Corollary 123. For the inverse, note that  $G/\pi^{-1}(B') \rightarrow G'/B'$  is bijective, which gives that  $\pi^{-1}(B')$  is parabolic as well as connected and solvable ( $\ker \pi$  and  $B'$  are connected and solvable).

$\pi^{-1}(RG')$  is a connected, solvable, normal subgroup of the torus  $RG$ .  $RG' = \pi(\pi^{-1}(RG'))$  is then a torus and so  $G'$  is reductive.  $\square$

By the Lemma,  $\#W = \#W(\overline{G}, \overline{T}) \stackrel{171}{=} 2$ . Pick a Borel  $B \supset T$ , so that  $\overline{B} \supset \overline{T}$  is a Borel.

$$1 \rightarrow RG \rightarrow B \rightarrow \overline{B} \rightarrow 1$$

being exact implies that

$$0 \rightarrow \text{Lie } RG \rightarrow \text{Lie } B \rightarrow \text{Lie } \overline{B} \rightarrow 0$$

is also exact.  $T$  again acts trivially on  $\text{Lie } RG$ .

$$\text{Lie } \overline{B} = \text{Lie } T \oplus \mathfrak{g}_{\pm\alpha} \implies \text{Lie } B = \mathfrak{t} \oplus \mathfrak{g}_{\pm\alpha}.$$

Also,

$$\text{Lie } B = \mathfrak{t} \oplus \text{Lie } B_u \implies \text{Lie } B_u = \mathfrak{g}_{\pm\alpha}$$

(iv)  $T_1$  exists, as  $\mathcal{D}G \trianglelefteq G$  (exercise). It is unique, as  $T_1 = (T \cap \mathcal{D}G)^0$ . (Another exercise:  $T_1 = T \cap \mathcal{D}G$ . Use that  $\mathcal{D}G$  is reductive.) Let  $y$  be a generator of  $X_*(T) \cong \mathbf{Z}$ . We have the containment

$$\text{Lie } \mathcal{D}G \subset \mathfrak{g} = \mathfrak{t} \oplus \mathfrak{g}_{\alpha} \oplus \mathfrak{g}_{-\alpha}$$

with  $T_1$  acting in the former and  $T$  on the latter.  $\mathcal{D}G$  being reductive implies - by Proposition 171

$$\Phi(\mathcal{D}G, T_1) = \{\pm\alpha|_{T_1}\}.$$

$\mathcal{D}G \cong \text{SL}_2$ :

$$T_1 = \left\{ \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \mid x \in k^\times \right\} \subset \text{SL}_2.$$

By the adjoint action (conjugation),  $T_1$  acts on

$$\text{Lie SL}_2 = M_2(k)_{\text{trace } 0} = k \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \oplus k \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Its roots are

$$\alpha : \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \mapsto x^2, \quad -\alpha : \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix} \mapsto x^{-2}.$$

Moreover, we can take

$$y = x \mapsto \begin{pmatrix} x & \\ & x^{-1} \end{pmatrix}$$

(or its inverse), which gives

$$\langle \alpha, y \rangle = \pm 2.$$

$\mathcal{DG} \cong \text{PSL}_2 \cong \text{GL}_2/\mathbf{G}_m$ :

$\overline{T}_1$  is equal to the image of  $D_2$  in  $\text{PSL}_2$ . By the adjoint action,  $T_1$  acts on

$$\text{Lie PSL}_2 = M_2(k)/k = k \left[ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right] \oplus k \left[ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right] \oplus k \left[ \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right].$$

Its roots are

$$\alpha : \left[ \begin{pmatrix} x_1 & \\ & x_2 \end{pmatrix} \right] \mapsto x_1 x_2^{-1}, \quad -\alpha : \left[ \begin{pmatrix} x_1 & \\ & x_2 \end{pmatrix} \right] \mapsto (x_1 x_2^{-1})^{-1} = x_1^{-1} x_2.$$

Moreover, we can take

$$y = x \mapsto \left[ \begin{pmatrix} x & \\ & 1 \end{pmatrix} \right]$$

(or its inverse), which gives

$$\langle \alpha, y \rangle = \pm 1.$$

Therefore, in any case,

$$\alpha^\vee := \frac{2y}{\langle \alpha, y \rangle} \in X_*(T_1)$$

and it is the unique cocharacter such that  $\langle \alpha, \alpha^\vee \rangle = 2$ .

If  $\lambda \in X_*(T)$ ,

$$s_\alpha \lambda - \lambda : \mathbf{G}_m \rightarrow T, \quad x \mapsto [n, \lambda(x)] = n \lambda(x) n^{-1} \lambda(x)^{-1},$$

where  $n \in N_G(T)$  is such that  $[n] = s_\alpha$ .  $s_\alpha \lambda - \lambda$  has image in  $(T \cap \mathcal{DG})^0 = T_1$ ; hence

$$s_\alpha \lambda - \lambda \in X_*(T_1) = \mathbf{Z}y.$$

Say  $s_\alpha \lambda - \lambda = \theta(\lambda)y$ . We have

$$\begin{aligned} \theta(\lambda)\langle \alpha, y \rangle &= \langle \alpha, s_\alpha \lambda - \lambda \rangle = \langle \alpha, s_\alpha \lambda \rangle - \langle \alpha, \lambda \rangle \\ &= \langle s_\alpha(\alpha), \lambda \rangle - \langle \alpha, \lambda \rangle, \quad \text{as this is true for } \overline{G} \text{ (Prop. 171), and } N_G(T)/T \cong N_{\overline{G}}(\overline{T})/\overline{T} \\ &= \langle -\alpha, \lambda \rangle - \langle \alpha, \lambda \rangle \\ &= -2\langle \alpha, \lambda \rangle \end{aligned}$$

Therefore,

$$\theta(\lambda) = \frac{-2\langle \alpha, \lambda \rangle}{\langle \alpha, y \rangle}$$

and

$$s_\alpha \lambda = \lambda + \theta(\lambda)y = \lambda - \frac{2\langle \alpha, \lambda \rangle}{\langle \alpha, y \rangle}y = \lambda - \langle \alpha, \lambda \rangle \alpha^\vee.$$

If  $\mu \in X^*(T)$ , then for all  $\lambda \in X_*(T)$

$$\langle s_\alpha \mu, \lambda \rangle = \langle \mu, s_\alpha \lambda \rangle = \langle \mu, \lambda \rangle - \langle \alpha, \lambda \rangle \langle \mu, \alpha^\vee \rangle = \langle \mu - \langle \mu, \alpha^\vee \rangle \alpha, \lambda \rangle$$

and so

$$s_\alpha \mu = \mu - \langle \mu, \alpha^\vee \rangle \alpha$$

□

**Lemma 176.**

(i) Let  $S \subset T$  be a subtorus such that  $\dim S = \dim T - 1$ . Then

$$\ker(\text{res} : X^*(T) \rightarrow X^*(S)) = \mathbf{Z}\mu$$

for some  $\mu \in X^*(T)$ .

(ii) If  $\nu \in X^*(T)$ ,  $m \in \mathbf{Z} - \{0\}$ , then  $(\ker \nu)^0 = (\ker m\nu)^0$ .

(iii) If  $\nu_1, \nu_2 \in X^*(T) - \{0\}$ , then

$$(\ker \nu_1)^0 = (\ker \nu_2)^0 \iff m\nu_1 = n\nu_2$$

for some  $m, n \in \mathbf{Z} - \{0\}$ .

*Proof.*

(i):  $\text{res}$  is surjective (exercise) and

$$X^*(T) \cong \mathbf{Z}^r, \quad X^*(S) \cong \mathbf{Z}^{r-1}.$$

(ii):

$$\underline{\text{“} \subset \text{”}}: \nu(t) = 1 \implies \nu(t)^n = 1.$$

$$\underline{\text{“} \supset \text{”}}: t \in (\ker m\nu)^0 \implies \nu(t)^n = 1, \text{ so } \nu((\ker m\nu)^0) \text{ is connected and finite, hence trivial.}$$

(iii):

$$\underline{\text{“} \Leftarrow \text{”}}: \text{Clear from (ii).}$$

$$\underline{\text{“} \Rightarrow \text{”}}: \text{Define } S = (\ker \nu_1)^0 = (\ker \nu_2)^0 \subset T, \text{ as in (i). Clearly, } \text{res}(\nu_1) = \text{res}(\nu_2) = 0, \text{ so } \nu_i \in \mathbf{Z}\mu.$$

The result follows. □

## 6.5 Root data.

**Definitions 177.** A **root datum** is a quadruple  $(X, \Phi, X^\vee, \Phi^\vee)$ , where

(i)  $X, X^\vee$  are free abelian groups of finite rank with a perfect bilinear pairing  $\langle \cdot, \cdot \rangle : X \times X^\vee \rightarrow \mathbf{Z}$

(ii)  $\Phi \subset X$  and  $\Phi^\vee \subset X^\vee$  are finite subsets with a bijection  $\Phi \rightarrow \Phi^\vee, \alpha \mapsto \alpha^\vee$

satisfying the following axioms:

(1)  $\langle \alpha, \alpha^\vee \rangle = 2$  for all  $\alpha \in \Phi$

(2)  $s_\alpha(\Phi) = \Phi$  and  $s_{\alpha^\vee}(\Phi^\vee) = \Phi^\vee$  for all  $\alpha \in \Phi$

where the “reflections” are given by

$$\begin{aligned} s_\alpha : X &\rightarrow X & s_{\alpha^\vee} : X^\vee &\rightarrow X^\vee \\ x &\mapsto x - \langle x, \alpha^\vee \rangle \alpha & y &\mapsto y - \langle \alpha, y \rangle \alpha^\vee \end{aligned}$$

A root datum is **reduced** if  $\mathbf{Q}\alpha \cap \Phi = \{\pm\alpha\}$  for all  $\alpha \in \Phi$ .

Recall that  $\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$ ,  $T_\alpha = (\ker \alpha)^0$ ,  $G_\alpha = \mathcal{Z}_G(T_\alpha)$ .

**Theorem 178.**

(i) For all  $\alpha \in \Phi$ ,  $G_\alpha$  is connected, reductive of semisimple rank 1.

- $\text{Lie } G_\alpha = \mathfrak{t} \oplus \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}$
- $\dim \mathfrak{g}_{\pm\alpha} = 1$
- $\mathbf{Q}\alpha \cap \Phi = \{\pm\alpha\}$

(ii) Let  $s_\alpha$  be the unique nontrivial element of  $W(G_\alpha, T) \subset W(G, T)$ . Then there exists  $\alpha^\vee \in X_*(T)$  such that  $\text{im } \alpha^\vee \subset \mathcal{D}G_\alpha$  and  $\langle \alpha, \alpha^\vee \rangle = 2$ . Moreover,

$$\begin{aligned} s_\alpha \mu &= \mu - \langle \mu, \alpha^\vee \rangle \alpha, & \text{for all } \mu \in X^*(T) \\ s_\alpha \lambda &= \lambda - \langle \alpha, \lambda \rangle \alpha^\vee, & \text{for all } \lambda \in X_*(T) \end{aligned}$$

(iii) Let  $\Phi^\vee = \{\alpha^\vee \mid \alpha \in \Phi\}$ . Then  $(X^*(T), \Phi, X_*(T), \Phi^\vee)$  is a reduced root datum.

(iv)  $W(G, T) = \langle s_\alpha \mid \alpha \in \Phi \rangle$

*Proof.*

(i). We saw above that  $G_\alpha$  is connected, reductive of semisimple rank 1.

$$\text{Lie } G_\alpha = \text{Lie } \mathcal{Z}_G(T_\alpha) \stackrel{98}{=} \mathfrak{z}_{\mathfrak{g}}(T_\alpha) = \mathfrak{t} \oplus \bigoplus_{\substack{\beta \in \Phi \\ \beta|_{T_\alpha}=1}} \mathfrak{g}_\beta$$

By Proposition 174,

$$\text{Lie } G_\alpha = \mathfrak{t} \oplus \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha}$$

with  $\dim \mathfrak{g}_{\pm\alpha} = 1$ . Hence, for all  $\beta \in \Phi$ ,

$$\begin{aligned} \beta|_{T_\alpha} = 1 &\iff \beta \in \{\pm\alpha\} \\ &\iff (\ker \alpha)^0 \subset (\ker \beta)^0 \\ &\iff (\ker \alpha)^0 = (\ker \beta)^0 \quad (\text{dimension considerations}) \\ &\iff \beta \in \mathbf{Q}\alpha \quad (\text{Lemma 82}) \end{aligned}$$

(ii): Follows from Proposition 174

(iii):

$\alpha \mapsto \alpha^\vee$  is bijective ( $\iff$  injective):

If  $\alpha^\vee = \beta^\vee$ , then

$$\begin{aligned} s_\alpha s_\beta(x) &= (x - \langle x, \beta^\vee \rangle \beta) - \langle (x - \langle x, \beta^\vee \rangle \beta), \alpha^\vee \rangle \alpha \\ &= x - \langle x, \alpha^\vee \rangle (\alpha + \beta) + \langle x, \alpha^\vee \rangle \langle \beta, \beta^\vee \rangle \alpha \\ &= x - \langle x, \alpha^\vee \rangle (\alpha + \beta) + 2\langle x, \alpha^\vee \rangle \alpha \\ &= x + \langle x, \alpha^\vee \rangle (\alpha - \beta) \end{aligned}$$

Therefore, if  $\langle \alpha - \beta, \alpha^\vee \rangle = 0$ , then for some  $n$

$$\begin{aligned} (s_\alpha s_\beta)^n = 1 &\implies \forall x, \quad x = (s_\alpha s_\beta)^n(x) = x + n\langle x, \alpha^\vee \rangle (\alpha - \beta) \\ &\implies \forall x, \quad 0 = n\langle x, \alpha^\vee \rangle (\alpha - \beta) \\ &\implies 0 = \alpha - \beta \\ &\implies \alpha = \beta \end{aligned}$$

$s_\alpha \Phi = \Phi$ :

The action of  $s_\alpha \in W$  on  $X^*(T)$  (and  $X_*(T)$ ) agrees with the action of  $s_\alpha$  (and  $s_{\alpha^\vee}$ ) in the definition of a root datum by (ii). Also, as noted above,  $W = N_G(T)/T$  preserves  $\Phi$ .

$s_{\alpha^\vee} \Phi^\vee = \Phi^\vee$ :

For  $w = [n] \in W$ , ( $n \in N_G(T)$ ),  $\beta \in \Phi$

$$w\beta(\cdot) = \beta(n^{-1}(\cdot)n) \implies \ker(w\beta) = n(\ker \beta)n^{-1} \implies T_{w\beta} = nT_\beta n^{-1}, G_{w\beta} = nG_\beta n^{-1}$$

From

$$\text{im}(w(\beta^\vee)) = \text{im}(n\beta^\vee n^{-1}) \subset n\mathcal{D}G_\beta n^{-1} = \mathcal{D}G_{w\beta}$$

and

$$\langle w\beta, w(\beta^\vee) \rangle = \langle \beta, \beta^\vee \rangle = 2$$

by (ii), we have that  $(w\beta)^\vee = w(\beta^\vee)$ . (iii) follows.

(iv): Induct on  $\dim G$ . Let  $w = [n] \in W$ ,  $n \in N_G(T)$ . As in the proof of Theorem 146 consider the homomorphism

$$\phi: T \rightarrow T, \quad t \mapsto [t, n] = ntn^{-1}t^{-1}.$$

$\text{im } \phi \neq T$ :

$S := (\ker \phi)^0 \neq 1$  is a torus and  $n \in \mathcal{Z}_G(S)$ . (Note that  $\mathcal{Z}_G(S)$  is connected, reductive by Corollary

158. Its roots are  $\{\alpha \in \Phi \mid \alpha|_S = 1\}$  and  $W(\mathcal{Z}_G(S), T) \subset W(G, T)$ .) If  $\mathcal{Z}_G(S) \neq G$ , we are done by induction.

If  $\mathcal{Z}_G(S) = G$ , then  $S \subset \mathcal{Z}_G$ . Define  $\overline{G} = G/S$ , which is reductive by Lemma 175, and  $\overline{T} = T/S$ , which is a maximal torus of  $\overline{G}$ . By induction, the (iv) holds for  $\overline{G}$ .

$$\Phi(G, T) = \Phi(\overline{G}, \overline{T}) \subset X^*(\overline{T}) \subset X^*(T).$$

It is an easy check that we have

$$N_G(T)/T = W(G, T) \xrightarrow{\sim} W(\overline{G}, \overline{T}) = N_{\overline{G}}(\overline{T})/\overline{T}$$

restricting to

$$W(G_\alpha, T) \xrightarrow{\sim} W(\overline{G}_\alpha, s_\alpha \mapsto s_\alpha).$$

Therefore, (iv) follows for  $\overline{G}$ .

$\text{im } \phi = T$ :  
 $\phi$  being surjective is equivalent to

$$\phi^* : X^*(T) \rightarrow X^*(T), \quad \mu \mapsto (w^{-1} - 1)\mu$$

is injective. Hence,  $w - 1 : V \rightarrow V$  is injective, thus bijective, where  $V = X^*(T) \otimes_{\mathbf{Z}} \mathbf{R}$ . Fix  $\alpha \in \Phi$ . Let  $x \in V - \{0\}$  be such that  $\alpha = (w - 1)x$ . Pick a  $W$ -invariant inner product  $(, ) : V \times V \rightarrow \mathbf{R}$  (averaging a general inner product over  $W$ ). Then

$$(x, x) = (wx, wx) = (x + \alpha, x + \alpha) = (x, x) + 2(x, \alpha) + (\alpha, \alpha) \implies 2(x, \alpha) = -(\alpha, \alpha).$$

Also,  $s_\alpha x = x + c\alpha$  (where  $c = -\langle x, \alpha^\vee \rangle \in \mathbf{Z}$ ) and, as  $s_\alpha^2 = 1$ ,

$$\begin{aligned} (x, \alpha) + c(\alpha, \alpha) &= (s_\alpha x, \alpha) = (x, s_\alpha(\alpha)) = -(x, \alpha) \implies 2(x, \alpha) = -c(\alpha, \alpha) \\ &\implies c = 1 \\ &\implies s_\alpha x = x + \alpha = wx \\ &\implies (s_\alpha w)x = x. \end{aligned}$$

Therefore, redefining  $\phi$  with  $s_\alpha w$  instead of  $w$ , we have that  $\text{im } \phi \neq T$ , and we are done by the previous case.  $\square$

**Remarks 179.**

- Let  $V$  be the subspace generated by  $\Phi$  in  $X \otimes \mathbf{R}$  (for  $X$  in a root datum). Then  $\Phi$  is a root system in  $V$ . (See §14.7 in Borel's *Linear Algebraic Groups*; references are there.) If  $V = X \otimes \mathbf{R}$  (which, in fact, is equivalent to  $G$  being semisimple), then  $(X, \Phi)$  uniquely determines  $(X, \Phi, X^\vee, \Phi^\vee)$ .
- The root datum of Theorem 178 does not depend (up to isomorphism) on the choice of  $T$ , as any two maximal tori are conjugate.

Facts:

1. Isomorphism Theorem: Two connected reductive groups are isomorphic  $\iff$  their root data are isomorphic.

2. Existence Theorem: Given a reduced root datum, there exists a reductive group that has the root datum.

(See Springer §9, §10.)

**Theorem 180.**

- (i) For all  $\alpha \in \Phi$  there is a unique connected closed  $T$ -stable unipotent subgroup  $U_\alpha \subset G$  such that  $\text{Lie } U_\alpha = \mathfrak{g}_\alpha$ . There exists an isomorphism  $u_\alpha : \mathbf{G}_a \xrightarrow{\sim} U_\alpha$  (unique up to scalar) such that

$$tu_\alpha(x)t^{-1} = u_\alpha(\alpha(t)x) \quad \text{for all } x \in \mathbf{G}_a, t \in T.$$

- (ii)  $G = \langle T, U_\alpha (\alpha \in \Phi) \rangle$  (i.e.,  $G$  is the smallest subgroup containing  $T$  and all of the  $U_\alpha$ )

- (iii)  $Z_G = \bigcap_{\alpha \in \Phi} \ker \alpha$

*Proof.*

(i): Let  $B_\alpha$  denote the Borel subgroup of  $G_\alpha$  containing  $T$  with  $\text{Lie } B_\alpha = \mathfrak{t} \oplus \mathfrak{g}_\alpha$  (Proposition 174, Theorem 178.) Then  $U_\alpha := (B_\alpha)_u$  satisfies all assumptions by Proposition 174. Also,  $\dim U_\alpha = \dim \mathfrak{g}_\alpha = 1$  and  $U_\alpha \cong \mathbf{G}_a$  by Theorem 58. Let  $u_\alpha : \mathbf{G}_a \rightarrow U_\alpha$  denote any isomorphism; any other differs by a scalar as  $\text{Aut } \mathbf{G}_a \cong \mathbf{G}_m$ . So  $tu_\alpha(x)t^{-1} = u_\alpha(\chi(t)x)$  for some  $\chi(t) \in k^\times$ . Via  $u_\alpha$ , identify  $U_\alpha \xrightarrow{t(\cdot)t^{-1}} U_\alpha$  with  $\mathbf{G}_a \xrightarrow{\chi(t)} \mathbf{G}_a$ . Since the derivative of the former is  $\mathfrak{g}_\alpha \xrightarrow{\text{Ad}(t)=\alpha(t)} \mathfrak{g}_\alpha$ , we see that the derivative of the latter is  $k \xrightarrow{\alpha(t)} k$ . However, by Theorem 76, we must have  $\alpha(t) = \chi(t)$  - and thus  $\alpha = \chi$ .

It remain to show that  $U_\alpha$  is unique. If  $U'_\alpha$  is another connected, closed,  $T$ -stable, and unipotent with  $\text{Lie } U'_\alpha = \mathfrak{g}_\alpha$ , by the same argument as above we get an isomorphism  $u'_\alpha : \mathbf{G}_a \rightarrow U'_\alpha$  such that  $tu'_\alpha(x)t^{-1} = u'_\alpha(\alpha(t)x)$ . Hence,  $U'_\alpha \subset G_\alpha$  (as  $\alpha(T_\alpha) = 1$ ).

$$\begin{aligned} T \text{ normalises } U'_\alpha &\implies TU'_\alpha \text{ is closed, connected, and solvable (exercise)} \\ &\implies TU'_\alpha \text{ is contained in a Borel containing } T \\ &\implies TU'_\alpha \subset B_\alpha, \quad \text{as } \text{Lie } U'_\alpha = \mathfrak{g}_\alpha \\ &\implies U'_\alpha = (TU'_\alpha)_u \subset (B_\alpha)_u = U_\alpha \\ &\implies U'_\alpha = U_\alpha \text{ (dimension)} \end{aligned}$$

- (ii): By Corollary 21,  $\langle T, U_\alpha (\alpha \in \Phi) \rangle$  is connected, closed. Its Lie algebra contains  $\mathfrak{t}$  and all of the  $\mathfrak{g}_\alpha$ , hence coincides with  $\mathfrak{g}$ . Thus

$$\dim \langle T, U_\alpha (\alpha \in \Phi) \rangle = \dim \mathfrak{g} = \dim G \implies \langle T, U_\alpha (\alpha \in \Phi) \rangle = G$$

- (iii):  $Z_G \subset T$  by Corollary 158 By (i),  $t \in T$  commutes with  $U_\alpha \iff \alpha(t) = 1$ , which implies that  $Z_G \subset \bigcap_{\alpha} \ker \alpha$ . The reverse inclusion follows by (ii).  $\square$

# Appendix. An example: the symplectic group

Set  $G = \mathrm{Sp}_{2n} = \{g \in \mathrm{GL}_{2n} \mid g^t J g = J\}$ , where  $J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$ .

**Fact.**  $G$  is connected. (See, for example, Springer 2.2.9 (1) or Borel 23.3.)

Define

$$\begin{aligned} T &= G \cap D_{2n} = \{\mathrm{diag}(x_1, \dots, x_{2n}) \mid \mathrm{diag}(x_1, \dots, x_{2n}) \cdot \mathrm{diag}(x_{n+1}, \dots, x_{2n}, x_1, \dots, x_n) = I\} \\ &= \{\mathrm{diag}(x_1, \dots, x_n, x_1^{-1}, \dots, x_n^{-1})\} \\ &\cong \mathbf{G}_m^n \end{aligned}$$

Clearly  $Z_G(T) = T$ , implying that  $T$  is a maximal torus and  $\mathrm{rk} G = n$ . Write  $\epsilon_i$ ,  $1 \leq i \leq n$ , for the morphisms

$$T \rightarrow \mathbf{G}_m, \quad \mathrm{diag}(x_1, \dots, x_n^{-1}) \mapsto x_i,$$

which form a basis of  $X^*(T)$ .

**Lemma 181.** *If  $\rho : G \rightarrow \mathrm{GL}(V)$  is a faithful (i.e., injective)  $G$ -representation that is semisimple, then  $G$  is reductive.*

*Proof.*

$U := R_u G$  is a connected, unipotent, normal subgroup of  $G$ . Write  $V = \bigoplus_{i=1}^r V_i$  with  $V_i$  irreducible ( $V$  is semisimple).  $V_i^U \neq 0$ , as  $U$  is unipotent (Proposition 39), and  $V_i^U \subset V_i$ , is  $G$ -stable, as  $U \trianglelefteq G$ :  $V_i^U = V_i$ . Hence,  $U$  acts trivially on  $V$ , and is thus trivial, since  $\rho$  is injective.  $\square$

We will show that the natural faithful representation  $G \hookrightarrow \mathrm{GL}_{2n}$  is irreducible and hence  $G$  is reductive. Let  $V = k^{2n}$  denote the underlying vector space with standard basis  $(e_i)_{i=1}^{2n}$ . We have  $V = \bigoplus_{i=1}^{2n} k e_i$  and, for all  $t \in T$ ,

$$t e_i = \begin{cases} \epsilon_i(t) e_i, & i \leq n \\ \epsilon_{i-n}(t)^{-1} e_i, & i > n \end{cases}$$

Any  $G$ -subrepresentation of  $V$  is a direct sum of  $T$ -eigenspaces; hence, it is enough to show that  $N_G(T)$  acts transitively on the  $k e_i$ , which is equivalent to it acting transitively on  $\{\pm \epsilon_1, \dots, \pm \epsilon_n\} \subset X^*(T)$ .

A calculation shows that the elements

$$g_i := \mathrm{diag}(I_{i-1}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, I_{n-2}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, I_{n-i-1}), \quad (1 \leq i < n)$$

lie in  $G$ , where  $\text{diag}(A_1, A_2, \dots)$  denotes a matrix with square blocks  $A_1, A_2, \dots$  along the diagonal in the given order. As well

$$g_n := \begin{pmatrix} \text{diag}(I_{n-1}, 0) & E_{nn} \\ -E_{nn} & \text{diag}(I_{n-1}, 0) \end{pmatrix},$$

lies in  $G$ , where  $E_{nn} \in M_n(k)$  has a 1 in the  $(n, n)$ -entry and 0's elsewhere. Note that the  $g_i \in N_G(T)$  for all  $i$  and  $g_i : \epsilon_i \mapsto \epsilon_{i+1}$ , for  $1 \leq i < n$ , and  $g_n : \epsilon_n \mapsto -\epsilon_n$  (with  $g_i \cdot \epsilon_j = \epsilon_j$  for  $i \neq j$ ). Hence,  $N_G(T)$  does act transitively on  $\{\pm\epsilon_i\}$ , so  $V$  is irreducible and  $G$  is reductive.

Lie Algebra:

If  $\psi : \text{GL}_{2n} \rightarrow \text{GL}_{2n}$ ,  $g \mapsto g^t J g$ , then  $d\psi_1 : M_{2n}(k) \rightarrow M_{2n}(k)$ ,  $X \mapsto X^t J + J X$  (as in the proofs of Propositions 77 and 78). Hence,

$$\mathfrak{g} \subset \{X \in M_{2n}(k) \mid X^t J + J X = 0\} =: \mathfrak{g}'.$$

Checking that  $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathfrak{g}'$  if and only if  $B^t = B, C^t = C$ , and  $D = -A^t$ , we see that

$$\dim \mathfrak{g}' = n^2 + 2 \binom{n+1}{2} = n(2n+1)$$

*Claim:*  $\dim G \geq n(2n+1)$

Define

$$\phi : \text{GL}_{2n} \rightarrow \mathbf{A}^{\binom{2n}{2}}, \quad g \mapsto ((g^t J g)_{ij})_{i < j}.$$

We have  $\phi^{-1}((J_{ij})_{i < j}) = G$ , (because  $g^t J g$  is antisymmetric). (This is still okay when  $p = 2$ .) So,

$$(2n)^2 = \dim \text{GL}_{2n} \stackrel{85}{=} \dim \overline{\phi(\text{GL}_{2n})} + \text{minimal fibre dimension} \leq \binom{2n}{2} + \dim G$$

and

$$\dim G \geq (2n)^2 - \binom{2n}{2} = n(2n+1).$$

Hence,

$$\dim \mathfrak{g} \leq n(2n+1) \leq \dim G = \dim \mathfrak{g} \implies \dim \mathfrak{g} = n(2n+1)$$

and so

$$\dim G = n(2n+1), \quad \text{and} \quad \mathfrak{g} = \{X \in M_{2n}(k) \mid X^t J + J X = 0\}.$$

Roots:

Write  $E_{ij}$  for the  $(2n) \times (2n)$  matrix with a 1 in the  $(i, j)$ -entry and 0's elsewhere. By the above,

$$\mathfrak{g} = \mathfrak{t} \oplus \left( \bigoplus_{i \neq j} k \begin{pmatrix} E_{ij} & 0 \\ 0 & -E_{ji} \end{pmatrix} \right) \oplus \left( \bigoplus_{i \leq j} k \begin{pmatrix} 0 & E_{ij} + E_{ji} \\ 0 & 0 \end{pmatrix} \right) \oplus \left( \bigoplus_{i \leq j} k \begin{pmatrix} 0 & 0 \\ E_{ij} + E_{ji} & 0 \end{pmatrix} \right)$$

(with  $E_{ij} + E_{ji}$  in the latter factors replaced with  $E_{ii}$  if  $i = j$  and  $p = 2$ ). Correspondingly,

$$\Phi = \{\epsilon_i - \epsilon_j \mid i \neq j\} \cup \{\epsilon_i + \epsilon_j \mid i \leq j\} \cup \{-\epsilon_i - \epsilon_j \mid i \leq j\}$$

( A check:  $\#\Phi = n(n-1) + \binom{n+1}{2} + \binom{n+1}{2} = 2n^2 = \dim \mathfrak{g} - \dim \mathfrak{t}$ .)

Coroots:

Let  $\epsilon_1^*, \dots, \epsilon_n^*$  denote the dual basis, so

$$\epsilon_i^*(x) = \text{diag}(1, \dots, x, \dots, x^{-1}, \dots, 1) = \text{diag}(I_{i-1}, x, I_{n-1}, x^{-1}, I_{n-i}).$$

We have

$$G_{\epsilon_i - \epsilon_j} = G \cap (D_{2n} + kE_{ij} + kE_{ji} + kE_{n+i, n+j} + kE_{n+j, n+i})$$

and so  $G_{\epsilon_i - \epsilon_j}$  is contained in

$$G \cap \{I_{2n} + (a-1)E_{ii} + bE_{ij} + cE_{ji} + (d-1)E_{jj} + (a'-1)E_{n+i, n+i} + b'E_{n+i, n+j} + c'E_{n+j, n+i} + (d'-1)E_{n+j, n+j}\}$$

where  $a, b, c, d, a', b', c', d'$  are such that  $ad - bc = 1 = a'd' - b'c'$ . It follows that

$$(\epsilon_i - \epsilon_j)^\vee = \epsilon_i^* - \epsilon_j^*.$$

Similarly,  $(\epsilon_i + \epsilon_j)^\vee = \epsilon_i^* + \epsilon_j^*$  and  $(-\epsilon_i - \epsilon_j)^\vee = -\epsilon_i^* - \epsilon_j^*$ .

$G$  is semisimple.  $RG = \mathcal{Z}_G^0 = \left( \bigcap_{\Phi} \ker \alpha \right)^0 = 1$